

# COUPLING RATE-INDEPENDENT AND RATE-DEPENDENT PROCESSES: EXISTENCE RESULTS

RICCARDA ROSSI AND MARITA THOMAS

ABSTRACT. We address the analysis of an abstract system coupling a rate-independent process with a rate-dependent nonlinear evolution equation. We propose suitable weak solution concepts and obtain existence results by passing to the limit in carefully devised time-discretization schemes. Our arguments combine techniques from the theory of *gradient systems* with the toolbox for *rate-independent* evolution, thus reflecting the mixed character of the problem. Finally, we discuss applications to a class of rate-independent processes in visco-elastic solids with inertia, and to a recently proposed model for damage with plasticity.

**Key words:** rate-independent processes; gradient systems; inertia; semistable energetic solutions; existence results; time discretization; generalized standard solids.

## 1. INTRODUCTION

The modeling of dissipative processes in mechanical systems via internal variables often leads to PDE systems which have the following structure

$$D_u \mathcal{E}(t, u(t), z(t)) = 0 \quad \text{in } \mathbf{V}^* \quad \text{for a.a. } t \in (0, T), \quad (1.1a)$$

$$\partial \mathcal{R}(z'(t)) + D_z \mathcal{E}(t, u(t), z(t)) = 0 \quad \text{in } \mathbf{Z}^* \quad \text{for a.a. } t \in (0, T). \quad (1.1b)$$

Here, the (separable) Banach spaces  $\mathbf{V}$  and  $\mathbf{Z}$  are the state spaces for the (slow) variable  $u$  and the (fast) internal variable  $z$ , with  $z'$  its time derivative. A guiding example is the deformation of a body under the influence of dissipative processes such as, e.g., damage, plasticity, phase transformations, or delamination. Assuming small strains, the deformation is described by its displacement field  $u$ , whereas, within the theory of generalized standard materials [HN75] (see also [Fré02a]), the changes of the elastic behavior due to the evolving dissipative processes are modeled by an *internal variable*  $z$ . The evolution of the system then results from a trade-off between the two competing mechanisms of energy conservation and energy dissipation, caused by time-dependent external loadings. Here and in what follows, the latter are included in the time-dependent energy functional  $\mathcal{E} : [0, T] \times \mathbf{V} \times \mathbf{Z} \rightarrow (-\infty, +\infty]$ , ( $D_u \mathcal{E}$  and  $D_z \mathcal{E}$  denoting its Gâteaux derivatives with respect to the variables  $u$  and  $z$ ); dissipation in  $z$  is evaluated through a positive, lower semicontinuous, and convex dissipation potential  $\mathcal{R} : \mathbf{Z} \rightarrow [0, +\infty]$ , which we assume in addition to be positively 1-homogeneous. Thus system (1.1) is invariant under rescalings of the time variable, i.e. it is *rate-independent*. We refer to [Mie05, Mie11, MR15] for surveys on the analysis of such class of systems.

In this paper we want to analyze the case in which the variable  $u$  additionally evolves subject to *viscous* dissipation (for example, according to Kelvin-Voigt rheology in the frame of material modeling). On top of that we also allow for the presence of inertia. Thus, (1.1a) is replaced by the evolution equation

$$\varrho u''(t) + \partial \mathcal{V}(u'(t)) + D_u \mathcal{E}(t, u(t), z(t)) \ni 0 \quad \text{in } \mathbf{V}^* \quad \text{for a.a. } t \in (0, T). \quad (1.1c)$$

Within this introduction, for expository reasons we will take the inertial term to be modulated by a constant  $\varrho \geq 0$ . The (lower semicontinuous, convex) dissipation potential  $\mathcal{V} : \mathbf{V} \rightarrow [0, +\infty)$  has superlinear growth at infinity, namely  $\lim_{\|v\|_{\mathbf{V}} \uparrow +\infty} \frac{\mathcal{V}(v)}{\|v\|_{\mathbf{V}}} = +\infty$ . The kinetic energy leading to the inertial term  $\varrho u''$  in (1.1c) is of the form

$$\mathcal{K}(u') = \frac{\varrho}{2} \|u'\|_{\mathbf{W}}^2 \quad \text{with } \mathbf{W} \text{ a Hilbert space such that } \mathbf{V} \subset \mathbf{W} \text{ continuously and densely} \quad (1.2)$$

and  $\mathbf{W} = \mathbf{W}^*$ , i.e.  $\mathbf{W}$  is identified with its dual  $\mathbf{W}^*$ . Observe that, for  $\varrho = 0$  system (1.1b, 1.1c) falls into the class of (*generalized*) *gradient systems*; in what follows, we will refer to it with the symbol  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ . In the case with inertia (i.e., if  $\varrho > 0$ ), (1.1b, 1.1c) contains a *reversible* term and thus ceases to be a *pure*

gradient system. We shall refer to it as a *damped inertial system* and denote it by  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ . In this paper, we will address both gradient systems and damped inertial systems.

Despite its *mixed* rate-independent/rate-dependent character, (1.1b, 1.1c) retains all the difficulties attached to the analysis of purely rate-independent systems. In particular, due to the linear growth at infinity of the potential  $\mathcal{R}$ , for the variable  $z$  as a function of time we only have BV-regularity, which does not guarantee the existence of the pointwise derivative  $z'(t)$  and thus calls for weak solvability concepts for (1.1b, 1.1c).

This issue was first addressed in [Rou09], focusing on a class of processes with a mixed rate-independent/rate-dependent character in generalized standard materials, modeled by a system of the type (1.1b, 1.1c), with a *quadratic* dissipation potential  $\mathcal{V}$ . In the spirit of the concept of *energetic solutions* for rate-independent systems, cf. [Mie05], an *energetic-type* weak solvability concept was proposed and analyzed in [Rou09]. This has provided a sound mathematical basis for various subsequent extensions, in particular to systems coupling rate-independent, rate-dependent, and *thermal* processes in thermo-visco-elastic materials, cf. [Rou10]. It has also paved the way to the recent [RT15], where a broad class of models encompassing rate-independent and rate-dependent processes coupled with temperature evolution and phase separation has been tackled.

In this paper, though, we will confine ourselves to the isothermal setting. We aim to revisit the analysis carried out in [Rou09] from a more abstract and general perspective. This will allow us to extend the existence results in [Rou09] to a much broader class of dissipation potentials  $\mathcal{V}$  and energy functionals  $\mathcal{E}$ . In particular, let us mention that the energies considered in [Rou09] need not be differentiable w.r.t. the variable  $z$ , as the energetic-type solution notion therein used does not involve the Gâteaux differential  $D_z \mathcal{E}$ , but they are taken to be sufficiently smooth w.r.t.  $u$ , and with suitable convexity properties. Here, instead, we focus on the case in which the functional  $u \mapsto \mathcal{E}(t, u, z)$  may be *nonconvex and nonsmooth*. Accordingly, we will replace the Gâteaux differential  $D_u \mathcal{E}$  in (1.1c) by the *Fréchet subdifferential* of  $\mathcal{E}$  with respect to  $u$ , namely the multivalued operator  $\partial_u^- \mathcal{E} : [0, T] \times \mathbf{V} \times \mathbf{Z} \rightrightarrows \mathbf{V}^*$  defined at a point  $(t, u, z)$  in the domain of  $\mathcal{E}$  by

$$\xi \in \partial_u^- \mathcal{E}(t, u, z) \quad \text{if and only if} \quad \liminf_{v \rightarrow u \text{ in } \mathbf{V}} \frac{\mathcal{E}(t, v, z) - \mathcal{E}(t, u, z) - \langle \xi, v - u \rangle_{\mathbf{V}}}{\|v - u\|_{\mathbf{V}}} \geq 0. \quad (1.3)$$

Indeed,  $\partial_u^- \mathcal{E}$  may be understood as a localization of the subdifferential of  $\mathcal{E}$  (w.r.t.  $u$ ) in the sense of convex analysis, and in fact it coincides with the latter object as soon as the functional  $u \mapsto \mathcal{E}(t, u, z)$  is *convex*, whereas if  $u \mapsto \mathcal{E}(t, u, z)$  is Gâteaux-differentiable, then  $\partial_u^- \mathcal{E}(t, u, z)$  reduces to the singleton  $\{D_u \mathcal{E}(t, u, z)\}$ . Therefore, we will study the subdifferential inclusion

$$\rho u''(t) + \partial \mathcal{V}(u'(t)) + \partial_u^- \mathcal{E}(t, u(t), z(t)) \ni 0 \quad \text{in } \mathbf{V}^* \quad \text{for a.a. } t \in (0, T), \quad (1.4a)$$

coupled with the (formally written) inclusion

$$\partial \mathcal{R}(z'(t)) + D_z \mathcal{E}(t, u(t), z(t)) \ni 0 \quad \text{in } \mathbf{Z}^* \quad \text{for a.a. } t \in (0, T). \quad (1.4b)$$

**1.1. Solution concepts for system (1.4).** As previously mentioned, already in the pure rate-independent context the lack of time regularity of  $z$  due to the rate-independent character of the flow rule (1.4b) makes it necessary to resort to a weak solvability concept. The one we are going to adopt in this coupled rate-independent/rate-dependent framework is the natural generalization of the energetic notion proposed for the specific class of systems in [Rou09], involving a *quadratic* dissipation  $\mathcal{V}$ .

In order to motivate our notion(s) of solution, let us momentarily continue to argue with the formally written system (1.4), observing that it can be rewritten as

$$\begin{aligned} -(\rho u''(t) + \xi(t)) &\in \partial \mathcal{V}(u'(t)) && \text{with } \xi(t) \in \partial_u^- \mathcal{E}(t, u(t), z(t)) && \text{in } \mathbf{V}^* && \text{for a.a. } t \in (0, T), \\ -D_z \mathcal{E}(t, u(t), z(t)) &\in \partial \mathcal{R}(z'(t)) && && \text{in } \mathbf{Z}^* && \text{for a.a. } t \in (0, T). \end{aligned} \quad (1.5)$$

Recall that, for a given convex and lower semicontinuous functional  $\Phi$  one has  $\zeta \in \partial \Phi(w)$  if and only if there holds  $\Phi(w) + \Phi^*(\zeta) = \langle \zeta, w \rangle$ , where the Fenchel-Moreau conjugate is defined by  $\Phi^*(\zeta) := \sup_w \{\langle \zeta, w \rangle - \Phi(w)\}$ . Therefore, introducing the conjugates  $\mathcal{V}^*$  and  $\mathcal{R}^*$ , (1.5) rewrites as

$$\begin{aligned} \mathcal{V}(u'(t)) + \mathcal{V}^*(-(\rho u''(t) + \xi(t))) &= -\langle (\rho u''(t) + \xi(t)), u'(t) \rangle_{\mathbf{V}} && \text{for a.a. } t \in (0, T), \\ \mathcal{R}(z'(t)) + \mathcal{R}^*(-D_z \mathcal{E}(t, u(t), z(t))) &= -\langle D_z \mathcal{E}(t, u(t), z(t)), z'(t) \rangle_{\mathbf{Z}} && \text{for a.a. } t \in (0, T). \end{aligned} \quad (1.6)$$

Now, taking into account the definition and properties of  $\mathcal{V}^*$  and  $\mathcal{R}^*$ , it is not difficult to see that the two separate identities in (1.6) imply

$$\begin{aligned} \mathcal{V}(u'(t)) + \mathcal{V}^*(-\varrho u''(t) + \xi(t)) + \mathcal{R}(z'(t)) &= -\langle (\varrho u''(t) + \xi(t)), u'(t) \rangle_{\mathbf{V}} - \langle D_z \mathcal{E}(t, u(t), z(t)), z'(t) \rangle_{\mathbf{Z}} \\ &= -\frac{d}{dt} \mathcal{E}(t, u(t), z(t)) + \partial_t \mathcal{E}(t, u(t), z(t)) - \mathcal{K}(u'(t)) \end{aligned} \quad (1.7a)$$

for a.a.  $t \in (0, T)$ , obtained by adding them up, using the chain rules for  $\mathcal{E}$  and for the kinetic energy  $\mathcal{K}$ , and the fact that, by the 1-homogeneity of  $\mathcal{R}$ , its conjugate  $\mathcal{R}^*$  is the indicator functional of the set  $K^* = \partial \mathcal{R}(0) = \{\eta \in \mathbf{Z}^* : \langle \eta, \zeta \rangle_{\mathbf{Z}} \leq \mathcal{R}(\zeta) \text{ for all } \zeta\}$ . Therefore, one has in fact to add to (1.7a) the constraint

$$-D_z \mathcal{E}(t, u(t), z(t)) \in \partial \mathcal{R}(0) \quad \text{in } \mathbf{Z}^* \quad \text{for every } t \in [0, T], \quad (1.7b)$$

which yields  $\mathcal{R}^*(-D_z \mathcal{E}(t, u(t), z(t))) = 0$ . In turn, (1.7a) & (1.7b) yield (1.6): in fact, rearranging terms in (1.7a), in view of (1.7b) one arrives at

$$\begin{aligned} \mathcal{V}(u'(t)) + \mathcal{V}^*(-\varrho u''(t) + \xi(t)) - \langle -(\varrho u''(t) + \xi(t)), u'(t) \rangle_{\mathbf{V}} \\ = -\mathcal{R}(z'(t)) - \mathcal{R}^*(-D_z \mathcal{E}(t, u(t), z(t))) + \langle -D_z \mathcal{E}(t, u(t), z(t)), z'(t) \rangle_{\mathbf{Z}}, \end{aligned}$$

whence the separate identities in (1.6).

We are now in the position to state our weak solvability notion for system (1.4): We call a pair  $(u, z) : [0, T] \rightarrow \mathbf{V} \times \mathbf{Z}$  a *semistable energetic solution* to the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ , resp. damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$  for  $\varrho > 0$ , if it fulfills

- (1) the subdifferential inclusion (1.4a);
- (2) the *semistability condition*

$$\mathcal{E}(t, u(t), z(t)) \leq \mathcal{E}(t, u(t), \tilde{z}) + \mathcal{R}(\tilde{z} - z(t)) \quad \text{for all } \tilde{z} \in \mathbf{Z} \text{ for all } t \in [0, T]; \quad (1.8a)$$

- (3) the *energy-dissipation inequality*

$$\begin{aligned} \frac{\varrho}{2} \|u'(t)\|_{\mathbf{W}}^2 + \int_0^t \mathcal{V}(u'(s)) + \mathcal{V}^*(-\xi(s) - \varrho u''(s)) ds + \text{Var}_{\mathcal{R}}(z; [0, t]) + \mathcal{E}(t, u(t), z(t)) \\ \leq \frac{\varrho}{2} \|u'(0)\|_{\mathbf{W}}^2 + \mathcal{E}(0, u(0), z(0)) + \int_0^t \partial_t \mathcal{E}(s, u(s), z(s)) ds \quad \text{for all } t \in [0, T], \end{aligned} \quad (1.8b)$$

with  $\xi(s)$  a selection in  $\partial_u^- \mathcal{E}(s, u(s), z(s))$  fulfilling (1.4a), i.e.  $\varrho u''(s) + \omega(s) + \xi(s) = 0$  with  $\omega(s) \in \partial \mathcal{V}(u'(s))$ , for almost all  $s \in (0, T)$ .

First of all, let us remark that due to  $\mathcal{V}(u'(t)) + \mathcal{V}^*(-\xi(s) - \varrho u''(s)) = \langle \partial \mathcal{V}(u'(t)), u'(t) \rangle_{\mathbf{V}}$  by convexity and in view of (1.4a), in the case of a quadratic dissipation potential  $\mathcal{V}$  the semistable energetic solutions in the sense of (1.8) coincide with the solution concept introduced in [Rou09, Def. 5.1] and there simply called *energetic solutions*. Clearly, the energy-dissipation inequality (1.8b) is the integrated, inequality version of (1.7a), whereas the semistability (1.8a) can be understood as a weak form of condition (1.7b). This solution concept well reflects the *mixed* character of system (1.4). On the one hand, in the spirit of the energetic formulation of (purely) rate-independent systems, it features an energy (in-)equality and a (semi)stability condition in which the rate-dependent character of the variable  $u$  is tracked by the fact that only the variable  $z$  is allowed to vary, whereas  $u$  is kept fixed. On the other hand, (1.8b) is reminiscent of the energy identity

$$\int_0^t (\mathcal{V}(u'(s)) + \mathcal{V}^*(-D_u \mathcal{E}(s, u(s)))) ds + \mathcal{E}(t, u(t)) = \mathcal{E}(0, u(0)) + \int_0^t \partial_t \mathcal{E}(s, u(s)) ds. \quad (1.9)$$

In fact, the origins of (1.9) (also referred to as *De Giorgi principle*, cf. [Mie16]) can be traced back to the seminal work by E. DE GIORGI on gradient flows in metric spaces, cf. [DGMT80, Amb95, AGS08]. We recall that the validity of (1.9) is *equivalent*, under suitable conditions on the energy functional  $\mathcal{E}$  (not depending on the variable  $z$  and Gâteaux-differentiable w.r.t.  $u$  for the sake of simplicity), to the fact that  $u$  is a solution to the (*pure*) gradient system

$$\partial \mathcal{V}(u'(t)) + D_u \mathcal{E}(t, u(t)) \ni 0 \quad \text{in } \mathbf{V}^* \quad \text{for a.a. } t \in (0, T), \quad (1.10)$$

cf. [RMS08, MRS13b].

Let us remark that the De Giorgi principle has also been exploited for the Vlasov-Fokker-Planck equation in [DPZ13], which also combines conservative with dissipative terms as it is the case for the damped inertial systems. Also there, as in (1.8b), the corresponding energy-dissipation balance features the thermodynamic

driving force  $-\xi(s) - \varrho u''(s)$  in the argument of  $\mathcal{V}^*$ . However, while [DPZ13] deduce their corresponding (1.8b) as an equality, we only require (1.8b) to hold as an inequality. On the one hand, this is the minimal requirement for suitable a priori estimates. On the other hand, as discussed in Section 3, under better regularity conditions (cf. Prop. 3.2, Prop. 3.5, Thm. 3.7), inequality (1.8b) can be upgraded to an energy-dissipation *equality*, so that the first law of thermodynamics is valid.

We will also work with a weaker solution notion, referred to as *weak semistable energetic* solution, where the validity of the subdifferential inclusion (1.4a) is no longer claimed and thus only the semistability (1.8a) and the energy-dissipation inequality (1.8b) (with  $\xi(s)$  a selection of  $\partial_u^- \mathcal{E}(s, u(s), z(s))$  for a.e.  $s \in (0, T)$ ) are required to hold, similarly as for energetic solutions to (purely) rate-independent system. Due to its intrinsically *variational* character, this concept will turn out to be particularly flexible for the *Evolutionary  $\Gamma$ -convergence analysis* for gradient and damped inertial systems in the forthcoming [RT16], in the same spirit as in [MRS08, Mie16].

The argument guaranteeing the passage from *weak semistable energetic* to *semistable energetic* solutions is drawn from the variational theory for gradient systems in [MRS13b]. Therein it was proved that, if the energy satisfies a suitable chain-rule inequality, then any curve fulfilling (1.9) a priori only in terms of an *inequality* “ $\leq$ ”, indeed complies with the energy *equality* (1.9), and it is thus a solution to (1.10). Similarly, in the forthcoming Proposition 3.2 we are going to show that any weak semistable energetic solution along which the (integral) chain-rule inequality

$$\begin{aligned} & \int_0^t \langle D_u \mathcal{E}(s, u(s), z(s)), u'(s) \rangle_{\mathbf{V}} ds \\ & \leq \mathcal{E}(t, u(t), z(t)) - \mathcal{E}(0, u(0), z(0)) - \int_0^t \partial_t \mathcal{E}(s, u(s), z(s)) ds + \text{Var}_{\mathcal{R}}(z; [0, t]) \end{aligned} \quad (1.11)$$

holds, is in fact a semistable energetic solution, thus fulfilling the subdifferential inclusion (1.4a), and such that (1.8b) holds as an *energy-dissipation balance*. Note that the integral chain rule inequality (1.11) here has been stated for simplicity in the case  $u \mapsto \mathcal{E}(t, u, z)$  is smooth. In Theorem 3.7 we will also provide some sufficient conditions on  $\mathcal{E}$  guaranteeing the validity of (1.11) along weak semistable energetic solutions. The proof of the latter result is in turn based on the combination of the semistability condition (1.8a) with Riemann sum techniques, typical of the theory of rate-independent systems, cf. [DMFT05, Mie05, MM05].

**1.2. Existence results.** To obtain existence results, we will follow an approach combining ideas for gradient flows and gradient systems, cf., e.g., [Amb95, AGS08, RS06, RMS08, MRS13b], with techniques for rate-independent evolution drawn from [MT04, Mie05, Mie11, MRS08]: Namely, we will enucleate a series of *abstract conditions* on the energy  $\mathcal{E}$  and on the dissipation potentials  $\mathcal{V}$  and  $\mathcal{R}$ , guaranteeing the existence (via limit passage in carefully devised time-discretization schemes) of:

**Theorem 4.9:** semistable energetic solutions to  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  for  $\mathcal{V}$  *quadratic*;

**Theorem 4.13:** weak semistable energetic solutions to  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  for *general*  $\mathcal{V}$  with superlinear growth;

**Theorem 5.4:** weak semistable energetic solutions to  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$  for *general*  $\mathcal{V}$  with superlinear growth;

**Theorem 5.6:** semistable energetic solutions to  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$  for  $\mathcal{V}$  *quadratic*.

Let us stress that Theorems 4.9 & 4.13 are concerned with pure gradient systems, i.e.  $\varrho = 0$  in (1.4a), while Theorems 5.4 & 5.6 treat the case with inertia,  $\varrho > 0$  in (1.4a). While postponing to Sections 2, 4, and 5 a detailed analysis of all of our hypotheses, let us briefly highlight here their most significant features, focusing on the case of the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ .

First of all, as standard for gradient flows, we require coercivity, i.e. that there exists  $\tau_o > 0$  such that for every  $(u_o, z_o) \in \mathbf{U} \times \mathbf{X}$

$$\text{the map } (u, z) \mapsto \mathcal{E}(t, u, z) + \tau_o \mathcal{V} \left( \frac{u - u_o}{\tau_o} \right) + \mathcal{R}(z - z_o) \text{ has sublevels bounded in } \mathbf{U} \times \mathbf{X}, \quad (1.12)$$

where the spaces  $\mathbf{U} \subset \mathbf{V}$  and  $\mathbf{X} \subset \mathbf{Z}$  with continuous embeddings. In fact, (1.12) is the minimal coercivity requirement to ensure that the *alternating (decoupled)* time-incremental minimization scheme

$$\begin{aligned} u_\tau^n &\in \underset{u \in \mathbf{V}}{\operatorname{Argmin}} \left( \tau \mathcal{V} \left( \frac{u - u_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, u, z_\tau^{n-1}) \right), \\ z_\tau^n &\in \underset{z \in \mathbf{Z}}{\operatorname{Argmin}} \left( \tau \mathcal{R} \left( \frac{z - z_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, u_\tau^n, z) \right) \end{aligned} \quad (1.13)$$

(cf. Remark 7.6 ahead for some comments on (1.13)), yielding discrete solutions to (1.4) in the case of a *general* dissipation potential  $\mathcal{V}$  with superlinear growth, does admit solutions. One then constructs approximate solutions to (1.4) by interpolating them suitably.

A crucial step in the existence proof is the derivation of a *discrete energy-dissipation* inequality satisfied by the approximate solutions: all a priori estimates stem from it, and one has to pass to the limit in this discrete inequality to obtain the time-continuous (1.8b). Again focusing on the case of gradient systems, we will exploit two methods of proof to obtain the discrete version of (1.8b).

- (1) For the case of a general dissipation potential, we will resort to specific techniques from the theory of *Minimizing Movements* for gradient flows, employing the notion of *variational interpolant* of the discrete solutions due to E. DE GIORGI, cf. [Amb95, AGS08].
- (2) For  $\mathcal{V}$  quadratic, we will make use of a suitable condition on the Fréchet subdifferential  $\partial_u^- \mathcal{E}$ , referred to as uniform Fréchet subdifferentiability, namely

$$\begin{aligned} \exists \Lambda \geq 0 \quad \forall t \in [0, T], \forall u, v \in \mathbf{V}, \forall z \in \mathbf{Z} \quad \forall \xi \in \partial_u^- \mathcal{E}(t, u, z) : \\ \mathcal{E}(t, v, z) - \mathcal{E}(t, u, z) \geq \langle \xi, v - u \rangle_{\mathbf{V}} - \Lambda \|v - u\|_{\mathbf{V}}^2. \end{aligned} \quad (1.14)$$

Essentially, (1.14) gives a *global* character to inequality (2.11) defining the Fréchet subdifferential. As we will see, it is guaranteed if, e.g.,  $u \mapsto \mathcal{E}(t, u, z)$  fulfills some  $\lambda$ -convexity property.

A key ingredient for passing to the limit in the *discrete energy-dissipation* inequality, and in the (discrete version) of the subdifferential inclusion (1.4a), will be a suitable closedness/continuity condition on  $(\partial_u^- \mathcal{E}, \mathcal{E})$ . It ensures that the graph of the Fréchet subdifferential  $\partial_u^- \mathcal{E}$  is closed in the strong-weak topology of  $\mathbf{V} \times \mathbf{V}^*$  along sequences with bounded energy, and in some qualified situations, leading to *enhanced* (weak) semistable energetic solutions, also that the energy functional is itself continuous along such sequences. In the case of general dissipation potentials  $\mathcal{V}$ , we will have to impose this requirement additionally. Nonetheless, we will see that closedness/continuity can be in fact derived as a consequence of the uniform Fréchet subdifferentiability condition (1.14) (which we will impose for quadratic  $\mathcal{V}$ ), in combination with a further *recovery sequence condition* on  $\mathcal{E}$ .

Finally, in order to prove the semistability (1.8a) we will exploit a version of the mutual recovery sequence condition proposed in [MRS08] for the fully rate-independent setting. This condition allows us to pass to the limit in the discrete form of the semistability condition by requiring that, given a sequence  $(u_n, z_n)_n \subset \mathbf{V} \times \mathbf{Z}$  semistable at times  $t_n$  and such that  $(t_n, u_n, z_n) \rightarrow (t, u, z)$  in suitable topologies, for every  $\tilde{z} \in \mathbf{Z}$  there exists a *recovery sequence*  $(\tilde{z}_n)_n \subset \mathbf{Z}$  converging to  $\tilde{z}$  and such that

$$\limsup_{n \rightarrow \infty} (\mathcal{R}(\tilde{z}_n - z_n) + \mathcal{E}(t_n, u_n, \tilde{z}_n) - \mathcal{E}(t_n, u_n, z_n)) \leq \mathcal{R}(\tilde{z} - z) + \mathcal{E}(t, u, \tilde{z}) - \mathcal{E}(t, u, z). \quad (1.15)$$

In this way, semistability (i.e. positivity of the r.h.s. in (1.15) for *any*  $\tilde{z} \in \mathbf{Z}$ ) is preserved in the limit passage.

**1.3. Applications.** To illustrate the outcome of our existence analysis for system (1.4), we have focused on applications to

- (1) the class of rate-independent processes in visco-elastic solids tackled in [Rou09], revisiting the existence results proved therein and extending them to systems modeling damage, with a *unidirectional* evolution;
- (2) a coupled damage-plasticity model recently proposed in [AMV14], whose analysis falls outside the scope of the results in [Rou09].

More precisely, in Section 6.1 we will consider systems pertaining to the class (1.4) and modeling the rate-independent evolution of an internal variable  $z$ , describing the elastic behavior of a body also subject to viscosity and inertia, at small strains. Thus, the displacement  $\mathbf{u}$  of the body will play the role of the rate-dependent variable  $u$ . Following [Rou09], we will confine the discussion to a *quadratic* dissipation potential on the strain rate  $\varepsilon(\mathbf{u}')$  and instead address quite a broad family of energy functionals of the form  $\mathcal{E}(t, \mathbf{u}, z) :=$

$\int_{\Omega} \varphi(\varepsilon(\mathbf{u}), z, \nabla z) - \mathbf{f}(t) \cdot \mathbf{u} dx$ , with  $\mathbf{f}$  a given external force. In [Rou09], the author proposed several classes of conditions on the energy density  $\varphi$  ensuring the existence of a semistable energetic solution to system (1.4). In Sec. 6.1 we will show that all the energies considered in [Rou09] fulfill the conditions of our abstract Thm. 4.9 (in the case where inertial effects are neglected), and Thm. 5.6, and thus we will recover the existence results from [Rou09]. In Sec. 6.2 we will further discuss an application to a model for adhesive contact between visco-elastic bodies subject to inertia. This seems to indicate that our abstract approach has a *unifying* and ultimately *simplifying* character.

It also paves the way to extensions of the analysis developed in [Rou09], tackled in Sec. 6.3. In particular, therein we will deduce the existence of *weak semistable energetic solutions* for rate-independent, *unidirectional*, damage processes in viscous solids driven by *non-quadratic* dissipation potential acting on  $\varepsilon(\mathbf{u}')$ , and by energies encompassing a BV, instead of a Sobolev, regularizing gradient term in the damage variable, and even featuring a non-smooth time-dependence.

Finally, in Sec. 6.4 we will show the *flexibility* of our approach by tackling a different type of mixed system, where both the rate-independent and the rate-dependent variables are internal variables. More specifically, we will consider a (purely) elastic body subject to damage, encompassing in the model the development of plasticity. While for the plastic tensor we will standardly assume a rate-independent evolution, we will take the dissipation potential acting on the damage rate to be quadratic. The displacement will be governed by the static balance of elastic energy. The resulting PDE system will be cast in the form (1.4) by introducing the *reduced energy* obtained by minimizing out the displacements. Then, with careful calculations we will check that Theorem 4.9 applies, yielding the existence of *semistable energetic solutions*.

**Plan of the paper.** In [Section 2](#) we set up the abstract functional analytic setting for (1.4) and fix the *basic conditions* on the energy  $\mathcal{E}$  and the dissipation potentials  $\mathcal{V}$  and  $\mathcal{R}$  that will be adopted throughout the paper. [Section 3](#) focuses on the notions of *semistable energetic* and *weak semistable energetic* solutions to (1.4), on their respective properties and mutual relations; finally, in Sec. 3.3 we show that, under uniform convexity of the energy the temporal regularity of the rate-independent variable  $z$  improves, thus extending to this mixed framework some of the results well known in the rate-independent context.

In [Section 4](#) we present our existence results, Theorems 4.9 & 4.13, for the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ , for quadratic  $\mathcal{V}$  and general  $\mathcal{V}$  with superlinear growth, respectively. We thoroughly discuss all of our abstract conditions on  $\mathcal{E}$ ,  $\mathcal{V}$ ,  $\mathcal{R}$ , and finally illustrate them on some examples in Sec. 4.3. The existence Thms. 5.4 and 5.6 for the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ , for  $\mathcal{V}$  general and  $\mathcal{V}$  quadratic, resp., are stated in [Section 5](#), where we also develop some comparison with the existence and approximation results by E. EMMRICH for abstract second order nonlinear evolution equations. [Section 6](#) is devoted to the applications.

The proofs of Theorems 4.9, 4.13, 5.4 & 5.6 are given throughout [Section 7](#), also relying on some tools from the theory of Young measures with values in Banach spaces, shortly recapped in [Appendix A](#).

In order to make navigation easier in the paper, we have summarized the four existence results with their crucial assumptions, examples and applications in Table 1.3 below.

## 2. SETUP

In the following, we will denote by  $\|\cdot\|_X$  the norm of a Banach space  $X$ , and by  $\langle \cdot, \cdot \rangle_X$  the duality pairing between  $X^*$  and  $X$ . If  $X$  is a Hilbert space, its inner product shall be denoted by  $(\cdot, \cdot)_X$ . With the symbol  $\mathbf{B}([0, T]; X)$  we will denote the space of bounded, measurable, *everywhere defined* functions on  $[0, T]$  with values in  $X$ .

**Topological setup.** The conditions on the map  $u \mapsto \mathcal{E}(t, u, z)$  will involve two

$$\text{(separable) reflexive Banach spaces } \mathbf{U} \subset \mathbf{V} \text{ with a continuous and dense embedding.} \quad (2.1a)$$

Furthermore, in the case inertial effects are included in system (1.4), there will come into play

$$\text{a Hilbert space } \mathbf{W}, \text{ identified with its dual } \mathbf{W}^*, \text{ such that } \begin{cases} \mathbf{U} \Subset \mathbf{W} & \text{compactly,} \\ \mathbf{V} \subset \mathbf{W} & \text{continuously and densely.} \end{cases} \quad (2.1b)$$

<b>Gradient system</b>	$\mathcal{V}$	Notion of $\partial_u^- \mathcal{E}$	Closedness of $\partial_u^- \mathcal{E}$	(EDI)	+Prop. 3.2/3.5/Thm. 3.7
Theorem 4.9 ( $\exists \star$ ) $\Downarrow$ Applications: Secs. 6.1, 6.3 (Thm. 6.3), 6.4	quadratic	(4.6) in Hyp. 4.2	(1) $\begin{cases} \mathbf{U} \Subset \mathbf{V}, \\ \text{Hyp. 4.6} \end{cases}$ $\Downarrow$ (4.12) in L. 4.7 ----- (2) $\begin{cases} \mathbf{U} \subset \mathbf{V}, \\ \text{Hyp. 4.8} \end{cases}$ $\begin{cases} (4.13a) \\ (4.13b) \end{cases}$	$\Rightarrow$ “ $\leq$ ” enhanced ----- $\Rightarrow$ “ $\leq$ ” enhanced ----- $\Rightarrow$ “ $\leq$ ” enhanced	$\Rightarrow$ “ $\equiv$ ” enhanced ----- $\Rightarrow$ “ $\equiv$ ” enhanced ----- $\Rightarrow$ “ $\equiv$ ” enhanced
Examples		Ex. 4.5, 4.15	for (1): Ex. 4.18 for (2): Ex. 4.16, 4.17		
Theorem 4.13 ( $\exists$ weak $\star$ ) $\Downarrow$ Application: Sec. 6.3 (Thm. 6.5)	superlinear	Def. (1.3)	(3) $\begin{cases} \mathbf{U} \subset \mathbf{V}, \\ \text{Hyp. 4.12} \end{cases}$ $\begin{cases} (4.24a) \\ (4.24b) \end{cases}$	$\Rightarrow$ “ $\leq$ ” ----- $\Rightarrow$ “ $\leq$ ” enhanced	$\Rightarrow$ “ $\equiv$ ” enhanced ----- $\Rightarrow$ “ $\equiv$ ” enhanced
Examples			for (3): Ex. 4.16		
<b>Damped inertial system</b>	$\mathcal{V}$	Notion of $\partial_u^- \mathcal{E}$	Closedness of $\partial_u^- \mathcal{E}$	(EDI)	+Prop. 3.2/3.5/Thm. 3.7
Theorem 5.4 ( $\exists$ weak $\star$ ) $\Downarrow$ Application: Sec. 6.3 (Thm. 6.7)	superlinear	(5.7) in Hyp. 5.2 satisfying (5.8) in Hyp. 5.3	(1) or (3)	$\Rightarrow$ “ $\leq$ ” ----- $\Rightarrow$ “ $\leq$ ” enhanced	$\Rightarrow$ “ $\equiv$ ” ----- $\Rightarrow$ “ $\equiv$ ” enhanced
Examples		(5.7): Ex. 4.5, 4.15; (5.8): Ex. 5.10	(1): Ex. 4.18; (3): Ex. 4.16		
Theorem 5.6 ( $\exists \star$ ) $\Downarrow$ Applications: Secs. 6.1, 6.2, 6.3 (Thm. 6.9)	quadratic $\mathbf{V} \Subset \mathbf{W}$	(5.7) in Hyp. 5.2 satisfying (5.8) in Hyp. 5.3	(2) $\begin{cases} \mathbf{U} \subset \mathbf{V}, \\ \text{Hyp. 4.8} \end{cases}$ $\begin{cases} (4.13a) \\ (4.13b) \end{cases}$	$\Rightarrow$ “ $\leq$ ” ----- $\Rightarrow$ “ $\leq$ ” enhanced	$\Rightarrow$ “ $\equiv$ ” enhanced ----- $\Rightarrow$ “ $\equiv$ ” enhanced
Examples		(5.7): Ex. 4.5, 4.15; (5.8): Ex. 5.10	Ex. 4.16, 4.17, 5.10		

TABLE 1. Existence results, main assumptions, applications, and examples: (weak)  $\star =$  (weak) *semistable energetic solution*, cf. Def. 3.1 (Def. 3.4); (EDI) = *energy dissipation inequality*, cf. (1.8b). Column 5 indicates whether the respective closedness condition in column 4 *enhances* (EDI) “ $\leq$ ” in the sense of Def. 3.1 & Def. 3.4, and, in combination with Prop. 3.2/3.5/Thm. 3.7, even to an enhanced *balance* “ $\equiv$ ”, cf. column 6.

Hence,

$$\mathbf{U} \subset \mathbf{V} \subset \mathbf{W} = \mathbf{W}^* \subset \mathbf{V}^* \subset \mathbf{U}^* \quad \text{with continuous and dense embeddings} \quad (2.2)$$

and

$$\langle w, u \rangle_{\mathbf{V}} = (w, u)_{\mathbf{W}} \quad \text{for all } u \in \mathbf{V} \text{ and } w \in \mathbf{W}. \quad (2.3)$$

We will then consider the *kinetic energy*

$$\mathcal{K}(v) := \frac{1}{2} \|v\|_{\mathbf{W}}^2 \quad \text{for all } v \in \mathbf{W}. \quad (2.4)$$

The assumptions on  $z \mapsto \mathcal{E}(t, u, z)$  will feature two Banach spaces

$$\mathbf{X} \subset \mathbf{Z} \text{ continuously, with } \mathbf{X} \text{ dual of a separable Banach space,} \quad (2.5a)$$

and  $\mathbf{X}$  endowed with a topology  $\sigma_{\mathbf{X}}$  such that

$$\begin{aligned} &\text{bounded sets in } \mathbf{X} \text{ are } \sigma_{\mathbf{X}}\text{-sequentially compact, and} \\ &z_n \xrightarrow{\sigma_{\mathbf{X}}} z \text{ in } \mathbf{X} \Rightarrow z_n \rightharpoonup z \text{ in } \mathbf{Z}. \end{aligned} \quad (2.5b)$$

**Remark 2.1.** Let us highlight that (2.5) encompasses the two following situations:

- (1) the space  $\mathbf{X}$  is reflexive: in this case, we take  $\sigma_{\mathbf{X}}$  to be the weak topology on  $\mathbf{X}$ ;
- (2) the space  $\mathbf{X}$  embeds *compactly* in  $\mathbf{Z}$ : in this case,  $\sigma_{\mathbf{X}}$  is the strong topology on  $\mathbf{Z}$ .

Our basic conditions on the functional  $\mathcal{E}$  mimic the conditions on energy  $\mathcal{E} = \mathcal{E}_t(u)$ , independent of  $z$ , proposed in [MRS13b].

**Hypothesis 2.2** (Basic conditions on  $\mathcal{E}$ ). *The functional  $\mathcal{E} : [0, T] \times \mathbf{V} \times \mathbf{Z} \rightarrow (-\infty, +\infty]$  has proper domain  $\text{dom}(\mathcal{E}) = [0, T] \times D_u \times D_z$ , with  $D_u \subset \mathbf{U}$  and  $D_z \subset \mathbf{X}$ . Moreover,*

$$\forall t \in [0, T] \quad \text{the map } (u, z) \mapsto \mathcal{E}(t, u, z) \quad \text{is weakly lower semicontinuous on } \mathbf{V} \times \mathbf{Z}, \quad (2.6a)$$

and  $\mathcal{E}$  is bounded from below, i.e.

$$\exists C_0 > 0 \quad \forall (t, u, z) \in [0, T] \times D_u \times D_z : \quad \mathcal{E}(t, u, z) \geq C_0; \quad (2.6b)$$

in fact, since  $\mathcal{E}$  is bounded from below, up to a shift we may always assume that it is bounded by a strictly positive constant. Furthermore, we require that

$$\begin{aligned} &\forall (u, z) \in D_u \times D_z \quad \text{the map } t \mapsto \mathcal{E}(t, u, z) \text{ is differentiable, with derivative } \partial_t \mathcal{E}(t, u, z) \text{ s.t.} \\ &\exists C_1, C_2 > 0 \quad \forall (t, u, z) \in [0, T] \times D_u \times D_z : \quad |\partial_t \mathcal{E}(t, u, z)| \leq C_1 (\mathcal{E}(t, u, z) + C_2) \text{ and fulfilling} \\ &\text{for all sequences } t_n \rightarrow t, u_n \rightarrow u \text{ in } \mathbf{V}, z_n \rightharpoonup z \text{ in } \mathbf{Z} \text{ with } \sup_n \mathcal{E}(t_n, u_n, z_n) \leq C \\ &\text{there holds } \limsup_{n \rightarrow \infty} \partial_t \mathcal{E}(t_n, u_n, z_n) \leq \partial_t \mathcal{E}(t, u, z). \end{aligned} \quad (2.6c)$$

Let us set

$$\mathcal{G}(u, z) := \sup_{t \in [0, T]} \mathcal{E}(t, u, z) \quad \text{for all } (u, z) \in D_u \times D_z \text{ and} \quad (2.7)$$

$$\mathcal{S}_E := \{(u, z) \in D_u \times D_z : \mathcal{G}(u, z) \leq E\} \quad \text{for } E > 0. \quad (2.8)$$

As a consequence of the power estimate in (2.6c) and of Gronwall's inequality there holds

$$\forall s, t \in [0, T], \forall (u, z) \in D_u \times D_z : \quad \mathcal{E}(t, u, z) + C_2 \leq \exp(C_1|t-s|) (\mathcal{E}(s, u, z) + C_2), \quad (2.9)$$

so that we in particular have the following estimate, which will play a crucial role in our analysis

$$\exists K_1, K_2 > 0 \quad \forall (t, u, z) \in [0, T] \times D_u \times D_z : \quad \mathcal{G}(u, z) \leq K_1 \mathcal{E}(t, u, z) + K_2. \quad (2.10)$$

Let us also recall that the Fréchet subdifferential  $\partial_u^- \mathcal{E} : [0, T] \times \mathbf{V} \times \mathbf{Z} \rightrightarrows \mathbf{V}^*$  of  $\mathcal{E}$  w.r.t.  $u$  is defined at  $(t, u, z) \in [0, T] \times D_u \times D_z$  by

$$\xi \in \partial_u^- \mathcal{E}(t, u, z) \Leftrightarrow \mathcal{E}(t, v, z) - \mathcal{E}(t, u, z) \geq \langle \xi, v - u \rangle_{\mathbf{V}} + o(\|v - u\|_{\mathbf{V}}) \text{ as } v \rightarrow u \text{ in } \mathbf{V}. \quad (2.11)$$

It is not difficult to check that for every  $(t, u, z) \in [0, T] \times D_u \times D_z$  the set  $\partial_u^- \mathcal{E}(t, u, z)$  is closed and convex.

For the dissipation potentials  $\mathcal{V}$  and  $\mathcal{R}$  we require:



**Hypothesis 2.3** (Basic conditions on  $\mathcal{V}$  and  $\mathcal{R}$ ). *The functional  $\mathcal{V} : \mathbf{V} \rightarrow [0, +\infty)$  is lower semicontinuous, convex, and fulfills*

$$\text{dom}(\mathcal{V}) = \mathbf{V}, \quad \mathcal{V}(0) = 0, \quad \lim_{\|v\|_{\mathbf{V}} \uparrow +\infty} \frac{\mathcal{V}(v)}{\|v\|_{\mathbf{V}}} = +\infty. \quad (2.12a)$$

*The functional  $\mathcal{R} : \mathbf{Z} \rightarrow [0, +\infty]$ , with domain  $\text{dom}(\mathcal{R})$ , is lower semicontinuous, convex, 1-positively homogeneous, i.e.*

$$\mathcal{R}(\lambda\zeta) = \lambda\mathcal{R}(\zeta) \text{ for all } \zeta \in \mathbf{Z} \text{ and } \lambda > 0, \quad \text{and } \mathcal{R}(\lambda\zeta) = 0 \text{ for all } \zeta \in \mathbf{Z} \text{ and } \lambda = 0, \quad (2.12b)$$

*and coercive*

$$\exists C_R > 0 \quad \forall \zeta \in \mathbf{Z} \quad \mathcal{R}(\zeta) \geq C_R \|\zeta\|_{\mathbf{Z}}. \quad (2.12c)$$

Recall that the lower semicontinuity of a functional together with convexity results in its *weak* lower semicontinuity. Hereafter, we will denote by  $\mathcal{V}^* : \mathbf{V}^* \rightarrow [0, +\infty)$  the convex conjugate of  $\mathcal{V}$ , defined by

$$\mathcal{V}^*(\xi) := \sup_{v \in \mathbf{V}} (\langle \xi, v \rangle_{\mathbf{V}} - \mathcal{V}(v)). \quad (2.13)$$

It follows from the fact that  $\text{dom}(\mathcal{V}) = \mathbf{V}$  that  $\mathcal{V}$  is continuous and that  $\mathcal{V}^*$  as well has superlinear growth at infinity (cf. e.g. [ET74, Chap. 1, Cor. 2.5]), namely

$$\lim_{\|\xi\|_{\mathbf{V}^*} \uparrow +\infty} \frac{\mathcal{V}^*(\xi)}{\|\xi\|_{\mathbf{V}^*}} = +\infty. \quad (2.14)$$

We will denote by  $\text{Var}_{\mathcal{R}}$  the total variation induced by  $\mathcal{R}$ . Given a curve  $z : [0, T] \rightarrow \mathbf{Z}$  and a subinterval  $[s, t] \subset [0, T]$ , we set

$$\text{Var}_{\mathcal{R}}(z; [s, t]) := \sup \left\{ \sum_{j=1}^N \mathcal{R}(z(r_j) - z(r_{j-1})) : s = r_0 < r_1 < \dots < r_{N-1} < r_N = t \right\}. \quad (2.15)$$

Due to (2.12c), if  $\text{Var}_{\mathcal{R}}(z; [0, T]) < +\infty$  then  $z \in \text{BV}([0, T]; \mathbf{Z})$  and the  $\mathcal{R}$ -total variation estimates the total variation induced by  $\|\cdot\|_{\mathbf{Z}}$ , viz.

$$\text{Var}_{\mathcal{R}}(z; [0, T]) \geq C_R \text{Var}(z; [0, T]).$$

**Remark 2.4** (State-dependent dissipation potentials). Most of the forthcoming results, and in particular the existence Theorems 4.9, 4.13, 5.4, 5.6, could be extended to the case that the dissipation potential  $\mathcal{V}$  depends on the state variable  $q = (u, z)$ . More specifically, along the lines of [MRS13b] we could consider a family  $(\mathcal{V}_q)_{q \in \mathbb{D}_u \times \mathbb{D}_z}$  of dissipation potentials such that

- (1) for every  $q \in \mathbb{D}_u \times \mathbb{D}_z$  the functional  $\mathcal{V}_q : \mathbf{V} \rightarrow [0, +\infty)$  is convex and lower semicontinuous;
- (2) the dissipation potentials  $(\mathcal{V}_q)_{q \in \mathbb{D}_u \times \mathbb{D}_z}$  and  $(\mathcal{V}_q^*)_{q \in \mathbb{D}_u \times \mathbb{D}_z}$  have superlinear growth at infinity, *uniformly* w.r.t.  $q$  in sublevels of the energy, i.e.

$$\forall S > 0 : \quad \begin{cases} \lim_{\|v\|_{\mathbf{V}} \rightarrow +\infty} \inf_{\mathcal{G}(q) \leq S} \mathcal{V}_q(v) = +\infty, \\ \lim_{\|\xi\|_{\mathbf{V}^*} \rightarrow +\infty} \inf_{\mathcal{G}(q) \leq S} \mathcal{V}_q^*(\xi) = +\infty; \end{cases} \quad (2.16a)$$

- (3) the dependence  $q \mapsto \mathcal{V}_q$  is continuous, on sublevels of the energy, in the sense of MOSCO-convergence (cf. e.g. [Att84]), i.e.

$$\begin{aligned} \forall S > 0 : \quad q_n \rightharpoonup q \text{ in } \mathbf{V} \times \mathbf{Z}, \quad \mathcal{G}(q_n) \leq S, \quad v_n \rightharpoonup v \text{ in } \mathbf{V} &\Rightarrow \liminf_{n \rightarrow \infty} \mathcal{V}_{q_n}(v_n) \geq \mathcal{V}_q(v), \\ \forall S > 0 : \quad q_n \rightharpoonup q \text{ in } \mathbf{V} \times \mathbf{Z}, \quad \mathcal{G}(q_n) \leq S, \quad v \in \mathbf{V} &\Rightarrow \begin{cases} \exists v_n \rightarrow v \text{ in } \mathbf{V}, \\ \lim_{n \rightarrow \infty} \mathcal{V}_{q_n}(v_n) = \mathcal{V}_q(v). \end{cases} \end{aligned} \quad (2.16b)$$

The extension of our results to a family  $(\mathcal{R}_q)_q$  of 1-positively homogeneous potentials seems to be more delicate, starting from the fact that it would not be completely clear how to define the total variation induced by  $(\mathcal{R}_q)_q$ . That is why, we choose to omit this issue in the fully general setting, whereas in Sec. 6.4 we will briefly describe how one of our existence results, Thm. 4.9, could be adapted to encompass a very specific dependence of  $\mathcal{R}$  on the *sole* variable  $u$ .

For the limit passage in (approximate versions) of the semistability condition, a key role will be played by the following recovery sequence condition, originally introduced in [MRS08], which is also termed *mutual* because it involves the energy  $\mathcal{E}$  and the dissipation  $\mathcal{R}$ .

**Hypothesis 2.5** (Mutual Recovery Sequence condition). *Let  $(t_n, u_n, z_n)_n \subset [0, T] \times D_u \times D_z$  fulfill for every  $n \in \mathbb{N}$  the semistability condition (1.8a), and suppose that  $t_n \rightarrow t$ ,  $(u_n, z_n) \rightharpoonup (u, z)$  in  $\mathbf{V} \times \mathbf{Z}$  with  $\sup_{n \in \mathbb{N}} \mathcal{G}(u_n, z_n) \leq C$ .*

*Then for every  $\tilde{z} \in \mathbf{Z}$  there exists  $\tilde{z}_n \rightarrow \tilde{z}$  in  $\mathbf{Z}$  such that*

$$\lim_{n \rightarrow \infty} (\mathcal{R}(\tilde{z}_n - z_n) + \mathcal{E}(t_n, u_n, \tilde{z}_n) - \mathcal{E}(t_n, u_n, z_n)) \leq \mathcal{R}(\tilde{z} - z) + \mathcal{E}(t, u, \tilde{z}) - \mathcal{E}(t, u, z). \quad (2.17)$$

**Remark 2.6.** In certain cases (cf. Remark 4.10), it will be sufficient to require a weaker variant of Hypothesis 2.5, in which we suppose that for every  $\tilde{z} \in \mathbf{Z}$  it is possible to construct a recovery sequence  $(\tilde{z}_n)_n$  fulfilling (2.17), whenever the sequence  $(t_n, u_n, z_n)_n \subset [0, T] \times D_u \times D_z$  fulfills the conditions of Hypothesis 2.5 and, in addition, we have the *energy convergence*

$$\mathcal{E}(t_n, u_n, z_n) \rightarrow \mathcal{E}(t, u, z). \quad (2.18)$$

We close this section stating the main coercivity assumption on the map  $\mathcal{E}(t, \cdot, \cdot)$ . Observe that, in the spirit of the *variational approach* to gradient flows (cf. [AGS08, Chap. 2, Sec. 2.1]), we require a property on the sublevels of the functional  $\mathcal{G}$  added with  $\mathcal{V}$  and  $\mathcal{R}$ . This coercivity/compactness requirement is at the core both of the existence of solutions to the time-incremental minimization scheme(s) we shall use to discretize (1.4), and of the compactness for the related family of approximate solutions to (1.4).

**Hypothesis 2.7** (Coercivity). *There exist  $\tau_o > 0$  such that for all  $u_o \in \mathbf{V}$  and  $z_o \in \mathbf{Z}$*

$$\text{the map } (u, z) \mapsto \mathcal{G}(u, z) + \tau_o \mathcal{V}\left(\frac{u - u_o}{\tau_o}\right) + \mathcal{R}(z - z_o) \text{ has sublevels bounded in } \mathbf{U} \times \mathbf{X}. \quad (2.19)$$

Observe that (2.19) guarantees the separate coercivity properties

$$\forall \bar{z} \in D_z \quad \text{the map } u \mapsto \mathcal{G}(u, \bar{z}) + \tau_o \mathcal{V}\left(\frac{u - u_o}{\tau_o}\right) \text{ has sublevels bounded in } \mathbf{U}, \quad (2.20a)$$

$$\forall \bar{u} \in D_u \quad \text{the map } z \mapsto \mathcal{G}(\bar{u}, z) + \mathcal{R}(z - z_o) \text{ has sublevels bounded in } \mathbf{X}. \quad (2.20b)$$

As a consequence of (2.20b) and of (2.5), a sequence  $(z_n)_n \subset \mathbf{Z}$  such that  $\sup_{n \in \mathbb{N}} \mathcal{G}(\bar{u}, \bar{z}_n) + \mathcal{R}(z_n - z_o) \leq C$  for some  $C > 0$  and  $\bar{u} \in D_u$  admits a subsequence converging with respect to the  $\sigma_{\mathbf{X}}$ -topology. Instead, with (2.20a) we are requiring that  $\mathcal{E}(t, \cdot, z)$  (added up with  $\mathcal{V}$ ) has bounded sublevels in  $\mathbf{U}$ , which may or may not be compactly embedded in  $\mathbf{V}$ . In other words, we encompass in our analysis these two cases:

- the energy  $\mathcal{E}(t, \cdot, z)$  and the dissipation potential  $\mathcal{V}$  have sublevels bounded in the same space;
- the energy  $\mathcal{E}(t, \cdot, z)$  has sublevels compact in the space  $\mathbf{V}$  of the dissipation  $\mathcal{V}$ .

The following examples are prototypes of these two situations. The different choices for the space  $\mathbf{V}$  reflect the different character of the equation ruling the evolution of the “viscous” variable  $u$ . To make this more transparent, we drop the dependence on the variable  $z$ .

**Example 2.8** (Linear visco-elasticity). *Let  $\Omega \subset \mathbb{R}^d$  be a bounded Lipschitz domain. We take  $\mathbf{V} = H_0^1(\Omega; \mathbb{R}^d)$ , the dissipation potential  $\mathcal{V}(\mathbf{v}) := \int_{\Omega} \frac{1}{2} \varepsilon(\mathbf{v}) : \mathbb{D} : \varepsilon(\mathbf{v}) \, dx$  with  $\mathbb{D} \in \mathbb{R}^{d \times d \times d \times d}$  positive definite and symmetric ( $\varepsilon(\mathbf{v}) = \frac{1}{2}(\nabla \mathbf{v} + \nabla \mathbf{v}^\top) \in \mathbb{R}^{d \times d}$  denoting the linearized strain tensor), and the energy functional  $\mathcal{E} : [0, T] \times \mathbf{V} \rightarrow \mathbb{R}$  defined by*

$$\mathcal{E}(t, \mathbf{u}) := \int_{\Omega} \frac{1}{2} \varepsilon(\mathbf{u}) : \mathbb{C} : \varepsilon(\mathbf{u}) \, dx - \int_{\Omega} \mathbf{f}(t) \cdot \mathbf{u} \, dx,$$

*with  $\mathbb{C} \in \mathbb{R}^{d \times d \times d \times d}$  positive definite and symmetric, and  $\mathbf{f} \in C^1([0, T]; H^{-1}(\Omega; \mathbb{R}^d))$  a given external loading. In this case,  $\mathcal{E}$  is coercive in the space  $\mathbf{U} = \mathbf{V} = H_0^1(\Omega; \mathbb{R}^d)$ . Choosing  $\mathbf{W} = L^2(\Omega; \mathbb{R}^d)$ , the viscous equation (1.4a) yields the classical linear visco-elasticity equation with inertia*

$$\ddot{\mathbf{u}} - \operatorname{div}(\mathbb{D} : \varepsilon(\dot{\mathbf{u}}) + \mathbb{C} : \varepsilon(\mathbf{u})) = \mathbf{f} \quad \text{in } \Omega \times (0, T),$$

*with homogeneous Dirichlet boundary conditions on  $\mathbf{u}$ .*

**Example 2.9** ( $L^2$ -gradient flow). We take  $\mathbf{V} = L^2(\Omega)$ , the dissipation potential

$$\mathcal{V}(v) := \int_{\Omega} \frac{1}{2}|v|^2 dx \quad \text{for every } v \in L^2(\Omega),$$

and the energy functional  $\mathcal{E} : [0, T] \times \mathbf{V} \rightarrow (-\infty, +\infty]$  defined by

$$\mathcal{E}(t, u) := \begin{cases} \int_{\Omega} (\frac{1}{2}|\nabla u|^2 + W(u)) dx - \int_{\Omega} f(t)u dx & \text{if } u \in H^1(\Omega), \\ +\infty & \text{otherwise,} \end{cases} \quad (2.21)$$

with  $W \in C^1(\mathbb{R})$  possibly nonconvex, and fulfilling the growth condition (cf. also Example 4.16)  $-c_1 \leq W(u) \leq c_2|u|^q + c_3$  for positive constants  $c_1, \dots, c_3$ , with the exponent  $q$  such that  $2(q-1) \leq 2^* = \frac{2d}{d-2}$  (to fix ideas, if  $d = 3$  we find the condition  $q \leq 4$ , e.g. fulfilled by the double-well potential  $W(u) = \frac{1}{4}(u^2 - 1)^2$ ). In this case,  $\mathcal{E}$  is coercive in the space

$$\mathbf{U} = H^1(\Omega) \Subset \mathbf{V} = L^2(\Omega),$$

and, in the case without inertia, (1.4a) reduces to the gradient flow

$$\dot{u} - \Delta u + W'(u) = f \quad \text{in } \Omega \times (0, T).$$

### 3. SOLUTION CONCEPTS AND THEIR PROPERTIES

In Secs. 3.1 and 3.2 we give the two solvability notions for system (1.4) that will be studied throughout the paper and thoroughly examine their properties, as well as their relation. Finally, in Sec. 3.3 we show that, under suitable uniform convexity properties of the map  $z \mapsto \mathcal{E}(t, u, z)$ , for any (enhanced) *weak* semistable energetic solution  $(u, z)$  the map  $t \mapsto z(t)$  enjoys additional regularity properties.

**3.1. Semistable energetic solutions.** The following notion extends the one proposed in [Rou09, Def. 5.1] for the case of rate-independent processes in viscous solids, to the case of the abstract system (1.4), and to a *general* dissipation potential  $\mathcal{V}$ .

**Definition 3.1** (Semistable energetic solution). *In the setting of Hypotheses 2.2 and 2.3, we call a pair  $(u, z) : [0, T] \rightarrow \mathbf{V} \times \mathbf{Z}$  a semistable energetic solution to the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  (to the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ , respectively), if*

$$u \in L^\infty(0, T; \mathbf{U}) \cap W^{1,1}(0, T; \mathbf{V}), \quad u' \in L^\infty(0, T; \mathbf{W}) \text{ in the damped inertial system case,} \quad (3.1a)$$

$$z \in L^\infty(0, T; \mathbf{X}) \cap \text{BV}([0, T]; \mathbf{Z}) \quad (3.1b)$$

fulfill the

- *subdifferential inclusion for  $u$*

$$u''(t) + \partial\mathcal{V}(u'(t)) + \partial_u^- \mathcal{E}(t, u(t), z(t)) \ni 0 \quad \text{in } \mathbf{V}^* \text{ for a.a. } t \in (0, T), \quad (3.2)$$

*viz.  $u''(t) + \omega(t) + \xi(t) = 0$ , with  $\xi(t) \in \partial_u^- \mathcal{E}(t, u(t), z(t))$  and  $\omega(t) \in \partial\mathcal{V}(u'(t))$  for almost all  $t \in (0, T)$ ;*

- *semistability condition*

$$\mathcal{E}(t, u(t), z(t)) \leq \mathcal{E}(t, u(t), \tilde{z}) + \mathcal{R}(\tilde{z} - z(t)) \quad \text{for all } \tilde{z} \in \mathbf{Z} \text{ for all } t \in [0, T]; \quad (3.3)$$

- *energy-dissipation inequality*

$$\begin{aligned} \frac{1}{2} \|u'(t)\|_{\mathbf{W}}^2 + \int_0^t \mathcal{V}(u'(r)) + \mathcal{V}^*(-\xi(r) - u''(r)) dr + \text{Var}_{\mathcal{R}}(z; [0, t]) + \mathcal{E}(t, u(t), z(t)) \\ \leq \frac{1}{2} \|u'(0)\|_{\mathbf{W}}^2 + \mathcal{E}(0, u(0), z(0)) + \int_0^t \partial_t \mathcal{E}(r, u(r), z(r)) dr \quad \text{for all } t \in [0, T], \end{aligned} \quad (3.4)$$

*with  $\xi(r) \in \partial_u^- \mathcal{E}(r, u(r), z(r))$  the selection from (3.2), i.e.,  $\xi(r)$  fulfills (3.2) for almost all  $r \in (0, T)$ .*

Finally, we call a semistable energetic solution  $(u, z)$  *enhanced* if it fulfills the energy-dissipation inequality (3.4) on any interval  $(s, t)$ , for all  $t \in (0, T]$  and for almost all  $s \in (0, t)$ .

Observe that, in the case  $\mathcal{V}$  is a quadratic functional and independent of the state variable (cf. (4.1) later on), the energy-dissipation inequality (3.4), also in view of (3.2), becomes

$$\begin{aligned} & \frac{1}{2} \|u'(t)\|_{\mathbf{W}}^2 + \int_0^t 2\mathcal{V}(u'(s)) \, ds + \text{Var}_{\mathcal{R}}(z; [0, t]) + \mathcal{E}(t, u(t), z(t)) \\ & \leq \frac{1}{2} \|u'(0)\|_{\mathbf{W}}^2 + \mathcal{E}(0, u(0), z(0)) + \int_0^t \partial_t \mathcal{E}(s, u(s), z(s)) \, ds \quad \text{for all } t \in [0, T]. \end{aligned} \quad (3.5)$$

Our next result generalizes [Rou09, Prop. 5.2] (cf. also [Rou10, Prop. 3.2]) to the case of  $\mathcal{E}(t, \cdot, z)$  possibly nonsmooth. It addresses the passage from semistable energetic to pointwise solutions of system (1.4). To avoid overburdening the paper, we will state it in the case of gradient systems, and leave to the reader the straightforward generalization to the case with inertia. In Proposition 3.2, we have to assume that  $\mathcal{E}(t, u, \cdot)$  is of class  $C^1$ , and that  $\mathcal{E}$  fulfills a suitable *chain-rule inequality* featuring the Fréchet differential of  $\mathcal{E}(t, u, \cdot)$  and the Fréchet subdifferential of  $\mathcal{E}(t, \cdot, z)$ . Moreover, we want to restrict the arguments to the case that  $z$  is differentiable, by improving  $z \in \text{BV}(0, T; \mathbf{Z})$  to  $z \in \text{AC}([0, T]; \mathbf{Z})$ . However, for general Banach spaces this does not yet give differentiability of  $t \mapsto z(t)$ . But, assuming additionally that the space  $\mathbf{Z}$  has the Radon-Nikodým property we have  $\text{AC}([0, T]; \mathbf{Z}) = W^{1,1}([0, T]; \mathbf{Z})$ , see [DU77], i.e.,  $z'$  exists a.e. as an element of  $L^1([0, T]; \mathbf{Z})$ . As a last requirement, we also have to exclude the case that  $\mathcal{R}$  takes the value  $+\infty$ , because the resulting constraint for the test functions in the semistability condition would not allow us to fully exploit it. Thus, in the setting outlined above, we show that, any semistable energetic solution solves system (1.4) pointwise a.e. in  $(0, T)$ .

**Proposition 3.2.** *Let  $\mathbf{Z}$  comply with the Radon-Nikodým property and  $\mathcal{R}$  take values in  $[0, +\infty)$ . We suppose that  $\mathcal{E}(t, u, \cdot) \in C^1(\mathbf{Z})$  for every  $(t, u) \in [0, T] \times \text{D}_u$ , and that the following chain-rule inequality holds:*

*for every curve  $(u, z) \in \text{AC}([0, T]; \mathbf{V} \times \mathbf{Z})$  and every  $\xi \in L^1(0, T; \mathbf{V}^*)$  with*

*$\xi(t) \in \partial_u^- \mathcal{E}(t, u(t), z(t))$  for a.a.  $t \in (0, T)$  and such that*

$$\sup_{t \in [0, T]} \mathcal{G}(u(t), z(t)) < +\infty, \quad \int_0^T \mathcal{V}(u'(t)) + \mathcal{V}^*(-\xi(t)) \, dt < +\infty, \quad (3.6)$$

*then  $t \mapsto \mathcal{E}(t, u(t), z(t))$  is absolutely continuous on  $[0, T]$  and*

$$\frac{d}{dt} \mathcal{E}(t, u(t), z(t)) \geq \partial_t \mathcal{E}(t, u(t), z(t)) + \langle \xi(t), u'(t) \rangle_{\mathbf{V}} + \langle \text{D}_z \mathcal{E}(t, u(t), z(t)), z'(t) \rangle_{\mathbf{Z}} \quad \text{for a.a. } t \in (0, T).$$

*Let  $(u, z)$  be a semistable energetic solution to the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  such that, in addition,  $z \in \text{AC}([0, T]; \mathbf{Z})$ . Then,  $(u, z)$  complies with the flow rule (1.4b) pointwise a.e. in  $(0, T)$  and (3.4) holds as an energy-dissipation balance along every subinterval  $[s, t] \subset [0, T]$ .*

*Proof.* Testing the semistability (3.3) with  $\tilde{z} = z(t) + \epsilon \zeta$ , where  $\zeta$  is an arbitrary element in  $\mathbf{Z}$  and  $\epsilon > 0$  is likewise arbitrary, and arguing in the very same way as in the proof of [Rou09, Prop. 5.2] leads to the inequality  $\langle -\text{D}_z \mathcal{E}(t, u(t), z(t)), \zeta \rangle_{\mathbf{V}} \leq \mathcal{R}(\zeta)$  for every  $\zeta \in \mathbf{Z}$ . In particular, we deduce

$$\langle -\text{D}_z \mathcal{E}(t, u(t), z(t)), z'(t) \rangle_{\mathbf{Z}} \leq \mathcal{R}(z'(t)) \quad \text{for a.a. } t \in (0, T). \quad (3.7)$$

Now, we consider the energy-dissipation inequality (3.4), written without the kinetic energy term, observing that, since  $z \in \text{AC}([0, T]; \mathbf{Z})$ , we have  $\text{Var}_{\mathcal{R}}(z; [0, t]) = \int_0^t \mathcal{R}(z'(s)) \, ds$ , see, e.g., [MR15, Prop. 3.26, p. 124]. We continue (3.4) by applying the chain-rule inequality from (3.6). Hence, we obtain the following chain of inequalities for every  $t \in [0, T]$

$$\begin{aligned} & \int_0^t \mathcal{V}(u'(s)) + \mathcal{V}^*(-\xi(s)) \, ds + \int_0^t \mathcal{R}(z'(s)) \, ds \leq \mathcal{E}(0, u(0), z(0)) - \mathcal{E}(t, u(t), z(t)) + \int_0^t \partial_t \mathcal{E}(s, u(s), z(s)) \, ds \\ & \leq \int_0^t (\langle -\xi(s), u'(s) \rangle_{\mathbf{V}} + \langle -\text{D}_z \mathcal{E}(s, u(s), z(s)), z'(s) \rangle_{\mathbf{Z}}) \, ds \\ & \leq \int_0^t \mathcal{V}(u'(s)) + \mathcal{V}^*(-\xi(s)) \, ds + \int_0^t \mathcal{R}(z'(s)) \, ds, \end{aligned} \quad (3.8)$$

where the last inequality follows from the definition (2.13) of  $\mathcal{V}^*$ , and from (3.7). Thus all inequalities hold as equalities, so that (3.4) holds as an equality at every  $t \in [0, T]$  and, a fortiori, on every subinterval  $[s, t] \subset [0, T]$ .

Moreover, rearranging terms we conclude that

$$\int_0^t (\mathcal{V}(u'(s)) + \mathcal{V}^*(-\xi(s)) - \langle -\xi(s), u'(s) \rangle_{\mathbf{V}}) ds = \int_0^t (\langle -D_z \mathcal{E}(s, u(s), z(s)), z'(s) \rangle_{\mathbf{Z}} - \mathcal{R}(z'(s))) ds.$$

Since the integrand in the l.h.s.-integral is positive by the definition (2.13) of  $\mathcal{V}^*$  and the integrand on the r.h.s. is negative by (3.7), we necessarily conclude that both integrals (and in fact both integrands, pointwise) equal zero. Hence

$$\langle -D_z \mathcal{E}(t, u(t), z(t)), z'(t) \rangle_{\mathbf{Z}} = \mathcal{R}(z'(t)) \quad \text{for a.a. } t \in (0, T). \quad (3.9)$$

Combining this with (3.7) we get that  $-D_z \mathcal{E}(t, u(t), z(t)) \in \partial \mathcal{R}(z'(t))$  for almost all  $t \in (0, T)$ . This concludes the proof.  $\square$

**Remark 3.3.** A close perusal of the proof of Prop. 3.2 reveals that the validity of subdifferential inclusion (3.2) has never been used. In fact, the very same argument as in the above lines allows us to prove that, if a pair  $(u, z)$  fulfilling (3.1a)–(3.1b) with  $z \in \text{AC}([0, T]; \mathbf{Z})$  complies with the semistability (3.3) and with the energy-dissipation inequality (3.4), and if the energy complies with the conditions from Prop. 3.2, then  $(u, z)$  also fulfills the flow rule (1.4b) and the subdifferential inclusion (3.2) pointwise a.e. in  $(0, T)$ . This is in accordance with the results from [MRS13b] for rate-dependent gradient systems.

**3.2. Weak semistable energetic solutions.** Let us now introduce in the following definition a much weaker solvability notion, where the pointwise subdifferential inclusion for  $u$  is dropped. It will be useful to bring this solution concept into play in connection with the evolutionary  $\Gamma$ -convergence analysis of gradient and damped inertial systems which will be investigated in the forthcoming [RT16].

**Definition 3.4** (Weak semistable energetic solution). *In the setting of Hypotheses 2.2 and 2.3, we say that a pair  $(u, z) : [0, T] \rightarrow \mathbf{V} \times \mathbf{Z}$  as in (3.1) is a weak semistable energetic solution to the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  (to the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ , respectively) if for all  $t \in [0, T]$  it complies with the semistability condition (3.3) and with the energy-dissipation inequality (3.4), with  $\xi(s)$  a selection in  $\partial_u^- \mathcal{E}(s, u(s), z(s))$  for a.a.  $s \in (0, T)$ .*

*We call a weak semistable energetic solution  $(u, z)$  enhanced if it fulfills the energy-dissipation inequality (3.4) on any interval  $(s, t)$ , for all  $t \in (0, T]$  and for almost all  $s \in (0, t)$ .*

Although the above concept is extremely weak, slightly adapting the proof of Proposition 3.2 (cf. Remark 3.3) it can be shown that, if the energy complies with the conditions from the latter result, then any weak semistable energetic solution such that in addition  $z \in \text{AC}([0, T]; \mathbf{Z})$  is also a semistable energetic solution, as well as a pointwise solution of system (3.2, 1.4b). In Proposition 3.5 and Theorem 3.7 ahead, we will provide a different set of conditions on the energy, still allowing us to pass from *weak* semistable energetic to *semistable energetic* solutions, but *without* the condition that  $z$  is absolutely continuous.

First of all, Prop. 3.5 guarantees that a weak semistable energetic solution  $(u, z)$  is in fact a semistable energetic solution, provided that  $(u, z)$  complies with a suitable “chain-rule inequality” in integral form, cf. (3.10) below. Then, in Thm. 3.7 we provide sufficient conditions on the energy  $\mathcal{E}$ , guaranteeing that any weak semistable energetic solution to  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  (to  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ , respectively) complies with (3.10). Let us point out that, although the derivation of (3.10) is at the core of the proofs of [Rou09, Prop. 5.4] as well as [Rou10, Prop. 4.3], its role in deriving the subdifferential inclusion (3.2) from the energy-dissipation inequality (3.4) has never been highlighted in such a generality before.

**Proposition 3.5.** *Assume Hypotheses 2.2 and 2.3. Then:*

- (1) *Let  $(u, z)$  be a weak semistable energetic solution to the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  and let  $\xi \in L^1(0, T; \mathbf{V}^*)$  be a selection  $\xi(t) \in \partial_u^- \mathcal{E}(t, u(t), z(t))$  for a.a.  $t \in (0, T)$ , fulfilling the energy-dissipation inequality (3.4). Suppose that  $(u, z)$  comply with the (integral) chain rule*

$$\begin{aligned} & \int_0^t \langle \xi(s), u'(s) \rangle_{\mathbf{V}} ds \\ & \leq \mathcal{E}(t, u(t), z(t)) - \mathcal{E}(0, u(0), z(0)) - \int_0^t \partial_t \mathcal{E}(s, u(s), z(s)) ds + \text{Var}_{\mathcal{R}}(z; [0, t]) \text{ for all } t \in [0, T]. \end{aligned} \quad (3.10)$$

*Then,*

- (i) the energy-dissipation inequality (3.4) turns into an identity, holding along any subinterval  $[s, t]$ , and indeed (3.10) holds with an equality sign;
- (ii)  $(u, z, \xi)$  comply with the subdifferential inclusion (3.2).

Hence,  $(u, z)$  is a semistable energetic solution.

- (2) In the case of a damped inertial system, assume in addition that for every  $(t, z) \in [0, T] \times \mathbf{D}_z$  the map  $u \mapsto \mathcal{E}(t, u, z)$  is Gâteaux-differentiable, and that  $\mathcal{V}$  and  $\mathcal{V}^*$  comply with the enhanced coercivity

$$\exists C_V, C'_V > 0 \quad \forall (v, \xi) \in \mathbf{V} \times \mathbf{V}^* \quad \mathcal{V}(v) + \mathcal{V}^*(\xi) \geq C_V |\langle \xi, v \rangle_{\mathbf{V}}| - C'_V. \quad (3.11)$$

Let  $(u, z)$  be a weak semistable energetic solution to the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$  such that  $u$  has the further regularity

$$u \in W^{2,1}(0, T; \mathbf{V}^*) \quad \text{and} \quad t \mapsto \mathbf{D}_u \mathcal{E}(t, u(t), z(t)) \in L^\infty(0, T; \mathbf{V}^*). \quad (3.12)$$

Then, the same conclusions as above hold, and  $(u, z)$  is a semistable energetic solution.

Observe that (3.11) is for example satisfied if the viscous dissipation potential has the form  $\mathcal{V}(v) = F(\|v\|_{\mathbf{V}})$ , with  $F$  a positive convex function with superlinear growth at infinity. In this case,  $\mathcal{V}^*(\xi) = F^*(\|\xi\|_{\mathbf{V}^*})$ , with  $F^*$  the conjugate of  $F$ , and (3.11) holds with  $C_V = 1$  and  $C'_V = 0$ .

*Proof.* We will confine the discussion to the case of a damped inertial system, as the argument for the gradient case is completely analogous. Since  $\int_0^t \mathcal{V}(u'(s)) + \mathcal{V}^*(-\mathbf{D}_u \mathcal{E}(s, u(s), z(s)) - u''(s)) \, ds \leq C$ , from (3.11) we gather that the map  $t \mapsto \langle -\mathbf{D}_u \mathcal{E}(t, u(t), z(t)) - u''(t), u'(t) \rangle_{\mathbf{V}}$  is in  $L^1(0, T)$ . Since  $t \mapsto \mathbf{D}_u \mathcal{E}(t, u(t), z(t))$  is in  $L^\infty(0, T; \mathbf{V}^*)$  we have that  $t \mapsto \langle -\mathbf{D}_u \mathcal{E}(t, u(t), z(t)), u'(t) \rangle_{\mathbf{V}}$  is in  $L^1(0, T)$ , hence we conclude that

$$t \mapsto \langle u''(t), u'(t) \rangle_{\mathbf{V}} \quad \text{is in } L^1(0, T). \quad (3.13)$$

Combining (3.10) with the energy-dissipation inequality (3.4) we conclude the chain of inequalities

$$\begin{aligned} & \int_0^t \mathcal{V}(u'(s)) + \mathcal{V}^*(-\mathbf{D}_u \mathcal{E}(s, u(s), z(s)) - u''(s)) \, ds \\ & \leq \frac{1}{2} \|u'(0)\|_{\mathbf{W}}^2 - \frac{1}{2} \|u'(t)\|_{\mathbf{W}}^2 - \int_0^t \langle \mathbf{D}_u \mathcal{E}(s, u(s), z(s)), u'(s) \rangle_{\mathbf{V}} \, ds = \int_0^t \langle -\mathbf{D}_u \mathcal{E}(s, u(s), z(s)) - u''(s), u'(s) \rangle_{\mathbf{V}} \, ds \end{aligned}$$

for all  $t \in [0, T]$ , where we have used that

$$\langle u''(t), u'(t) \rangle_{\mathbf{V}} = \frac{1}{2} \frac{d}{dt} \|u'\|_{\mathbf{W}}^2(t) \quad \text{for a.a. } t \in (0, T)$$

since  $(\mathbf{V}, \mathbf{W}, \mathbf{V}^*)$  is a Hilbert (or Gelfand) triple, cf. (2.3), as well as (3.13). Therefore, we find

$$\int_0^t (\mathcal{V}(u'(s)) + \mathcal{V}^*(-\mathbf{D}_u \mathcal{E}(s, u(s), z(s)) - u''(s)) - \langle -\mathbf{D}_u \mathcal{E}(s, u(s), z(s)) - u''(s), u'(s) \rangle_{\mathbf{V}}) \, ds \leq 0. \quad (3.14)$$

Now, it follows from the definition of the conjugate  $\mathcal{V}^*$  that also the opposite of inequality (3.14) holds. Thus, we conclude that (3.14) holds as an equality a.e. in  $(0, T)$ . Hence,

$$-u''(s) - \mathbf{D}_u \mathcal{E}(s, u(s), z(s)) \in \partial \mathcal{V}(u'(s)) \quad \text{for a.a. } s \in (0, T),$$

whence (3.2). Furthermore, from the above arguments it follows that (3.4) and (3.10) hold as equalities on every interval  $[0, t] \subset [0, T]$ . This concludes the proof.  $\square$

**Remark 3.6.** A perusal of the proof of Prop. 3.2 allows us to conclude that, along a weak semistable energetic solution  $(u, z)$  with the property  $(u, z) \in \text{AC}([0, T]; \mathbf{V} \times \mathbf{Z})$ , the chain rule inequality in (3.6) and the integral chain rule (3.10) are equivalent.

In the proof of Thm. 3.7 below, we derive the chain-rule inequality (3.10), in the case the map  $u \mapsto \mathcal{E}(t, u, z)$  is Gâteaux-differentiable but *without requiring*  $z \in \text{AC}([0, T]; \mathbf{Z})$ , from the semistability condition (3.3), mimicking the Riemann-sum procedure from the proof of [Rou09, Prop. 5.4], see also [Rou10, Prop. 4.3], which in turn is based on the argument first developed in [DMFT05].

**Theorem 3.7.** *Assume Hypotheses 2.2 and 2.3, that for every  $(t, z) \in [0, T] \times \mathbf{D}_z$  the map  $u \mapsto \mathcal{E}(t, u, z)$  is Gâteaux-differentiable, and that*

$$\forall M > 0 \exists S > 0 \quad \forall t \in [0, T], \forall u, u_1, u_2 \in \mathbf{D}_u, \bar{z} \in \mathbf{D}_z : \quad (3.15a)$$

$$\begin{cases} (u, \bar{z}) \in \mathcal{S}_M \Rightarrow \|\mathbf{D}_u \mathcal{E}(t, u, \bar{z})\|_{\mathbf{V}^*} \leq S, \\ (u_1, \bar{z}), (u_2, \bar{z}) \in \mathcal{S}_M \Rightarrow \|\mathbf{D}_u \mathcal{E}(t, u_1, \bar{z}) - \mathbf{D}_u \mathcal{E}(t, u_2, \bar{z})\|_{\mathbf{V}^*} \leq S \|u_1 - u_2\|_{\mathbf{V}}, \end{cases}$$

with  $\mathcal{S}_M$  as in (2.8), and that  $\partial_t \mathcal{E}$  satisfies analogous Lipschitz estimates, i.e.

$$\forall \tilde{M} > 0 \exists \tilde{S} > 0 \quad \forall t_1, t_2 \in [0, T], \forall u_1, u_2 \in \mathbf{D}_u, \bar{z} \in \mathbf{D}_z : \quad (3.15b)$$

$$(u_1, \bar{z}), (u_2, \bar{z}) \in \mathcal{S}_{\tilde{M}} \Rightarrow \begin{cases} |\partial_t \mathcal{E}(t_1, u_1, \bar{z}) - \partial_t \mathcal{E}(t_2, u_1, \bar{z})| \leq \tilde{S} |t_1 - t_2|, \\ |\partial_t \mathcal{E}(t_1, u_1, \bar{z}) - \partial_t \mathcal{E}(t_1, u_2, \bar{z})| \leq \tilde{S} \|u_1 - u_2\|_{\mathbf{V}}. \end{cases}$$

Let  $(u, z)$  be a weak semistable energetic solution to the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  (to the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ , respectively). Then,  $(u, z)$  complies with (3.10) and the energy-dissipation inequality (3.4) holds as an identity.

*Proof.* Preliminarily, we observe that along a weak semistable energetic solution  $(u, z)$  there holds

$$\sup_{t \in [0, T]} \mathcal{G}(u(t), z(t)) \leq C. \quad (3.16)$$

This can be checked by arguing in the very same way as in the proof of the forthcoming Proposition 7.2, namely starting from the energy-dissipation inequality (3.4) and exploiting estimate (2.10). Therefore, thanks to the first of (3.15a), we have that the map  $t \mapsto \mathbf{D}_u \mathcal{E}(t, u(t), z(t))$  is in  $L^\infty(0, T; \mathbf{V}^*)$ .

Now, let  $\eta > 0$  be fixed, and let  $(\mathbf{t}_i)_{i=1}^{N_\eta}$  be a partition of the interval  $[0, t]$ , with  $0 < \mathbf{t}_i - \mathbf{t}_{i-1} \leq \eta$  for all  $i = 1, \dots, N_\eta$ . We test the semistability (3.3) at time  $t = \mathbf{t}_{i-1}$  with  $\tilde{z} = z(\mathbf{t}_i)$ , obtaining

$$\mathcal{E}(\mathbf{t}_{i-1}, u(\mathbf{t}_{i-1}), z(\mathbf{t}_{i-1})) \leq \mathcal{E}(\mathbf{t}_{i-1}, u(\mathbf{t}_{i-1}), z(\mathbf{t}_i)) + \mathcal{R}(z(\mathbf{t}_i) - z(\mathbf{t}_{i-1})).$$

Using that

$$\begin{aligned} \mathcal{E}(\mathbf{t}_{i-1}, u(\mathbf{t}_{i-1}), z(\mathbf{t}_i)) &= \mathcal{E}(\mathbf{t}_i, u(\mathbf{t}_i), z(\mathbf{t}_i)) + \mathcal{E}(\mathbf{t}_i, u(\mathbf{t}_{i-1}), z(\mathbf{t}_i)) - \mathcal{E}(\mathbf{t}_i, u(\mathbf{t}_i), z(\mathbf{t}_i)) \\ &\quad + \mathcal{E}(\mathbf{t}_{i-1}, u(\mathbf{t}_{i-1}), z(\mathbf{t}_i)) - \mathcal{E}(\mathbf{t}_i, u(\mathbf{t}_{i-1}), z(\mathbf{t}_i)) \end{aligned}$$

we deduce by the chain rule

$$\begin{aligned} \mathcal{E}(\mathbf{t}_{i-1}, u(\mathbf{t}_{i-1}), z(\mathbf{t}_{i-1})) &\leq \mathcal{E}(\mathbf{t}_i, u(\mathbf{t}_i), z(\mathbf{t}_i)) + \mathcal{R}(z(\mathbf{t}_i) - z(\mathbf{t}_{i-1})) \\ &\quad - \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} \langle \mathbf{D}_u \mathcal{E}(\mathbf{t}_i, u(s), z(\mathbf{t}_i)), u'(s) \rangle_{\mathbf{V}} ds - \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} \partial_t \mathcal{E}(s, u(\mathbf{t}_{i-1}), z(\mathbf{t}_i)) ds, \end{aligned}$$

which we add up over the index  $i = 1, \dots, N_\eta$ . Taking into account that  $\sum_{i=1}^{N_\eta} \mathcal{R}(z(\mathbf{t}_i) - z(\mathbf{t}_{i-1})) \leq \text{Var}_{\mathcal{R}}(z; [0, t])$ , we arrive at

$$\begin{aligned} \mathcal{E}(0, u(0), z(0)) &\leq \mathcal{E}(t, u(t), z(t)) + \text{Var}_{\mathcal{R}}(z; [0, t]) - \sum_{i=1}^{N_\eta} \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} \langle \mathbf{D}_u \mathcal{E}(\mathbf{t}_i, u(s), z(\mathbf{t}_i)), u'(s) \rangle_{\mathbf{V}} ds \\ &\quad - \sum_{i=1}^{N_\eta} \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} \partial_t \mathcal{E}(s, u(\mathbf{t}_{i-1}), z(\mathbf{t}_i)) ds. \end{aligned}$$

We now deal with the two latter terms. Observe that

$$\begin{aligned} &\sum_{i=1}^{N_\eta} \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} \langle \mathbf{D}_u \mathcal{E}(\mathbf{t}_i, u(s), z(\mathbf{t}_i)), u'(s) \rangle_{\mathbf{V}} ds \\ &= \sum_{i=1}^{N_\eta} \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} \langle \mathbf{D}_u \mathcal{E}(\mathbf{t}_i, u(\mathbf{t}_i), z(\mathbf{t}_i)), u'(\mathbf{t}_i) \rangle_{\mathbf{V}} ds + \sum_{i=1}^{N_\eta} \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} \langle \mathbf{D}_u \mathcal{E}(\mathbf{t}_i, u(s), z(\mathbf{t}_i)) - \mathbf{D}_u \mathcal{E}(\mathbf{t}_i, u(\mathbf{t}_i), z(\mathbf{t}_i)), u'(s) \rangle_{\mathbf{V}} ds \\ &+ \sum_{i=1}^{N_\eta} \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} \langle \mathbf{D}_u \mathcal{E}(\mathbf{t}_i, u(\mathbf{t}_i), z(\mathbf{t}_i)), u'(s) - u'(\mathbf{t}_i) \rangle_{\mathbf{V}} ds \doteq S_1^\eta + S_2^\eta + S_3^\eta. \end{aligned} \quad (3.17)$$

Now, in view of estimate (3.15a) (which is applicable thanks to (3.16)), we have that

$$|S_2^\eta| \leq C \sum_{i=1}^{N_\eta} \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} \|u(s) - u(\mathbf{t}_i)\|_{\mathbf{V}} \|u'(s)\|_{\mathbf{V}} ds \leq C \|u'\|_{L^1(0,T;\mathbf{V})} \max_{i=1,\dots,N_\eta} \max_{s \in [\mathbf{t}_{i-1}, \mathbf{t}_i]} \|u(s) - u(\mathbf{t}_i)\|_{\mathbf{V}}. \quad (3.18)$$

Since  $u \in W^{1,1}(0,T;\mathbf{V})$ , we have  $\max_{i=1,\dots,N_\eta} \max_{s \in [\mathbf{t}_{i-1}, \mathbf{t}_i]} \|u(s) - u(\mathbf{t}_i)\|_{\mathbf{V}} \rightarrow 0$  as  $\eta \downarrow 0$ , hence  $S_2^\eta \rightarrow 0$ . Moreover,

$$|S_3^\eta| \leq \|D_u \mathcal{E}(\cdot, u(\cdot), z(\cdot))\|_{L^\infty(0,T;\mathbf{V}^*)} \sum_{i=1}^{N_\eta} \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} \|u'(s) - u'(\mathbf{t}_i)\|_{\mathbf{V}} ds \quad (3.19)$$

where the latter estimate is again a consequence of (3.15a). We may choose the partition  $(\mathbf{t}_i)_{i=1}^{N_\eta}$  in such a way that  $\sum_{i=1}^{N_\eta} \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} \|u'(s) - u'(\mathbf{t}_i)\|_{\mathbf{V}} ds \rightarrow 0$  as  $\eta \downarrow 0$ , hence also  $S_3^\eta \rightarrow 0$ , and such that

$$S_1^\eta \rightarrow \int_0^t \langle D_u \mathcal{E}(s, u(s), z(s)), u'(s) \rangle_{\mathbf{V}} ds,$$

to which, ultimately, the term on the left-hand side of (3.17) converges.

Analogously,

$$\begin{aligned} & \sum_{i=1}^{N_\eta} \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} \partial_t \mathcal{E}(s, u(\mathbf{t}_{i-1}), z(\mathbf{t}_i)) ds \\ &= \sum_{i=1}^{N_\eta} \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} \partial_t \mathcal{E}(\mathbf{t}_i, u(\mathbf{t}_i), z(\mathbf{t}_i)) ds + \sum_{i=1}^{N_\eta} \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} (\partial_t \mathcal{E}(s, u(\mathbf{t}_i), z(\mathbf{t}_i)) - \partial_t \mathcal{E}(\mathbf{t}_i, u(\mathbf{t}_i), z(\mathbf{t}_i))) ds \\ & \quad + \sum_{i=1}^{N_\eta} \int_{\mathbf{t}_{i-1}}^{\mathbf{t}_i} (\partial_t \mathcal{E}(s, u(\mathbf{t}_{i-1}), z(\mathbf{t}_i)) - \partial_t \mathcal{E}(s, u(\mathbf{t}_i), z(\mathbf{t}_i))) ds \doteq S_4^\eta + S_5^\eta + S_6^\eta. \end{aligned}$$

With a suitable choice of the partition we have that  $S_4^\eta \rightarrow \int_0^t \partial_t \mathcal{E}(s, u(s), z(s)) ds$ , while exploiting (3.15b) and arguing similarly as above we have that  $S_5^\eta, S_6^\eta \rightarrow 0$ . Hence, (3.10) ensues.  $\square$

### 3.3. Improved temporal regularity of semistable energetic solutions by uniform convexity of $\mathcal{E}$ .

Under stronger convexity properties of the energy functional, it has been shown in [MT04, MR07, TM10] in the fully rate-independent setting that energetic solutions enjoy better temporal regularity. In particular, uniform convexity of the energy functional in the pair  $(u, z)$  implies temporal Hölder-, or even Lipschitz continuity, [TM10, Thm. 4.5]. In the coupled rate-independent/rate-dependent setting, however, the uniform convexity of  $\mathcal{E}(t, u(t), \cdot)$  in general only ensures the *continuity in time* of the semistable variable  $z$ , and in some special cases its Hölder continuity. To give a more precise outline, in what follows uniform convexity is only required with respect to the  $z$ -variable and the condition is formulated with respect to an additional Banach space  $\mathbf{S} \supset \mathbf{X}$ , which may or may not coincide with  $\mathbf{Z}$  or  $\mathbf{X}$ . Namely, we impose that

$$\begin{aligned} & \exists \alpha \geq 2 \exists C_* > 0 \forall t \in [0, T], \forall u \in \mathbf{V}, \forall z_0, z_1 \in \mathbf{D}_z, \forall \theta \in [0, 1], \text{ setting } z_\theta = \theta z_1 + (1 - \theta) z_0 : \\ & \mathcal{E}(t, u, z_\theta) + \mathcal{R}(z_\theta - z_0) + C_* \theta (1 - \theta) \|z_1 - z_0\|_{\mathbf{S}}^\alpha \\ & \leq \theta \mathcal{E}(t, u, z_0) + (1 - \theta) \mathcal{E}(t, u, z_1) + \mathcal{R}(z_1 - z_0), \end{aligned} \quad (3.20)$$

which the reader may compare with [TM10, condition (4.4a)]. We refer to [TM10, Sec. 4, 5] for more details and examples on the uniform convexity inequality (3.20) featuring the exponent  $\alpha \geq 2$ .

Now, let  $(u, z) : \mathbf{V} \times \mathbf{Z}$  be a weak semistable energetic solution. Choosing  $\theta \in (0, 1)$ ,  $u = u(t)$ ,  $z_0 = z(t)$  and  $z_1 = z(s)$ , with fixed  $s < t \in [0, T]$ , in (3.20) yields

$$\begin{aligned} 0 & \leq \mathcal{E}(t, u(t), z_\theta) + \mathcal{R}(z_\theta - z(t)) - \mathcal{E}(t, u(t), z(t)) \\ & \leq \theta (\mathcal{E}(t, u(t), z(s)) + \mathcal{R}(z(s) - z(t)) - \mathcal{E}(t, u(t), z(t)) - C_* (1 - \theta) \|z(s) - z(t)\|_{\mathbf{S}}^\alpha). \end{aligned} \quad (3.21)$$

Then, dividing by  $\theta > 0$  and letting  $\theta \rightarrow 1$  results in the following improved semistability inequality:

$$\mathcal{E}(t, u(t), z(t)) + C_* \|z(t) - z(s)\|_{\mathbf{S}}^\alpha \leq \mathcal{E}(t, u(t), z(s)) + \mathcal{R}(z(s) - z(t)). \quad (3.22)$$

Via the energy-dissipation inequality (3.4), under further continuity assumptions on the partial derivatives  $\partial_t \mathcal{E}$  and  $D_u \mathcal{E}$  one may therefore obtain an estimate of the form  $\|z(t) - z(s)\|_{\mathbf{S}}^\alpha \leq \omega(t - s)$  with  $\omega$  a modulus



of continuity also depending on  $u'$  and  $u''$ . To deduce this estimate, however, it is crucial that the energy-dissipation inequality (3.4) is satisfied on *every* subinterval  $[s, t] \subset [0, T]$ , which of course is a defining property of energetic solutions in the fully rate-independent setting, but a strengthening of the notion of semistable energetic solutions in the coupled setting. Moreover, it has to be stressed, that ultimately concluding the continuity estimate for  $\|z(t) - z(s)\|_{\mathbf{S}}^{\alpha}$  requires the Gâteaux differentiability of  $\mathcal{E}(t, \cdot, z)$ , and, in the case with inertia, additionally that  $u \in W^{2,1}(0, T; \mathbf{V}^*)$ . All in all, depending on the assumptions imposed on  $\mathcal{E}(t, \cdot, \cdot)$ , we obtain the temporal regularity results (R1)–(R2) stated in Thm. 3.8 below. In particular, let us mention in advance that (R2) guarantees Hölder continuity with an exponent  $h < 1/2$ .

**Theorem 3.8** (Continuity of weak semistable energetic solution by uniform convexity). *Let  $(u, z) : \mathbf{V} \times \mathbf{Z}$  be a weak semistable energetic solution for the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ , resp. the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ . Assume that*

- (A1)  $(u, z)$  satisfies the energy-dissipation inequality (3.4) even on all subintervals  $[s, t] \subset [0, T]$ ,
- (A2)  $\mathcal{E}(t, u(t), \cdot) : \mathbf{Z} \rightarrow [0, +\infty]$  is uniformly convex as in (3.20), featuring a Banach space  $\mathbf{S}$ , with  $\mathbf{X} \subset \mathbf{S}$  continuously,
- (A3)  $\mathcal{E}(t, \cdot, z)$  is Gâteaux-differentiable for every  $(t, z) \in [0, T] \times D_z$ ,
- (A4) in the case of a damped inertial system,  $u \in W^{2,1}(0, T; \mathbf{V}^*)$  (cf. (3.12)) and  $t \mapsto |\langle u''(t), u'(t) \rangle_{\mathbf{V}}| \in L^1(0, T)$ .

Then, the following regularity results hold true:

- (R1) In addition to (A1)–(A4) assume that the energy functional  $\mathcal{E}$  satisfies the following Hölder continuity condition: There are constants  $c_* > 0, \beta_u \in (0, 1]$  such that for all  $s, t \in [0, T]$ , all  $(u_0, z_1), (u_1, z_1) \in \mathcal{S}_E$ :

$$|\mathcal{E}(t, u_1, z_1) - \mathcal{E}(t, u_0, z_1)| \leq c_* \|u_1 - u_0\|_{\mathbf{V}}^{\beta_u}. \quad (3.23)$$

Then,  $z$  complies with the following estimate:

$$C_* \|z(t) - z(s)\|_{\mathbf{S}}^{\alpha} \leq C|t - s| + \int_s^t |\langle u''(r), u'(r) \rangle_{\mathbf{V}}| dr + c_* \left( \int_s^t \|u'(r)\|_{\mathbf{V}} dr \right)^{\beta_u} \quad (3.24)$$

and thus has the regularity  $z \in C^0([0, T]; \mathbf{S})$ .

- (R2) In addition to (A1)–(A4) & (3.23) assume that  $\mathcal{E}(t, \cdot, z)$  complies with the following gradient estimate (cf. also Hyp. 5.3, (5.8)): There exist constants  $C_3, C_4, C_5 > 0$  and  $\sigma \in [1, \infty)$  such that

$$\|D_u \mathcal{E}(t, u, z)\|_{\mathbf{V}^*}^{\sigma} \leq C_3 \mathcal{E}(t, u, z) + C_4 \|u\|_{\mathbf{V}} + C_5 \quad \text{for all } (t, u, z) \in [0, T] \times D_u \times D_z. \quad (3.25)$$

Then,  $z$  complies with the following estimate:

$$C_* \|z(t) - z(s)\|_{\mathbf{S}}^{\alpha} \leq C|t - s| + c_* \left( \int_s^t \|u'(r)\|_{\mathbf{V}} dr \right)^{\beta_u} \quad (3.26)$$

and thus has the regularity  $z \in C^{0,h}([0, T]; \mathbf{S})$ . Moreover, if the dissipation  $\mathcal{V}$  has  $p$ -growth for some  $p > 1$ , namely  $\mathcal{V}(v) \geq \tilde{C}_1 \|v\|_{\mathbf{V}}^p - \tilde{C}_2$  for some  $\tilde{C}_1, \tilde{C}_2 > 0$ , then

$$z \in C^{0,h}([0, T]; \mathbf{S}) \quad \text{with the Hölder-exponent } h = \beta_u / (p' \alpha) < 1/2, \quad (3.27)$$

where  $p' = p/(p-1)$ .

*Proof.* The proof carries over the steps of [TM10, Thm. 4.5] to the present coupled setting.

Starting out from the improved semistability inequality (3.22) we deduce the following estimate using the energy-dissipation inequality (3.4):

$$\begin{aligned}
C_* \|z(t) - z(s)\|_{\mathbf{S}}^{\beta_u} &\leq \mathcal{E}(s, u(s), z(t)) + \mathcal{R}(z(t) - z(s)) - \mathcal{E}(s, u(s), z(s)) \\
&\leq \mathcal{E}(t, u(t), z(t)) + \text{Var}_{\mathcal{R}}(z; [s, t]) - \mathcal{E}(s, u(s), z(s)) + \mathcal{E}(s, u(s), z(t)) - \mathcal{E}(t, u(t), z(t)) \\
&\leq \frac{1}{2} (\|u'(s)\|_{\mathbf{W}}^2 - \|u'(t)\|_{\mathbf{W}}^2) - \int_s^t \mathcal{V}(u'(r)) + \mathcal{V}^*(-D_u \mathcal{E}(r, u(r), z(r)) - u''(r)) \, dr \\
&\quad + \int_s^t \partial_t \mathcal{E}(r, u(r), z(r)) - \partial_t \mathcal{E}(r, u(s), z(t)) \, dr + \mathcal{E}(t, u(s), z(t)) - \mathcal{E}(t, u(t), z(t)) \\
&= - \int_s^t \langle u''(r), u'(r) \rangle_{\mathbf{V}} \, dr - \int_s^t \mathcal{V}(u'(r)) + \mathcal{V}^*(-D_u \mathcal{E}(r, u(r), z(r)) - u''(r)) \, dr \\
&\quad + \int_s^t \partial_t \mathcal{E}(r, u(r), z(r)) - \partial_t \mathcal{E}(r, u(s), z(t)) \, dr + \mathcal{E}(t, u(s), z(t)) - \mathcal{E}(t, u(t), z(t))
\end{aligned} \tag{3.28}$$

**Ad (R1):** Using the power control (2.6c) on the terms involving  $\partial_t \mathcal{E}$  and (3.23) on the energy difference  $\mathcal{E}(t, u(s), z(t)) - \mathcal{E}(t, u(t), z(t))$ , we can further estimate (3.28) as follows:

$$\begin{aligned}
(3.28) &\leq \int_s^t |\langle u''(r), u'(r) \rangle_{\mathbf{V}}| \, dr + \int_s^t (C_1(\mathcal{E}(r, u(r), z(r)) + \mathcal{E}(r, u(s), z(t))) + 2C_2) \, dr + c_* \|u(s) - u(t)\|_{\mathbf{V}}^{\beta_u} \\
&\leq \int_s^t |\langle u''(r), u'(r) \rangle_{\mathbf{V}}| \, dr + C|t - s| + c_* \left( \int_s^t \|u'(r)\|_{\mathbf{V}} \, dr \right)^{\beta_u},
\end{aligned}$$

where we have used that  $\mathcal{E}(r, u(r), z(r)) + \mathcal{E}(r, u(s), z(t)) \leq 2E$ , since  $(u(r), z(r)), (u(s), z(t)) \in \mathcal{S}_E$ . Moreover, we exploited that  $0 \leq \int_s^t \mathcal{V}(u'(r)) + \mathcal{V}^*(-D_u \mathcal{E}(r, u(r), z(r)) - u''(r)) \, dr \leq C$  by the energy-dissipation inequality (3.4), so that the negative of this term can be dropped to find an estimate from above. Since  $\langle u''(\cdot), u'(\cdot) \rangle_{\mathbf{V}} \in L^1(0, T)$ , the first integral term on the right-hand side is absolutely continuous, whereas the second one is continuous due to  $u' \in L^1(0, T; \mathbf{V})$ . This proves estimate (3.24) and yields  $z \in C^0([0, T]; \mathbf{S})$ .

**Ad (R2):** Instead of dropping the negative viscous term, as we did for (R1), we now apply the Young-Fenchel inequality, i.e.

$$- \int_s^t \mathcal{V}(u'(r)) + \mathcal{V}^*(-D_u \mathcal{E}(r, u(r), z(r)) - u''(r)) \, dr \leq \int_s^t \langle u''(r) + D_u \mathcal{E}(r, u(r), z(r)), u'(r) \rangle_{\mathbf{V}} \, dr.$$

Inserting this in (3.28), and subsequently making use of the gradient estimate (3.25), (3.23), and the power control (2.6c) we find:

$$\begin{aligned}
(3.28) &\leq \int_s^t \langle D_u \mathcal{E}(r, u(r), z(r)), u'(r) \rangle_{\mathbf{V}} \, dr \\
&\quad + \int_s^t \partial_t \mathcal{E}(r, u(r), z(r)) - \partial_t \mathcal{E}(r, u(s), z(t)) \, dr + \mathcal{E}(t, u(s), z(t)) - \mathcal{E}(t, u(t), z(t)) \\
&\leq C|t - s| + c_* \left( \int_s^t \|u'(r)\|_{\mathbf{V}} \, dr \right)^{\beta_u},
\end{aligned} \tag{3.29}$$

where we argued as for (R1) to estimate the energy difference and energies within sublevels  $\mathcal{S}_E$ . Hence, we have estimate (3.26) and  $z \in C^0([0, T]; \mathbf{S})$ .

To verify the Hölder continuity in the case of  $\mathcal{V}$  of  $p$ -growth, we continue the chain of inequalities (3.29) by applying Hölder's inequality with exponent  $p$  to the second term on the r.h.s., and thus we find

$$(3.28) \leq C|s - t| + c_* \left( \int_s^t \|u'(\zeta)\|_{\mathbf{V}} \, d\zeta \right)^{\beta_u} \leq |s - t|^{\beta_u/p'} (C|s - t|^{1-\beta_u/p'} + c_* \|u'\|_{L^p(0, T; \mathbf{V})}^{\beta_u}) \leq C|s - t|^{\beta_u/p'}.$$

Hence, we conclude (3.27).  $\square$

#### 4. EXISTENCE RESULTS FOR THE GRADIENT SYSTEM $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$

Throughout this section, we will focus on the analysis of gradient systems.

In Section 4.1 we will first address the case in which the dissipation potential  $\mathcal{V}$  is quadratic. In this context, our main assumption on the energy  $\mathcal{E}$  will be a sort of *uniform Fréchet subdifferentiability* condition

w.r.t. the variable  $u$ , drawn from [MRS13a] and implied, e.g., by  $\lambda$ -convexity type properties of the mapping  $u \mapsto \mathcal{E}(t, u, z)$ . Hence, in our first existence result, Theorem 4.9, we are going to show that the approximate solutions constructed by a carefully devised time-discretization scheme converge to a *semistable energetic* solution of the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ .

In Section 4.2 we will extend our analysis to a reasonably broad class of convex dissipation potentials  $\mathcal{V}$  with superlinear growth at infinity. For the limit passage from time-discrete to time-continuous, we will resort to an approach, and to ideas, from the *variational theory for gradient flows*. In this context, we will not require  $\lambda$ -convexity properties on the energy functional  $\mathcal{E}$ . However, we will need to *directly* impose a closedness condition on  $\partial_u^- \mathcal{E}$  more stringent than the closedness property guaranteed by the uniform subdifferentiability condition. Hence, we will give our second existence result, Theorem 4.13, stating that the approximate solutions constructed by time discretization converge to a *weak* semistable energetic solution to the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ , which clearly turns into a *semistable energetic* one under the additional condition (3.15) on  $\mathcal{E}$ .

In Section 4.3 we will compare the two sets of conditions on  $\mathcal{E}$  in two examples. Furthermore, we will hint at possible extensions of Theorems 4.9 and 4.13.

**4.1. Case 1:  $\mathcal{V}$  quadratic.** Throughout this section, we will confine the discussion to *quadratic* dissipation potentials for  $u$ , namely we will suppose that

$$\mathcal{V}(v) = \frac{1}{2}a(v, v) \quad \text{with } a : \mathbf{V} \times \mathbf{V} \rightarrow \mathbb{R} \text{ a continuous and coercive bilinear form,} \quad (4.1)$$

thus inducing on  $\mathbf{V}$  a Hilbert space structure. Hence, for every  $v \in \mathbf{V}$

$$\partial \mathcal{V}(v) = \{Av\} \quad \text{with } A : \mathbf{V} \rightarrow \mathbf{V}^* \text{ the linear operator associated with } a.$$

Still, we will continue to use the notation  $\partial \mathcal{V}$ , also to make the comparison with the results of Sec. 4.2 more transparent. We now detail the time-discretization scheme which will give rise to the approximate solutions for the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ .

**Time-discretization scheme.** Let  $\tau = T/N_\tau > 0$  be the time step, inducing a partition

$$\mathcal{P}_\tau = \{0 = t_0 < t_1 < \dots < t_{N_\tau-1} < t_{N_\tau} = T\}$$

of the interval  $[0, T]$ . We shall denote by  $\bar{\tau}_\tau$  and  $\underline{\tau}_\tau$  the left-continuous and right-continuous piecewise constant interpolants associated with the partition  $\mathcal{P}_\tau$ , namely

$$\bar{\tau}_\tau(0) = \underline{\tau}_\tau(0) := 0, \quad \bar{\tau}_\tau(t) := t_n \quad \text{for } t \in (t_{n-1}, t_n], \quad \underline{\tau}_\tau(t) := t_{n-1} \quad \text{for } t \in [t_{n-1}, t_n). \quad (4.2)$$

Of course, for every  $t \in [0, T]$  we have  $\bar{\tau}_\tau(t) \downarrow t$  and  $\underline{\tau}_\tau(t) \uparrow t$  as  $\tau \downarrow 0$ .

Starting from the Cauchy data  $(u_0, z_0) \in D_u \times D_z$  (cf. (4.17) below), we construct discrete solutions  $(u_\tau^n, z_\tau^n)_{n=1}^{N_\tau}$  by *alternating (decoupled)* time-incremental minimization (cf. the forthcoming Remark 7.6). More precisely, we first find  $z_\tau^n$  by minimizing with respect to  $z$  the sum of the energy and of the dissipation potential  $\mathcal{R}$ , with  $u_\tau^{n-1}$  and  $z_\tau^{n-1}$  given from the previous step. Then, we construct  $u_\tau^n$  by minimizing, now with respect to  $u$ , the sum of the energy and of the dissipation  $\mathcal{V}$ , with  $z_\tau^n$  now given.

**Problem 4.1.** Let  $(u_\tau^0, z_\tau^0) := (u_0, z_0) \in D_u \times D_z$ , and for  $n = 1, \dots, N_\tau$ , find

$$z_\tau^n \in \underset{z \in \mathbf{Z}}{\text{Argmin}} \left( \tau \mathcal{R} \left( \frac{z - z_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, u_\tau^{n-1}, z) \right), \quad (4.3a)$$

$$u_\tau^n \in \underset{u \in \mathbf{V}}{\text{Argmin}} \left( \tau \mathcal{V} \left( \frac{u - u_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, u, z_\tau^n) \right). \quad (4.3b)$$

With the direct method in the Calculus of Variations, exploiting Hypotheses 2.2 and 2.7, it is not difficult to check that for all  $0 < \tau < \tau_o$  Problem 4.1 admits a solution  $(u_\tau^n, z_\tau^n)_{n=1}^{N_\tau}$ . Then, we construct the approximate solutions as the left-continuous and right-continuous interpolants of  $(u_\tau^n, z_\tau^n)_{n=1}^{N_\tau}$ , defined by

$$\bar{u}_\tau(t) := u_\tau^n, \quad \bar{z}_\tau(t) := z_\tau^n \quad \text{for } t \in (t_{n-1}, t_n], \quad \underline{u}_\tau(t) := u_\tau^{n-1}, \quad \underline{z}_\tau(t) := z_\tau^{n-1} \quad \text{for } t \in [t_{n-1}, t_n). \quad (4.4)$$

We also consider the piecewise linear interpolants

$$u_\tau(t) := \frac{t - t_{n-1}}{\tau} u_\tau^n + \frac{t_n - t}{\tau} u_\tau^{n-1}, \quad z_\tau(t) := \frac{t - t_{n-1}}{\tau} z_\tau^n + \frac{t_n - t}{\tau} z_\tau^{n-1}, \quad \text{for } t \in [t_{n-1}, t_n). \quad (4.5)$$

**Uniform Fréchet subdifferentiability.** For the derivation of all a priori estimates on the approximate solutions, it will be crucial to obtain a *discrete* form of the energy-dissipation inequality (3.4), cf. (7.30) ahead. Its proof will be based on a suitable condition on the Fréchet subdifferential  $\partial_u^- \mathcal{E} : \mathbf{V} \rightrightarrows \mathbf{V}^*$ , that can be interpreted as a *uniform Fréchet subdifferentiability* of the energy  $u \mapsto \mathcal{E}(t, u, z)$  (in this connection, we refer to the discussions in [MRS13b, Sec. 2] and [MRS13a, Sec. 3] where analogous properties were introduced and exploited). We may understand it like this: While the inequality (2.11) defining the Fréchet subdifferential has a *local* character, with (4.6) below we make it *global*, at the price of having a negative term on the right-hand side. Observe that for our purposes it will be sufficient to require this property on energy sublevels.

**Hypothesis 4.2** (Uniform Fréchet subdifferentiability). *For all  $E > 0$  there exists  $\Lambda_E \geq 0$  such that*

$$\begin{aligned} \mathcal{E}(t, v, z) - \mathcal{E}(t, u, z) &\geq \langle \xi, v - u \rangle_{\mathbf{V}} - \Lambda_E \|v - u\|_{\mathbf{V}}^2 \\ &\text{for all } t \in [0, T], (u, z), (v, z) \in \mathcal{S}_E \text{ and for all } \xi \in \partial_u^- \mathcal{E}(t, u, z). \end{aligned} \quad (4.6)$$

**Remark 4.3.** Observe that, the validity of inequality (4.6) implies on its own that  $\xi \in \partial_u^- \mathcal{E}(t, u, z)$ , cf. (2.11).

Notice that, if (4.6) holds with a remainder term of the type  $\Lambda_E \|v - u\|_{\mathbf{V}} \|v - u\|_{\mathbf{Y}}$ , and  $\mathbf{Y}$  a space such that  $\mathbf{V} \subset \mathbf{Y}$  with a continuous embedding, then (4.6) holds with the term  $\tilde{\Lambda}_E \|v - u\|_{\mathbf{V}}^2$  for some other  $\tilde{\Lambda}_E > 0$ . This can be checked by observing that there exists  $C_{\mathbf{V}, \mathbf{Y}} > 0$  such that  $\|v - u\|_{\mathbf{Y}} \leq C_{\mathbf{V}, \mathbf{Y}} \|v - u\|_{\mathbf{V}}$  for every  $u, v \in \mathcal{S}_E$ .

A sufficient condition for (4.6) is provided by the  $\lambda$ -convexity of the energy  $u \mapsto \mathcal{E}(t, u, z)$ , on energy sublevels, namely

**Lemma 4.4.** *Let  $\mathcal{E} : [0, T] \times \mathbf{V} \times \mathbf{Z} \rightarrow (-\infty, +\infty]$  fulfill Hypothesis 2.2 and*

$$\begin{aligned} \forall E > 0 \quad \exists \Lambda_E \geq 0 \quad \forall t \in [0, T], (u, z), (v, z) \in \mathcal{S}_E, \forall \theta \in (0, 1) : \\ \mathcal{E}(t, (1 - \theta)u + \theta v, z) &\leq (1 - \theta)\mathcal{E}(t, u, z) + \theta\mathcal{E}(t, v, z) + \Lambda_E \theta(1 - \theta) \|v - u\|_{\mathbf{V}}^2. \end{aligned} \quad (4.7)$$

*Then,  $\mathcal{E}$  complies with Hypothesis 4.2.*

The *proof* follows the very same lines as that in [MRS13a, Lemma 3.26], hence it is omitted.

**Example 4.5.** *Let  $\phi : \mathbb{R}^{d \times d} \rightarrow \mathbb{R}$ ,  $\phi(e) := \operatorname{tr} e|e|$  with  $\operatorname{tr} e = \sum_{i=1}^d e_{ii}$  for any  $e = (e_{ij})_{i,j=1}^d \in \mathbb{R}^{d \times d}$ ,  $1 < d \in \mathbb{N}$ . The function  $\phi$  is not convex, as can be checked e.g. for  $d = 2$ , and with the diagonal matrices  $e := \operatorname{diag}(1/2, 1/2)$  and  $\tilde{e} = \operatorname{diag}(-1, 1)$ . For  $\theta = 1/2$ , this choice gives  $\frac{1}{2}\sqrt{\frac{5}{8}} = \phi(\theta e + (1 - \theta)\tilde{e}) > \theta\phi(e) + (1 - \theta)\phi(\tilde{e}) = \frac{1}{2}\sqrt{\frac{1}{2}}$ . But  $\phi$  is  $\Lambda$ -convex for any  $\Lambda > \sqrt{d}$ . To verify this statement, we calculate that*

$$\phi''(e)[\tilde{e}, \tilde{e}] = 2 \operatorname{tr} \tilde{e} \frac{e:\tilde{e}}{|e|} + \operatorname{tr} e \frac{|\tilde{e}|^2}{|e|} - \frac{\operatorname{tr} e}{|e|} \left( \frac{e:\tilde{e}}{|e|} \right)^2. \quad (4.8)$$

*By distinguishing cases and by straightforward calculations, for any choice of  $e, \tilde{e} \in \mathbb{R}^{d \times d}$  it can be shown that*

$$\phi''(e)[\tilde{e}, \tilde{e}] \geq -2|\operatorname{tr} \tilde{e}||\tilde{e}| \geq -2\sqrt{d}|\tilde{e}|^2. \quad (4.9)$$

*where the second inequality stems from*

$$|\operatorname{tr} e| \leq \sum_{i=1}^d |e_{ii}| \leq \sqrt{d} \left( \sum_{i=1}^d (e_{ii}^2) \right)^{1/2} \leq \sqrt{d} \left( \sum_{i,j=1}^d (e_{ij}^2) \right)^{1/2} = \sqrt{d}|e|.$$

*In fact, (4.9) proves that  $\phi$  is  $\Lambda$ -convex for any  $\Lambda \geq \sqrt{d}$ . From this we conclude that the corresponding integral functional defined on a domain  $\Omega \subset \mathbb{R}^d$ ,*

$$\Phi : \mathbf{V} \rightarrow (-\infty, +\infty), \quad \Phi(\mathbf{u}) = \int_{\Omega} \phi(\varepsilon(\mathbf{u})) \, dx \quad \text{with } \phi(\varepsilon(\mathbf{u})) := |\operatorname{tr} \varepsilon(\mathbf{u})| |\varepsilon(\mathbf{u})| \quad (4.10)$$

*satisfies  $\Lambda$ -convexity (4.7) on  $\mathbf{V} = H_0^1(\Omega; \mathbb{R}^d)$  for any  $\Lambda > \sqrt{d}$ . We also refer to Sec. 4.3 for further examples.*

As mentioned in the above lines, the primary role of condition (4.6) is in the derivation of a *discrete energy-dissipation inequality*. Furthermore,

- (1) In the case  $\mathbf{U} \Subset \mathbf{V}$ , in combination with Hypothesis 4.6 below, condition (4.6) in Hypothesis 4.2 will guarantee a closedness property of the graph of  $\partial_u^- \mathcal{E}$  with respect to strong-weak topology, and a continuity property of  $\mathcal{E}$  (cf. Lemma 4.7 ahead), that will play a key role in the passage to the time-continuous limit.

- (2) If only the *continuous* embedding  $\mathbf{U} \subset \mathbf{V}$  is assumed, it will be necessary to require, in addition to Hypothesis 4.2, a version of the closedness property of  $\partial_u^- \mathcal{E}$  in Bochner spaces, cf. Hypothesis 4.8 below.

**Case 1: A recovery sequence condition for  $\mathcal{E}(t, \cdot, z)$ .** In the case  $\mathbf{U} \Subset \mathbf{V}$ , we impose the following

**Hypothesis 4.6** (Recovery sequence condition for  $\mathcal{E}(t, \cdot, z)$ ). *For all sequences  $(t_n)_n \subset [0, T]$ ,  $(u_n)_n \subset \mathbf{V}$ , and  $(z_n)_n \subset \mathbf{Z}$  such that*

$$t_n \rightarrow t, \quad u_n \rightarrow u \text{ in } \mathbf{V}, \quad z_n \rightarrow z \text{ in } \mathbf{Z}, \quad \sup_n \mathcal{G}(u_n, z_n) \leq C$$

and for every  $\tilde{u} \in D_u$  there exists  $\tilde{u}_n \rightarrow \tilde{u}$  in  $\mathbf{V}$  such that

$$\limsup_{n \rightarrow \infty} (\mathcal{E}(t_n, \tilde{u}_n, z_n) - \mathcal{E}(t_n, u_n, z_n)) \leq \mathcal{E}(t, \tilde{u}, z) - \mathcal{E}(t, u, z). \quad (4.11)$$

We postpone the proof of the following result to Section 7.2.

**Lemma 4.7.** *Suppose that  $\mathbf{U} \Subset \mathbf{V}$  and let  $\mathcal{E}$  fulfill Hypotheses 2.2, 4.2 and 4.6. Then, for all sequences  $(t_n, u_n, z_n, \xi_n) \subset [0, T] \times \mathbf{V} \times \mathbf{Z} \times \mathbf{V}^*$  there holds*

$$\left\{ \begin{array}{l} t_n \rightarrow t, \\ u_n \rightarrow u \quad \text{in } \mathbf{V}, \\ z_n \rightarrow z \quad \text{in } \mathbf{Z}, \\ \sup_n \mathcal{G}(u_n, z_n) \leq C, \\ \xi_n \rightarrow \xi \quad \text{in } \mathbf{V}^* \\ \xi_n \in \partial_u^- \mathcal{E}(t_n, u_n, z_n) \end{array} \right\} \Rightarrow \xi \in \partial_u^- \mathcal{E}(t, u, z) \quad \text{and} \quad \mathcal{E}(t_n, u_n, z_n) \rightarrow \mathcal{E}(t, u, z). \quad (4.12)$$

**Case 2: A “Bochner-space” version of closedness à la Minty of  $\partial_u^- \mathcal{E}$ .** In the case  $\mathbf{U} \subset \mathbf{V}$  continuously, it is no longer possible to deduce from Hypotheses 4.2 & 4.6 the closedness property (4.12). Therefore we have to require a suitable variant of it. Observe that (4.12) has a “static”, or “pointwise-in-time” character, in that it only involves sequences converging in the spaces  $\mathbf{V}$ ,  $\mathbf{Z}$ , and  $\mathbf{V}^*$ , and it is in fact suited to the application of Young measure arguments in the limit passage from time-discrete to time-continuous, cf. the proof of Theorem 4.13.

Instead, the forthcoming Hypothesis 4.8 considers sequences of time-dependent functions, converging in Lebesgue/Sobolev-Bochner spaces. It will be *directly* used in the limit passage in the discrete version of the equation for the rate-dependent variable  $u$ . Indeed, (4.13a) below also features a lim sup condition, in the spirit of Minty’s trick within the theory of maximal monotone operators (see e.g. [Att84, Lemma 3.57, p. 356]), that can be verified by arguing on the discrete  $u$ -equation, checking that all terms apart from the one to identify either pass to the limit, or can be handled by lower semicontinuity. For this, the quadratic character of  $\mathcal{V}$  will play a crucial role.

Finally, in Hyp. 4.8 we will distinguish between a *weak*, and a *strong* form of the closedness property, depending on whether we also obtain convergence of the energies.

**Hypothesis 4.8** (Closedness of  $\partial_u^- \mathcal{E}$  à la Minty). *For all sequences  $t_n : [0, T] \rightarrow [0, T]$ ,  $(u_n)_n \subset L^\infty(0, T; \mathbf{U}) \cap H^1(0, T; \mathbf{V})$ ,  $(z_n)_n \subset L^\infty(0, T; \mathbf{X}) \cap \text{BV}([0, T]; \mathbf{Z})$ , and  $(\xi_n)_n \subset L^2(0, T; \mathbf{V}^*)$ ,*

$$\text{if } \left\{ \begin{array}{l} t_n \rightarrow t \quad \text{pointwise a.e. in } (0, T), \\ u_n \xrightarrow{*} u \quad \text{in } L^\infty(0, T; \mathbf{U}) \cap H^1(0, T; \mathbf{V}), \\ z_n \xrightarrow{*} z \text{ in } L^\infty(0, T; \mathbf{X}), \quad z_n(t) \xrightarrow{\sigma\mathbf{X}} z(t) \text{ in } \mathbf{X} \quad \text{for all } t \in [0, T], \\ \exists C > 0 \quad \forall n \in \mathbb{N} \quad \forall t \in [0, T] : \quad \mathcal{G}(u_n(t), z_n(t)) \leq C, \\ \xi_n \rightarrow \xi \quad \text{in } L^2(0, T; \mathbf{V}^*), \\ \xi_n(t) \in \partial_u^- \mathcal{E}(t_n(t), u_n(t), z_n(t)) \quad \text{for a.a. } t \in (0, T), \\ \limsup_{n \rightarrow \infty} \int_0^T \langle \xi_n, u_n \rangle_{\mathbf{V}} dt \leq \int_0^T \langle \xi, u \rangle_{\mathbf{V}} dt, \end{array} \right. \quad (4.13a)$$

then there holds  $\xi(t) \in \partial_u^- \mathcal{E}(t(t), u(t), z(t))$  for a.a.  $t \in (0, T)$ .

We will also consider a stronger form of (4.13a), namely for all sequences

$$(t_n)_n, (u_n)_n, (z_n)_n, (\xi_n)_n \text{ as in (4.13a) there holds} \quad (4.13b)$$

$$\xi(t) \in \partial_u^- \mathcal{E}(t(t), u(t), z(t)), \quad \text{and} \quad \mathcal{E}(t_n(t), u_n(t), z_n(t)) \rightarrow \mathcal{E}(t(t), u(t), z(t)) \quad \text{for a.a. } t \in (0, T).$$

In Section 4.3 we will provide the example of an energy complying with Hypotheses 4.2, 4.6, and 4.8.

**Statement of our first existence result.** Under the basic conditions given in Sec. 2, the uniform Fréchet subdifferentiability from Hyp. 4.2, and ( $\mathbf{U} \subseteq \mathbf{V}$  & Hyp. 4.6) or, in *alternative*, Hyp. 4.8, Theorem 4.9 below states that, up to a subsequence, any family of approximate solutions constructed via time discretization, converge to a semistable energetic solution  $(u, z)$  of the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  for  $\mathcal{V}$  quadratic, provided that the initial data  $(u_0, z_0)$  fulfill the semistability (3.3) at  $t = 0$ , i.e.

$$\mathcal{E}(0, u_0, z_0) \leq \mathcal{E}(0, u_0, \tilde{z}) + \mathcal{R}(\tilde{z} - z_0) \quad \text{for all } \tilde{z} \in \mathbf{Z}. \quad (4.14)$$

Observe that (4.14) is necessary since with our concept of (weak) semistable energetic solution we claim that the semistability holds at *all* times. We also detail in which spaces the approximate solutions converge, distinguishing a first set of convergences which will hold also in the case of  $\mathcal{V}$  with general superlinear growth (cf. Theorem 4.13 ahead), from convergence (4.16), related to the quadratic character of  $\mathcal{V}$ . Under suitable conditions we will also obtain the energy convergence (4.19), which will allow us to prove that the pair  $(u, z)$  is an *enhanced* semistable energetic solution, cf. Def. 3.1.

**Theorem 4.9** (Semistable energetic solutions for  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ ,  $\mathcal{V}$  quadratic). *Assume Hypotheses 2.2, 2.3, 2.5, and 2.7. In addition, suppose that  $\mathcal{V}$  is quadratic, cf. (4.1), that  $\partial_{\underline{u}}^- \mathcal{E}$  complies with Hyp. 4.2, and that*

- (1) *either  $\mathbf{U} \subseteq \mathbf{V}$  and in addition Hyp. 4.6 is valid;*
- (2) *or, Hyp. 4.8 holds.*

Let  $u_0 \in \mathbf{D}_u$  and  $z_0 \in \mathbf{D}_z$  be fixed, fulfilling (4.14), and let  $(\bar{u}_\tau, \underline{u}_\tau, u_\tau, \bar{z}_\tau, \underline{z}_\tau, z_\tau)_\tau$  be a family of approximate solutions constructed via the time-discretization scheme in Problem 4.1.

Then, for every sequence  $\tau_k \downarrow 0$  as  $k \rightarrow \infty$  there exist a (not relabeled) subsequence, and functions  $(u, z)$  as in (3.1) such that, in addition,  $u \in H^1(0, T; \mathbf{V})$  and the following convergences hold as  $k \rightarrow \infty$

$$u_{\tau_k} \rightharpoonup u \quad \text{in } W^{1,1}(0, T; \mathbf{V}), \quad (4.15a)$$

$$\bar{u}_{\tau_k}, u_{\tau_k} \xrightarrow{*} u \quad \text{in } L^\infty(0, T; \mathbf{U}), \quad (4.15b)$$

$$\bar{u}_{\tau_k}(t), \underline{u}_{\tau_k}(t), u_{\tau_k}(t) \rightharpoonup u(t) \quad \text{in } \mathbf{U} \text{ for all } t \in [0, T], \quad (4.15c)$$

$$\bar{z}_{\tau_k}, z_{\tau_k} \xrightarrow{*} z \quad \text{in } L^\infty(0, T; \mathbf{X}), \quad (4.15d)$$

$$\bar{z}_{\tau_k}(t), z_{\tau_k}(t) \xrightarrow{\sigma \mathbf{X}} z(t) \quad \text{in } \mathbf{X} \text{ for all } t \in [0, T], \quad (4.15e)$$

$$(\bar{z}_{\tau_k}(t) - \underline{z}_{\tau_k}(t)) \xrightarrow{\sigma \mathbf{X}} 0 \quad \text{in } \mathbf{X} \text{ for almost all } t \in (0, T), \quad (4.15f)$$

$$\bar{z}_{\tau_k}(t), z_{\tau_k}(t) \rightharpoonup z(t) \quad \text{in } \mathbf{Z} \text{ for all } t \in [0, T], \quad (4.15g)$$

(while for  $(\underline{z}_{\tau_k}(t))_k$  the analogues of (4.15e) and (4.15g) hold for almost all  $t \in (0, T)$ ), and, moreover,

$$u_{\tau_k} \rightharpoonup u \quad \text{in } H^1(0, T; \mathbf{V}). \quad (4.16)$$

The pair  $(u, z)$  is a semistable energetic solution to the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ , satisfying the initial conditions

$$u(0) = u_0, \quad z(0) = z_0. \quad (4.17)$$

Furthermore,

- (i) *under the conditions in (1), we also have the additional convergences*

$$\bar{u}_{\tau_k}(t), u_{\tau_k}(t), \underline{u}_{\tau_k}(t) \rightarrow u(t) \text{ in } \mathbf{V} \text{ for all } t \in [0, T], \quad (4.18)$$

$$\mathcal{E}(\bar{u}_{\tau_k}(t), \bar{u}_{\tau_k}(t), \bar{z}_{\tau_k}(t)) \rightarrow \mathcal{E}(t, u(t), z(t)) \quad \text{for a.a. } t \in (0, T), \quad (4.19)$$

and the pair  $(u, z)$  is an *enhanced semistable energetic solution*, complying with

$$\begin{aligned} & \int_s^t \mathcal{V}(u'(r)) + \mathcal{V}^*(-\xi(r)) dr + \text{Var}_{\mathcal{R}}(z; [s, t]) + \mathcal{E}(t, u(t), z(t)) \\ & \leq \mathcal{E}(s, u(s), z(s)) + \int_s^t \partial_t \mathcal{E}(r, u(r), z(r)) dr \quad \text{for all } t \in (0, T] \text{ for } s = 0, \text{ and a.a. } s \in (0, t). \end{aligned} \quad (4.20)$$

- (ii) *If Hyp. 4.8 holds in the stronger form (4.13b), then (4.19) and (4.20) are valid and  $(u, z)$  is enhanced.*

**Remark 4.10.** As it will be clear from the proof of Theorem 4.9, performed in Section 7.2, under the conditions guaranteeing the energy convergence (4.19), it would be sufficient to adopt the *weaker* variant of Hypothesis 2.5 introduced in Remark 2.6 in order to obtain the existence of semistable energetic solutions. In that case, however, the semistability condition (3.3) would only hold at *almost all*  $t \in (0, T)$ .

Let us also mention in advance that the same observation is valid for Theorems 4.9, 5.4, and 5.6 ahead.

**4.2. Case 2:  $\mathcal{V}$  with general superlinear growth.** We now address the analysis of system (1.4) for a fairly broad class of dissipation potentials  $\mathcal{V}$  that fulfill, in addition to condition (2.12a), the following property

$$\mathcal{V}^*(\xi_1) = \mathcal{V}^*(\xi_2) \quad \text{for all } \xi_1, \xi_2 \in \partial\mathcal{V}(u) \quad \text{for all } u \in \mathbf{V}. \quad (4.21)$$

It was observed in [MRS13b, Rmk. 2.1] that the dissipation potential  $\mathcal{V}$  complies with (4.21) at  $u \in \mathbf{V}$  if and only if the mapping  $\lambda \mapsto \mathcal{V}(\lambda u)$  is differentiable at  $\lambda = 1$ . Therefore, condition (4.21) is satisfied for example when  $\mathcal{V}$  is a linear combination of (positively) homogeneous, or differentiable, convex potentials.

Condition (4.21) will have a technical role in the derivation of a discrete energy-dissipation inequality, different from the one proved for  $\mathcal{V}$  quadratic, for the approximate solutions arising from time discretization, cf. Lemma 7.1 ahead. The proof of such inequality, developed in [MRS13b], extends the *variational techniques* for gradient flows from [Amb95, AGS08] to the case of systems driven by general dissipation potentials with superlinear growth, complying with (4.21).

**Time-discretization scheme.** In the case of a general dissipation potential  $\mathcal{V}$ , we construct the discrete solutions by a scheme different from the one in Problem 4.1, cf. again Remark 7.6 for more comments. In fact, first we solve a minimum problem for  $u$ , with  $u_\tau^{n-1}$  and  $z_\tau^{n-1}$  given from the previous step, and then we solve for  $z$ , with  $u_\tau^n$  now given.

**Problem 4.11.** Let  $(u_\tau^0, z_\tau^0) := (u_0, z_0) \in D_u \times D_z$ , and for  $n = 1, \dots, N_\tau$ , find

$$u_\tau^n \in \underset{u \in \mathbf{V}}{\operatorname{Argmin}} \left( \tau \mathcal{V} \left( \frac{u - u_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, u, z_\tau^{n-1}) \right), \quad (4.22a)$$

$$z_\tau^n \in \underset{z \in \mathbf{Z}}{\operatorname{Argmin}} \left( \tau \mathcal{R} \left( \frac{z - z_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, u_\tau^n, z) \right). \quad (4.22b)$$

With the direct method in the Calculus of Variations, exploiting Hypotheses 2.2 and 2.7, it is not difficult to check that for all  $0 < \tau < \tau_o$  Problem 4.11 admits a solution  $(u_\tau^n, z_\tau^n)_{n=1}^{N_\tau}$ .

As in Section 4.1, we construct approximate solutions by taking the piecewise constant and linear interpolants of the elements  $(u_\tau^n, z_\tau^n)_{n=1}^{N_\tau}$ . Nonetheless, for the derivation of the approximate version of the energy-dissipation inequality (3.4) in the forthcoming Lemma 7.1, it will be crucial to resort to yet another notion of interpolant of  $(u_\tau^n)_{n=1}^{N_\tau}$ , which was first introduced by E. DE GIORGI within the *Minimizing Movements* theory (see [Amb95, AGS08]). It is defined in the following way: the map  $t \mapsto \tilde{u}_\tau(t)$  is Lebesgue measurable in  $(0, T)$  and satisfies

$$\begin{cases} \tilde{u}_\tau(0) = u_0, \\ \tilde{u}_\tau(t) \in \underset{u \in \mathbf{V}}{\operatorname{Argmin}} \left\{ r \mathcal{V} \left( \frac{u - u_\tau^{n-1}}{r} \right) + \mathcal{E}(t, u, z_\tau^{n-1}) \right\} \quad \text{for } t = t_{n-1} + r \in (t_{n-1}, t_n]. \end{cases} \quad (4.23)$$

Hereafter, we will call  $\tilde{u}_\tau$  *variational interpolant*. We refer to [RS06, Rmk. 4.4], [MRS13b, Sec. 4] for a thorough discussion on the existence of the above measurable selection. Observe that when  $t = t_n$  the minimum problems (4.22a) and (4.23) coincide, hence we may assume  $\bar{u}_\tau(t_n) = \underline{u}_\tau(t_n) = u_\tau(t_n) = \tilde{u}_\tau(t_n)$  for all  $n = 1, \dots, N_\tau$ .

**A closedness condition.** In Section 7.1 we will see that the variational interpolant  $\tilde{u}_\tau$  and the piecewise constant one  $\bar{z}_\tau$  comply with the discrete energy-dissipation inequality (7.8). The latter will be the starting point in the derivation of all a priori estimates, and in it we will pass to the limit to conclude that any limit pair  $(u, z)$  of the approximate solutions complies with the energy-dissipation inequality (3.4). Let us stress that, the usage of the variational techniques from the theory of gradient flows enables us to obtain (7.8) without assuming the uniform Fréchet subdifferentiability condition from Hypothesis 4.2.

However, in order to pass to the limit in (7.8) as  $\tau \downarrow 0$ , we will exploit in a key way a closedness property of the graph of  $\partial_u^- \mathcal{E}$  that has the same “static” character as (4.12) and thus applies to “pointwise-in-time” convergences. Likewise, it will be exploited in combination with a Young measure argument. Nonetheless, as

we are no longer requiring  $\mathbf{U} \Subset \mathbf{V}$  like in Lemma 4.7, boundedness of the energies no longer turns the weak convergence in  $\mathbf{V}$  into strong. Hence, the sequence  $(u_n)_n$  featuring in (4.24a) below is supposed to be weakly converging in  $\mathbf{V}$ , only. As for Hyp. 4.8, we will also set apart the case in which, in addition, energy convergence is guaranteed.

**Hypothesis 4.12** (Closedness of  $\partial_u^- \mathcal{E}$ , Continuity of  $\mathcal{E}$ ). *For all sequences  $(t_n, u_n, z_n)_n \subset [0, T] \times D_u \times D_z$  and  $(\xi_n)_n \subset \mathbf{V}^*$*

$$\left\{ \begin{array}{l} t_n \rightarrow t, u_n \rightharpoonup u \text{ in } \mathbf{V}, z_n \rightarrow z \text{ in } \mathbf{Z}, \sup_n \mathcal{G}(u_n, z_n) \leq C, \\ \xi_n \rightharpoonup \xi \text{ in } \mathbf{V}^*, \\ \xi_n \in \partial_u^- \mathcal{E}(t_n, u_n, z_n) \end{array} \right\} \Rightarrow \xi \in \partial_u^- \mathcal{E}(t, u, z). \quad (4.24a)$$

We will also consider the following enhanced closedness/continuity condition:

$$\begin{aligned} & \text{for all sequences } (t_n, u_n, z_n)_n \text{ and } (\xi_n)_n \text{ in the conditions of (4.24a) there holds} \\ & \xi \in \partial_u^- \mathcal{E}(t, u, z) \text{ and } \lim_{n \rightarrow \infty} \mathcal{E}(t_n, u_n, z_n) = \mathcal{E}(t, u, z). \end{aligned} \quad (4.24b)$$

Observe that, since  $\mathcal{E}$  is lower semicontinuous, it is sufficient to have  $\limsup_{n \rightarrow \infty} \mathcal{E}(t_n, u_n, z_n) \leq \mathcal{E}(t, u, z)$  for (4.24b) to hold. Clearly, (4.24b) implies (4.12).

**Our second existence result.** The proof of Theorem 4.13 will be carried out in Sec. 7.1.

**Theorem 4.13** (Weak semistable energetic solutions for  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ ,  $\mathcal{V}$  general). *Assume Hypotheses 2.2, 2.3, 2.5, and 2.7. In addition, suppose that  $\mathcal{V}$  complies with (4.21), and that  $\partial_u^- \mathcal{E}$  from (2.11) complies with (4.24a) from Hyp. 4.12. Let  $(\bar{u}_\tau, \underline{u}_\tau, u_\tau, \tilde{u}_\tau, \bar{z}_\tau, \underline{z}_\tau, z_\tau)_\tau$  be a family of approximate solutions constructed via the time-discretization scheme in Problem 4.11, starting from data  $u_0 \in D_u$  and  $z_0 \in D_z$  fulfilling (4.14).*

*Then, for every sequence  $\tau_k \downarrow 0$  as  $k \rightarrow \infty$  there exist a (not relabeled) subsequence, and functions  $(u, z)$  as in (3.1) such that convergences (4.15) hold, as well as*

$$\tilde{u}_{\tau_k} \overset{*}{\rightharpoonup} u \text{ in } L^\infty(0, T; \mathbf{U}), \quad \tilde{u}_{\tau_k}(t) \rightarrow u(t) \text{ in } \mathbf{U} \text{ for all } t \in [0, T], \quad (4.25)$$

*and the pair  $(u, z)$  is a weak semistable energetic solution to the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ , fulfilling the initial conditions  $u(0) = u_0$  and  $z(0) = z_0$ .*

*Furthermore, if the enhanced closedness condition (4.24b) holds, then  $(u, z)$  fulfills (4.20) and is thus an enhanced weak semistable energetic solution.*

Finally, the following result is a consequence of Prop. 3.5 and of Thm. 3.7.

**Corollary 4.14.** *Under the assumptions of Theorem 4.13, if in addition  $\mathcal{E}$  complies with (3.15), then  $(u, z)$  from Thm. 4.13 is a semistable energetic solution to the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ , complying with the energy-dissipation balance along every subinterval  $[s, t] \subset [0, T]$ .*

**4.3. Discussion of the closedness conditions and examples.** In this section, we illustrate and compare Hypotheses 4.2, 4.6, 4.8, and 4.12 on a prototypical class of energy functionals.

We start by discussing the closedness hypotheses/properties so far introduced, namely

- (1) the “static” closedness properties (4.12) from Lemma 4.7 and (4.24a), (4.24b) from Hypothesis 4.12;
- (2) the “Bochner-space” version of closedness à la Minty from Hyp. 4.8, in its two forms (4.13a) and (4.13b).

We have the obvious implications

$$\begin{aligned} (4.24b) & \not\Rightarrow (4.24a), \\ & \not\Rightarrow (4.12), \end{aligned} \quad (4.13b) \Rightarrow (4.13a).$$

Instead, there seems to be no relation between the “static” and the “Bochner-space” closedness properties. The former do not imply the latter, because the weak convergences in Bochner/Sobolev spaces imposed in (4.13a) do not yield pointwise in time convergences, and thus do not allow for the application of the “static” properties. In turn, the validity of (4.13a)/(4.13b), which of course could be restricted to functions constant in time, does not entail any of (4.12), (4.24a), (4.24b), because of the lim sup-condition featuring in (4.13a)/(4.13b).



This is well reflected by the following examples. There, since the closedness properties essentially concern the dependence of  $\mathcal{E}$  on the variable  $u$ , we will consider a very simple (yet significant) dependence on the variable  $z$ , aiming at a more transparent discussion and postponing to Section 6 more complex examples.

**Admissible energy functionals.** Following the discussion in [MRS13a, Sec. 5], we consider energies  $\mathcal{E} : [0, T] \times L^2(\Omega) \times L^1(\Omega) \rightarrow (-\infty, +\infty]$  of the form

$$\mathcal{E}(t, u, z) := \begin{cases} \int_{\Omega} g(z)\beta(\nabla u) + W(u) dx - \langle f(t), u \rangle_{W^{1,p}(\Omega)} & \text{if } u \in \mathbf{D} \\ +\infty & \text{if } u \in L^2(\Omega) \setminus \mathbf{D} \end{cases} \quad (4.26)$$

for all  $t \in [0, T]$  and  $z \in L^1(\Omega)$ , where  $\Omega \subset \mathbb{R}^d$  is a bounded Lipschitz domain, the nonlinear functions  $g$ ,  $\beta$ , and  $W$  fulfill

$$g \in \text{BC}(\mathbb{R}) \text{ and } \exists c_0 > 0 \forall z \in \mathbb{R} : g(z) \geq c_0; \quad (4.27a)$$

$$\beta : \mathbb{R}^d \rightarrow [0, +\infty) \text{ is differentiable, convex and} \quad (4.27b)$$

$$\exists p > 1, \exists c_1, c_2, c_3 > 0 \forall A \in \mathbb{R}^d : c_1|A|^p - c_2 \leq \beta(A) \leq c_3(|A|^p + 1); \quad (4.27c)$$

$$W : \mathbb{R} \rightarrow (-\infty, +\infty] \text{ is bounded from below and } \lambda\text{-convex for some } \lambda \in \mathbb{R}; \quad (4.27d)$$

(in (4.27a) BC stands for *bounded and continuous*), while  $f \in C^1([0, T]; W^{1,p}(\Omega)^*)$ . To fix ideas we consider zero Dirichlet boundary conditions and thus set

$$\mathbf{D} := \{u \in W_0^{1,p}(\Omega) : W(u) \in L^1(\Omega)\} = \mathbf{D}_u. \quad (4.28)$$

It can be easily verified that  $\partial_t \mathcal{E}(t, u, z) = -\langle f'(t), u \rangle_{W^{1,p}(\Omega)}$  complies with (2.6c). Taking into account that  $W$  is bounded from below and exploiting the Poincaré inequality it is also immediate to check that  $\mathcal{E}$  is itself bounded from below, and the sublevels of the functional  $\mathcal{G}(u) := \sup_{t \in [0, T]} \mathcal{E}(t, u)$  are bounded in  $\mathbf{U} := W^{1,p}(\Omega)$ . Thus, the choice of the ambient space  $\mathbf{V}$  determines restrictions on  $p$ , namely

$$\text{if } \mathbf{V} = L^2(\Omega), \text{ we have } \mathbf{U} = W^{1,p}(\Omega) \Subset \mathbf{V} \text{ for all } p > 2d/(2+d); \quad (4.29a)$$

$$\text{if } \mathbf{V} = H^1(\Omega), \text{ we have } \mathbf{U} = W^{1,p}(\Omega) \subset \mathbf{V} \text{ for all } p \geq 2. \quad (4.29b)$$

We now discuss the uniform Fréchet subdifferentiability condition in Hypothesis 4.2. In fact, the energy in Example 4.15 below also complies with the Fréchet subdifferentiability condition in the forthcoming Hyp. 5.2.

**Example 4.15.** *Since  $W$  is  $\lambda$ -convex (we may suppose  $\lambda < 0$  without loss of generality), we have for every  $u, v \in \mathbf{D}$  and  $\theta \in [0, 1]$*

$$\mathcal{E}(t, (1-\theta)u + \theta v, z) \leq (1-\theta)\mathcal{E}(t, u, z) + \theta\mathcal{E}(t, v, z) - \frac{\lambda}{2}\theta(1-\theta)\|v-u\|_{L^2(\Omega)}^2. \quad (4.30)$$

Hence, (4.7) (and, a fortiori, (4.6)) holds with  $\mathbf{V} = L^2(\Omega)$  and the constant  $\Lambda = \frac{|\lambda|}{2}$  independent of the energy sublevel.

Note that in the special case  $\beta(\nabla u) = \frac{1}{2}|\nabla u|^2$ , arguing as in [MRS13a, Ex. 5.1], we can improve (4.30) using the 1-convexity of  $\beta$ . Hence, in place of  $-\frac{\lambda}{2}\theta(1-\theta)\|v-u\|_{L^2(\Omega)}^2$ , on the r.h.s. of (4.30) we find

$$-\frac{1}{2}\theta(1-\theta)(\|\nabla(v-u)\|_{L^2(\Omega)}^2 + \lambda\|v-u\|_{L^2(\Omega)}^2) \leq \frac{1}{2}\theta(1-\theta)((-1-\rho\lambda)\|\nabla(v-u)\|_{L^2(\Omega)}^2 - \lambda C_{\rho}\|v-u\|_{L^1(\Omega)}^2)$$

thanks to the Gagliardo-Nirenberg inequality  $\|w\|_{L^2(\Omega)} \leq C_{\text{GN}}\|w\|_{L^1(\Omega)}^{2/(d+2)}\|\nabla w\|_{L^2(\Omega)}^{d/(d+2)} + \|w\|_{L^1(\Omega)}$  and to Young's inequality, with  $\rho > 0$  arbitrary and  $C_{\rho} > 0$  accordingly determined. Therefore, choosing  $\rho \leq 1/|\lambda|$ , we conclude (4.7) (and, a fortiori, (4.6)) in a stronger form (cf. Remark 4.3), with the remainder term involving the square norm of  $\mathbf{Y} := L^1(\Omega)$ , and  $\Lambda_E = \frac{1}{2}C_{\rho}|\lambda|$ .

Within the subsequent examples we illustrate the closedness property (4.12) and Hypotheses 4.8, 4.12. We start with Hypotheses 4.12 and we see that only a comparably small subclass of the energy functionals complying with (4.26) & (4.27) fulfill (4.24a), (4.24b), while consecutive generalizations still meet Hypothesis 4.8 and property (4.12).

**Example 4.16.** We consider  $\mathcal{E} : [0, T] \times \mathbf{V} \times \mathbf{Z} \rightarrow (-\infty, +\infty]$ ,

$$\mathcal{E}(t, u, z) = \int_{\Omega} \left( \frac{g(z)}{2} |\nabla u|^2 + \frac{1}{4}(u^2 - 1)^2 \right) dx - \langle f(t), u \rangle_{W^{1,2}(\Omega)} \quad \text{if } u \in \mathbf{D}, \text{ cf. (4.28),} \quad (4.31)$$

and  $+\infty$  otherwise. In this way, we omit other contributions to  $\mathcal{E}$  that would only depend on the variable  $z$  as they do not affect the Fréchet subdifferential of  $\mathcal{E}$  w.r.t.  $u$ ; nonetheless, let us implicitly assume that such contributions contain a regularizing gradient term and that, therefore,

$$\mathbf{X} = W^{1,q}(\Omega) \quad \text{for some } q > 1, \quad \text{while } \mathbf{Z} = L^1(\Omega). \quad (4.32)$$

We see that the energy functional (4.31) falls into the class defined by (4.26) & (4.27) for the special case of  $p = 2$ .

Let us first take  $\mathbf{V} = L^2(\Omega)$ . It is not difficult to check that the Fréchet subdifferential of  $u \mapsto \mathcal{E}(t, u, z)$  w.r.t. the  $L^2(\Omega)$ -topology has domain

$$\begin{aligned} \text{dom}(\partial_u^- \mathcal{E}) &= \{(t, u, z) \in [0, T] \times H^1(\Omega) \times L^1(\Omega) : -\text{div}(g(z)\nabla u) + u^3 - u - f(t) \in L^2(\Omega)\} \text{ and} \\ \partial_u^- \mathcal{E}(t, u, z) &= \{-\text{div}(g(z)\nabla u) + u^3 - u - f(t)\} \text{ for all } (t, u, z) \in \text{dom}(\partial_u^- \mathcal{E}). \end{aligned} \quad (4.33)$$

In order to investigate Hyp. 4.12, consider  $u_n \rightharpoonup u$  in  $L^2(\Omega)$  with  $\sup_n \mathcal{G}(u_n, z_n) \leq C$ . Since  $\mathcal{E}(t, u, z) \geq c \int_{\Omega} |\nabla u|^2 dx - C$  for some  $c, C > 0$ , we conclude  $u_n \rightharpoonup u$  in  $H^1(\Omega)$ , hence  $u_n \rightarrow u$  in  $L^{2^*-\epsilon}(\Omega)$  for all  $\epsilon \in (0, 2^* - 1]$ . Thus,  $u_n^3 - u_n \rightharpoonup u^3 - u$  in  $L^2(\Omega)$ . By continuity of  $f$ ,  $t_n \rightarrow t$  implies  $f(t_n) \rightarrow f(t)$  in  $L^2(\Omega)$ . From  $\xi_n \rightharpoonup \xi$  in  $L^2(\Omega)$  we infer that  $(-\text{div}(g(z_n)\nabla u_n))_n$  is bounded in  $L^2(\Omega)$ , hence it weakly converges to some  $\zeta$  in  $L^2(\Omega)$ . We now aim to show that  $\zeta = -\text{div}(g(z)\nabla u)$ . For this, we directly employ the definition of subdifferential:

$$\begin{aligned} \int_{\Omega} \frac{g(z)}{2} |\nabla v|^2 dx - \int_{\Omega} \frac{g(z)}{2} |\nabla u|^2 dx &\geq \limsup_{n \rightarrow \infty} \left( \int_{\Omega} \frac{g(z_n)}{2} |\nabla v|^2 dx - \int_{\Omega} \frac{g(z_n)}{2} |\nabla u_n|^2 dx \right) \\ &\geq \liminf_{n \rightarrow \infty} \langle -\text{div} g(z_n)\nabla u_n, v - u_n \rangle_{\mathbf{V}} = \langle \zeta, v - u \rangle_{\mathbf{V}}. \end{aligned} \quad (4.34)$$

For the first estimate we have employed the weak sequential lower semicontinuity of the functional, the second inequality stems from the definition of the subdifferential, while the last equality is due to the convergences  $-\text{div} g(z_n)\nabla u_n \rightharpoonup \xi$  in  $L^2(\Omega)$  and  $u_n \rightarrow u$  strongly in  $L^2(\Omega)$  thanks to  $\mathbf{U} \Subset \mathbf{V}$ . Altogether, the above relations give  $\zeta = -\text{div}(g(z)\nabla u)$ . The just established convergence and closedness relations imply  $\xi = -\text{div}(g(z)\nabla u) + u^3 - u - f(t)$ , as well as  $\int_{\Omega} \frac{g(z_n)}{2} |\nabla u_n|^2 dx \rightarrow \int_{\Omega} \frac{g(z)}{2} |\nabla u|^2 dx$ , whence  $u_n \rightarrow u$  in  $H^1(\Omega)$  strongly, hence the convergence of the energies. We have thus established the stronger property (4.24b), and thus closedness condition (4.12), as well.

Now we consider  $\mathbf{V} = \mathbf{U} = H^1(\Omega)$ . Due to the quadratic nature of the leading order term in the energy functional we immediately find that  $u_n \rightharpoonup u$  in  $\mathbf{V}$  implies that  $-\text{div}(g(z_n)\nabla u_n) \rightharpoonup -\text{div}(g(z)\nabla u)$  in  $\mathbf{V}^*$ , hence, the validity of the closedness condition (4.24a). But since we here in general cannot infer the strong convergence of  $u_n \rightarrow u$  in  $\mathbf{V}$ , the stronger form (4.24b) is not valid in this context.

It can also be verified that, still with  $\mathbf{V} = \mathbf{U} = H^1(\Omega)$ , closedness Hypothesis 4.8 in its weaker form (4.13a) is valid with the energy from (4.31). Furthermore, thanks to the limsup-condition contained therein, also taking into account that the functionals  $\mathcal{F}_n(u) := \int_0^T \int_{\Omega} \frac{g(z_n)}{2} |\nabla u|^2 dx dt$  Mosco-converge (cf. [Att84]) in  $L^2(0, T; H^1(\Omega))$  to the functional  $\mathcal{F}(u) := \int_0^T \int_{\Omega} \frac{g(z)}{2} |\nabla u|^2 dx dt$ , we find that, for sequences  $(u_n, z_n)$  in the conditions of (4.13a), there holds  $\int_{\Omega} \frac{g(z_n(t))}{2} |\nabla u_n(t)|^2 dx \rightarrow \int_{\Omega} \frac{g(z(t))}{2} |\nabla u(t)|^2 dx$  for almost all  $t \in (0, T)$  (cf. also the argument from (4.34)). In turn, this implies that  $u_n(t) \rightarrow u(t)$  strongly in  $\mathbf{V}$ . For  $\Omega \subset \mathbb{R}^d$  with  $d = 2, 3, 4$  we also find that  $|u_n(t)|^4 \rightarrow |u(t)|^4$  strongly in  $L^1(\Omega)$ , which yields energy convergence. Therefore, closedness condition (4.13b) hold true.

Observe that in the above example, the linear character of the higher order term  $-\text{div}(g(z)\nabla u)$  contributing to the Fréchet subdifferential  $\partial_u^- \mathcal{E}$  has played a major role.

In the next example, we generalize the class of energy functionals to identify a wider setting in which Hypothesis 4.8 is satisfied. For simplicity we particularize the discussion to the cases  $\beta(A) = \frac{1}{p}|A|^p$  and  $W(u) = (u^2 - 1)^2/4$ , but it could be extended to a uniformly convex  $\beta$  with  $p$ -growth, cf. condition (6.4) in Sec. 6.1 ahead, and to a  $\lambda$ -convex potential  $W$  complying with suitable growth conditions. Since Hyp. 4.8 comes into play in the case the compact embedding  $\mathbf{U} \Subset \mathbf{V}$  is no longer required, we will focus on the case

$\mathbf{V} = H^1(\Omega)$  and  $\mathbf{U} \subset \mathbf{V}$  continuously, but not compactly. The example also illustrates that Hyp. 4.12 is in general not satisfied.

**Example 4.17.** We take  $\mathbf{V} = H^1(\Omega)$  and, for  $p \geq 2$  (cf. (4.29b))

$$\mathcal{E}(t, u, z) = \int_{\Omega} \left( \frac{g(z)}{p} |\nabla u|^p + \frac{1}{4}(u^2 - 1)^2 \right) dx - \langle f(t), u \rangle_{W^{1,p}(\Omega)} \quad \text{if } u \in \mathbf{D}, \text{ cf. (4.28),}$$

and  $+\infty$  otherwise. We choose the spaces  $\mathbf{X}$  and  $\mathbf{Z}$  as in (4.32). It can be checked that, for  $(t, u, z) \in [0, T] \times \mathbf{D} \times L^1(\Omega)$

$$\begin{aligned} \partial_u^- \mathcal{E}(t, u, z) \neq \emptyset &\Leftrightarrow -\operatorname{div}(g(z)|\nabla u|^{p-2}\nabla u) + u^3 - u - f(t) \in H^1(\Omega)^* \\ \text{and in that case } \partial_u^- \mathcal{E}(t, u, z) &= \{-\operatorname{div}(g(z)|\nabla u|^{p-2}\nabla u) + u^3 - u - f(t)\}. \end{aligned}$$

To verify Hypothesis 4.8, observe that, for a sequence  $u_n \xrightarrow{*} u$  in  $L^\infty(0, T; W^{1,p}(\Omega)) \cap H^1(0, T; H^1(\Omega))$ , whence  $u_n \rightarrow u$  strongly in  $C^0([0, T]; W^{1-\epsilon,p}(\Omega))$  for all  $\epsilon \in (0, 1)$ , there holds

$$u_n^3 - u_n \rightarrow u^3 - u \quad \text{in } L^2((0, T) \times \Omega). \quad (4.35)$$

Furthermore, let  $(t_n)_n$  a sequence in the conditions of (4.13a). From  $f \in C^1([0, T]; W^{1,p}(\Omega)^*)$  it follows that

$$f(t_n) \rightarrow f(t) \quad \text{in } L^\infty(0, T; W^{1,p}(\Omega)^*). \quad (4.36)$$

Finally, for a sequence  $(z_n)_n$  converging to  $z$  as indicated in (4.13a), by the compact embedding  $W^{1,q}(\Omega) \Subset L^q(\Omega)$  we have  $z_n(t) \rightarrow z(t)$  in  $L^q(\Omega)$  for all  $t \in [0, T]$  and hence, since  $g \in \text{BC}(\mathbb{R})$ , we have  $g(z_n) \rightarrow g(z)$  in  $L^\sigma((0, T) \times \Omega)$  for all  $1 \leq \sigma < \infty$ .

Now, let us consider the sequence  $\xi_n = -\operatorname{div}(g(z_n)|\nabla u_n|^{p-2}\nabla u_n) + u_n^3 - u_n - f(t_n)$ , weakly converging to some  $\xi$  in  $L^2(0, T; \mathbf{V}^*)$ . In view of (4.35) and (4.36),  $-\operatorname{div}(g(z_n)|\nabla u_n|^{p-2}\nabla u_n)$  weakly converges in  $L^2(0, T; W^{1,p}(\Omega)^*)$  to  $\zeta := \xi - W'(u) + f(t)$ . We now aim to show that  $\zeta = -\operatorname{div}(g(z)|\nabla u|^{p-2}\nabla u)$ . For this, we use the condition  $\limsup_{n \rightarrow \infty} \int_0^T \langle \xi_n, u_n \rangle_{\mathbf{V}} dt \leq \int_0^T \langle \xi, u \rangle_{\mathbf{V}} dt$  which, in view of (4.35) and of (4.36), in particular implies that

$$\limsup_{n \rightarrow \infty} \int_0^T \langle -\operatorname{div}(g(z_n)|\nabla u_n|^{p-2}\nabla u_n), u_n \rangle_{W^{1,p}(\Omega)} dt \leq \int_0^T \langle \zeta, u \rangle_{W^{1,p}(\Omega)} dt. \quad (4.37)$$

Taking into account that the functionals  $\mathcal{F}_n(u) := \int_0^T \int_{\Omega} \frac{g(z_n)}{p} |\nabla u|^p dx dt$  MOSCO-converge (cf. [Att84]) in  $L^p(0, T; W^{1,p}(\Omega))$  to  $\mathcal{F}(u) := \int_0^T \int_{\Omega} \frac{g(z)}{p} |\nabla u|^p dx dt$ , thanks to the lim sup inequality in (4.37) it is sufficient to conclude that  $\zeta = -\operatorname{div}(g(z)|\nabla u|^{p-2}\nabla u)$  almost everywhere in  $(0, T)$ . Hence,  $\xi = -\operatorname{div}(g(z)|\nabla u|^{p-2}\nabla u) + u^3 - u - f(t)$  a.e. in  $(0, T)$ , which concludes the proof of (4.13a).

Furthermore, from (4.37) we also infer that  $\int_0^T \int_{\Omega} \frac{g(z_n)}{p} |\nabla u_n|^p dx dt \rightarrow \int_0^T \int_{\Omega} \frac{g(z)}{p} |\nabla u|^p dx dt$ , whence the convergence of the energies in (4.13b).

But for  $p > 2$  observe that Hyp. 4.12 does not hold: Due to the nonlinear character of the higher order term, we are no longer in the position to directly infer from  $-\operatorname{div}(g(z_n)|\nabla u_n|^{p-2}\nabla u_n) \rightharpoonup \zeta$  in  $\mathbf{V}^*$  and  $u_n \rightharpoonup u$  in  $\mathbf{V}$  that  $\zeta = -\operatorname{div}(g(z)|\nabla u|^{p-2}\nabla u)$ .

With the next example, let us now turn to Hypothesis 4.6, which is assumed if  $\mathbf{U} \Subset \mathbf{V}$ .

**Example 4.18.** If  $\mathbf{V} = L^2(\Omega)$  and  $\mathbf{Z} = L^1(\Omega)$ , to verify the validity of Hypothesis 4.6, for every  $\tilde{u} \in \mathbf{D}_u \subset W^{1,p}(\Omega)$ , it is sufficient to take the recovery sequence  $\tilde{u}_n \equiv \tilde{u}$ . Then, for  $t_n \rightarrow t$  and  $z_n \rightarrow z$  in  $L^1(\Omega)$  we have  $g(z_n) \rightarrow g(z)$  in  $L^q(\Omega)$  for all  $q \in [1, \infty)$ , and by dominated convergence  $\int_{\Omega} g(z_n)\beta(\nabla \tilde{u}) dx \rightarrow \int_{\Omega} g(z)\beta(\nabla \tilde{u}) dx$ . By the continuity of  $f$ , we have  $\mathcal{E}(t_n, \tilde{u}, z_n) \rightarrow \mathcal{E}(t, \tilde{u}, z)$  and (4.11) ensues. Thanks to Lemma 4.7 closedness of the subdifferential and energy convergence ensue.

We also refer to Section 6.3 for further examples on the closedness hypotheses, also in combination with the other hypotheses.

5. EXISTENCE RESULTS FOR THE DAMPED INERTIAL SYSTEM  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ 

In this section we address the case with inertia, and investigate the weak solvability of the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ , supplemented with the initial conditions

$$u(0) = u_0, \quad u'(0) = v_0, \quad z(0) = z_0 \quad (5.1)$$

with  $(u_0, z_0) \in D_u \times D_z$  and  $v_0 \in \mathbf{W}$ . Combining the assumptions and techniques from Sections 4.1 and 4.2, we will obtain the existence of weak semistable energetic solutions for a general convex  $\mathcal{V}$  with superlinear growth, cf. Thm. 5.4, and of semistable energetic solutions for a quadratic dissipation potential  $\mathcal{V}$ , cf. Thm. 5.6.

**Time discretization.** We construct our discrete solutions via the following scheme

**Problem 5.1.** Let  $(u_\tau^0, z_\tau^0) := (u_0, z_0) \in D_u \times D_z$  and set

$$u_\tau^{-1} := u_0 - \tau v_0, \quad (5.2)$$

and for  $n = 1, \dots, N_\tau$ , find

$$z_\tau^n \in \underset{z \in \mathbf{Z}}{\text{Argmin}} \left( \tau \mathcal{R} \left( \frac{z - z_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, u_\tau^{n-1}, z) \right), \quad (5.3a)$$

$$u_\tau^n \in \underset{u \in \mathbf{V}}{\text{Argmin}} \left( \frac{1}{\tau^2} \|u - 2u_\tau^{n-1} + u_\tau^{n-2}\|_{\mathbf{W}}^2 + \tau \mathcal{V} \left( \frac{u - u_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, u, z_\tau^n) \right). \quad (5.3b)$$

Observe that Euler-Lagrange equation for the minimum problem (5.3b) now reads (taking into account that  $\mathbf{W} = \mathbf{W}^* \subset \mathbf{V}^*$ , cf. the proof of Prop. 7.7 for more details)

$$\frac{u_\tau^n - 2u_\tau^{n-1} + u_\tau^{n-2}}{\tau^2} + \partial \mathcal{V} \left( \frac{u - u_\tau^{n-1}}{\tau} \right) + \partial_u^- \mathcal{E}(t_n, u_\tau^n, z_\tau^n) \ni 0 \quad \text{in } \mathbf{V}^*. \quad (5.4)$$

We will denote by  $v_\tau$  the piecewise linear interpolant of the values  $((u_\tau^n - u_\tau^{n-1})/\tau)_{n=1}^{N_\tau}$ , namely

$$v_\tau(t) = \frac{t - t_{n-1}}{\tau} \frac{u_\tau^n - u_\tau^{n-1}}{\tau} + \frac{t_n - t}{\tau} \frac{u_\tau^{n-1} - u_\tau^{n-2}}{\tau} \quad \text{for } t \in (t_{n-1}, t_n]. \quad (5.5)$$

Therefore,  $v_\tau'(t) = \frac{1}{\tau^2} (u_\tau^n - u_\tau^{n-1} - (u_\tau^{n-1} - u_\tau^{n-2})) = \frac{1}{\tau^2} (u_\tau^n - 2u_\tau^{n-1} + u_\tau^{n-2})$  for all  $t \in (t_{n-1}, t_n]$ , and (5.4) rephrases as

$$v_\tau'(t) + \partial \mathcal{V}(v_\tau'(t)) + \bar{\xi}_\tau(t) \ni 0 \quad \text{with } \bar{\xi}_\tau(t) \in \partial_u^- \mathcal{E}(\bar{v}_\tau(t), \bar{u}_\tau(t), \bar{z}_\tau(t)) \quad \text{for a.a. } t \in (0, T). \quad (5.6)$$

We now introduce and motivate the two enhanced conditions on the Fréchet subdifferential of  $\mathcal{E}$  that we will have to impose in our existence results for  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ .

**Enhanced Fréchet subdifferentiability and subgradient estimate.** In order to recover the discrete version of the energy-dissipation inequality (3.4), we shall have to resort to a slightly stronger and more general version of the uniform Fréchet subdifferentiability property from Hypothesis 4.2. Indeed, with (5.7) below, on the one hand we require estimate (4.6) in Hyp. 4.2 to hold with a constant independent of the energy sublevel. On the other hand, we replace  $\|\cdot\|_{\mathbf{V}}^2$  by  $\|\cdot\|_{\mathbf{W}\mathcal{V}(\cdot)^\nu}$ , with an exponent  $\nu \in (0, 1)$  which we have to envisage as related to the growth properties of  $\mathcal{V}$ . For instance, in the case  $\mathcal{V}(v) = \frac{1}{2}\|v\|_{\mathbf{V}}^2$ , then we will have  $\nu = \frac{1}{2}$ , cf. also the forthcoming Example 5.10. Again, in the case  $\mathcal{V}(v) = \frac{1}{2}\|v\|_{\mathbf{V}}^2$ , taking into account that  $\mathbf{V} \subset \mathbf{W}$  continuously, we are in fact strengthening (4.6), cf. Remark 4.3.

**Hypothesis 5.2** (Enhanced Fréchet subdifferentiability). *There exist  $\nu \in (0, 1)$  and a constant  $\Lambda > 0$  such that*

$$\begin{aligned} \forall t \in [0, T], \quad \forall (u, z), (v, z) \in D_u \times D_z \quad \forall \xi \in \partial_u^- \mathcal{E}(t, u, z) : \\ \mathcal{E}(t, v, z) - \mathcal{E}(t, u, z) \geq \langle \xi, v - u \rangle_{\mathbf{V}} - \Lambda \|v - u\|_{\mathbf{W}\mathcal{V}(v-u)^\nu}. \end{aligned} \quad (5.7)$$

In addition, observe that in (5.6) the inertial term makes it necessary to estimate the term  $\bar{\xi}_\tau \in \partial_u^- \mathcal{E}(\cdot, \bar{u}_\tau, \bar{z}_\tau)$  independently, since no comparison estimate is possible. In this direction, the following condition requires that the  $\mathbf{V}^*$ -norm of the elements in the Fréchet subdifferential  $\partial_u^- \mathcal{E}$  to be estimated by the energy itself. As we will see later on, this condition e.g. holds for the energy functional in a model for adhesive contact in visco-elastic materials. This will also determine the spatial regularity for  $u''$ , with  $u$  the semistable energetic solution to the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$  arising in the time-continuous limit of scheme (5.3).

**Hypothesis 5.3** (Subgradient estimate). *There exist constants  $C_3, C_4, C_5 > 0$  and  $\sigma \in [1, \infty)$  such that*

$$\|\xi\|_{\mathbf{V}^*}^\sigma \leq C_3 \mathcal{E}(t, u, z) + C_4 \|u\|_{\mathbf{V}} + C_5 \quad \text{for all } \xi \in \partial_u^- \mathcal{E}(t, u, z) \text{ and all } (t, u, z) \in [0, T] \times D_{\mathbf{u}} \times D_{\mathbf{z}}. \quad (5.8)$$

A discussion of Hypothesis 5.3 in terms of two examples is given at the end of this section.

**An existence result for general  $\mathcal{V}$ .** We now give our first result for system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ , stating the existence of *weak semistable energetic solutions*. In addition to the basic conditions from Section 2 and to Hypotheses 5.2 & 5.3, we will also need to enforce a suitable closedness condition of  $\partial_u^- \mathcal{E}$  in order to pass to the time-continuous limit and identify the elements in the Fréchet subdifferential  $\partial_u^- \mathcal{E}$ . This will be done by

- either directly requiring Hypothesis 4.12;
- or, by imposing  $\mathbf{U} \Subset \mathbf{V}$ . Then, the desired closedness property for  $\partial_u^- \mathcal{E}$  will derive from combining Hypotheses 4.6 and the uniform Fréchet subdifferentiability 5.2. Indeed, using that  $\mathcal{V}$  is continuous, it is not difficult to check that the proof of Lemma 4.7 goes through in this setting as well, guaranteeing the closedness property (4.12).

**Theorem 5.4** (Weak semistable energetic solutions to  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ ,  $\mathcal{V}$  general). *Assume Hypotheses 2.2, 2.3, 2.5, 2.7 and, in addition, Hypotheses 5.2 and 5.3. Moreover, assume*

- (1) *either Hypothesis 4.12,*
- (2) *or,  $\mathbf{U} \Subset \mathbf{V}$  with Hypothesis 4.6.*

*Then, for every sequence  $\tau_k \downarrow 0$  as  $k \rightarrow \infty$  there exist a (not relabeled) subsequence, and functions  $(u, z)$  as in (3.1) such that, in addition,  $u \in W^{2,1}(0, T; \mathbf{V}^*)$ , and convergences (4.15) hold. Moreover,*

$$v'_{\tau_k} \xrightarrow{*} u' \quad \text{in } L^\infty(0, T; \mathbf{W}), \quad (5.9a)$$

$$v_{\tau_k}(t), u'_{\tau_k}(t) \rightarrow u'(t) \quad \text{in } \mathbf{W} \quad \text{for all } t \in [0, T], \quad (5.9b)$$

$$v'_{\tau_k} \rightarrow u'' \quad \text{in } L^1(0, T; \mathbf{V}^*), \quad (5.9c)$$

*and the pair  $(u, z)$  is a weak semistable energetic solution to the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ , fulfilling the initial conditions (5.1) with initial data  $(u_0, z_0) \in D_{\mathbf{u}} \times D_{\mathbf{z}}$  fulfilling (4.14).*

*Furthermore, if Hypothesis 4.12 holds with the additional continuity condition (4.24b), or under Hypothesis 4.6 in the case  $\mathbf{U} \Subset \mathbf{V}$ , then we have the additional convergence (4.19), as well as (4.18) in the case of Hyp. 4.6 with  $\mathbf{U} \Subset \mathbf{V}$ .*

Observe, however, that, even in the cases in which we get the energy convergence (4.19), we cannot conclude that  $(u, z)$  is an enhanced weak energetic solution, cf. Remark 7.9 ahead. Still, we have the analogue of Corollary 4.14.

**Corollary 5.5.** *Under the assumptions of Theorem 5.4, if in addition  $\mathcal{E}$  complies with the smoothness properties (3.15), then  $(u, z)$  from Theorem 5.4 is a semistable energetic solution, complying with the energy-dissipation balance along every subinterval  $[s, t] \subset [0, T]$ .*

**An existence result for  $\mathcal{V}$  quadratic.** In the case of a *quadratic* dissipation potential  $\mathcal{V}$ , we will be able to show the existence of *semistable energetic solutions* to  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$  in the special setting that

$$\mathbf{V} \Subset \mathbf{W}. \quad (5.10)$$

This will guarantee sufficient compactness information for (the sequence approximating)  $u'$  to allow for the limit passage in the (discrete) viscous equation. Clearly, in view of the compact embedding  $\mathbf{U} \Subset \mathbf{W}$  from (2.1b), (5.10) is for example fulfilled in the case in which  $\mathbf{V} = \mathbf{U}$ , cf. Example 2.8. Therefore, to get the closedness of  $\partial_u^- \mathcal{E}$  we will resort to the closedness condition à la Minty from Hypothesis 4.8.

**Theorem 5.6** (Semistable energetic solutions for  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ ,  $\mathcal{V}$  quadratic). *Assume Hypotheses 2.2, 2.3, 2.5, 2.7, and let  $\mathcal{V}$  be quadratic, cf. (4.1). In addition, assume Hypotheses 4.8, 5.2, 5.3, and (5.10).*

*Then, for every sequence  $\tau_k \downarrow 0$  as  $k \rightarrow \infty$  starting from initial data  $(u_0, z_0)$  fulfilling (4.14) there exist a (not relabeled) subsequence, and functions  $(u, z)$  as in (3.1) such that, in addition,  $u \in H^1(0, T; \mathbf{V}) \cap H^2(0, T; \mathbf{V}^*)$  and convergences (4.15) hold, as well as (5.9). Moreover,*

$$v'_{\tau_k} \rightarrow u'' \quad \text{in } L^2(0, T; \mathbf{V}^*), \quad u'_{\tau_k} \rightarrow u' \quad \text{in } L^p(0, T; \mathbf{W}) \quad \text{for all } 1 \leq p < \infty \quad (5.11)$$

and the pair  $(u, z)$  is a semistable energetic solution to the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ , fulfilling (5.1).

Finally, if Hyp. 4.8 holds in the stronger form (4.13b), then the energy-dissipation inequality holds in the form

$$\begin{aligned} & \frac{1}{2} \|u'(t)\|_{\mathbf{W}}^2 + \int_s^t \mathcal{V}(u'(r)) + \mathcal{V}^*(-\xi(r) - u''(r)) dr + \text{Var}_{\mathcal{R}}(z; [s, t]) + \mathcal{E}(t, u(t), z(t)) \\ & \leq \frac{1}{2} \|u'(s)\|_{\mathbf{W}}^2 + \mathcal{E}(s, u(s), z(s)) + \int_s^t \partial_t \mathcal{E}(r, u(r), z(r)) dr \quad \text{for all } t \in (0, T] \text{ for } s = 0, \text{ and a.a. } s \in (0, t), \end{aligned} \quad (5.12)$$

with  $\xi(t)$  a selection in  $\partial_u^- \mathcal{E}(t, u(t), z(t))$  also fulfilling

$$u''(t) + \partial \mathcal{V}(u'(t)) + \xi(t) = 0 \quad \text{for a.a. } t \in (0, T), \quad (5.13)$$

and  $(u, z)$  is thus an enhanced semistable energetic solution.

The proofs of Theorems 5.4 and 5.6 will be carried out in Sec. 7.3.

**Remark 5.7.** In Remark 7.8 we will check that for  $\mathcal{V}$  quadratic it would indeed be possible to work under a weaker version of the Fréchet subdifferentiability from Hypothesis 5.2, namely

$$\begin{aligned} & \forall t \in [0, T], \forall (u, z), (v, z) \in D_u \times D_z \quad \forall \xi \in \partial_u^- \mathcal{E}(t, u, z) : \\ & \mathcal{E}(t, v, z) - \mathcal{E}(t, u, z) \geq \langle \xi, v - u \rangle_{\mathbf{V}} - \Lambda \|v - u\|_{\mathbf{V}}^2, \end{aligned} \quad (5.14)$$

i.e. the former (4.6), but with a constant *independent* of the energy sublevel.

**Remark 5.8.** In the case of quadratic  $\mathcal{V}$ , it would also be possible to weaken the subgradient estimate in Hyp. 5.3 by replacing  $\|\xi\|_{\mathbf{V}^*}$  with  $\|\xi\|_{\mathbf{U}^*}$ , with  $\xi$  an element in the Fréchet subdifferential  $\partial_u^- \mathbf{U} \mathcal{E}(t, u, z)$  of  $\mathcal{E}$  in the duality between  $\mathbf{U}^*$  and  $\mathbf{U}$ . Accordingly, taking into account that for every  $v \in \mathbf{V}$ ,  $\partial \mathcal{V}(v) \subset \mathbf{V}^* \subset \mathbf{U}^*$ , (3.2) could be formulated in a weaker way as a subdifferential inclusion in  $\mathbf{U}^*$ , and one would have  $u \in H^2(0, T; \mathbf{U}^*)$ .

The proof of this extension follows the same lines as the argument for Thm. 5.6, with suitable modifications such as the closedness argument, to be formulated in the duality between  $\mathbf{U}^*$  and  $\mathbf{U}$ .

**Remark 5.9** (Comparison with the results by E. Emmrich & coworkers). In the papers [ET11, EŠ11, EŠ13], E. EMMRICH and coauthors have obtained a series of deep results on the existence of solutions (whose basics were already established in the seminal paper [LS65]), and on the full discretization (in time *and* space) for this class of second order nonlinear evolution equations

$$u'' + Au' + Bu = f \quad \text{in } V^*, \text{ a.e. in } (0, T). \quad (5.15)$$

Here,  $V = V_A \cap V_B$  and  $V_A, V_B$  are the separable reflexive Banach spaces on which the two operators  $A$  and  $B$  are defined. The authors neither suppose  $V_A \subset V_B$  (continuously), nor do they require  $V_B \subset V_A$ , thus allowing for a wide class of applications, although (5.15) is not coupled to the rate-independent evolution of an additional internal variable  $z$ .

In [ET11, EŠ11] the following assumptions are made on the *time-dependent* operators  $A$  and  $B$ : It is required that  $A : V_A \rightarrow V_A^*$  has a main part  $d$ -monotone, hemicontinuous, (cf. e.g. [Rou05, Chap. 2, Definitions 2.1, 2.3]) and satisfying a certain growth condition, that is perturbed by a non-monotone but locally Hölder continuous operator, with suitable growth. The main part of the operator  $B : V_B \rightarrow V_B^*$  is linear, symmetric, and strongly positive, and it is also perturbed by a locally Hölder continuous operator, with suitable growth. The analysis in [ET11, EŠ11] can thus be compared to ours in the (particular) case in which the energy functional as a function of  $u$  is quadratic up to a lower order perturbation. Observe that, while we restrict to the subdifferential of a possibly *nonsmooth, but convex* dissipation potential  $\mathcal{V}$ , the operator  $A$  in (5.15) could be the differential of a convex smooth potential with a nonconvex, but lower order, perturbation.

Recently, in [EŠ13], the authors instead consider the case in which  $B$  is the Gâteaux differential of an energy bounded from below, with superlinear growth at infinity in the sense of (2.12a). It is assumed that  $B$  maps bounded sets of  $V_B$  into bounded sets of  $V_B^*$  and that it is demicontinuous. However, in this case the analysis is restricted to the case in which  $A$  is the linear operator associated with a bilinear form inducing an inner product on  $V_A$  and a norm  $\|\cdot\|_A$  equivalent to  $\|\cdot\|_{V_A}$ . This corresponds to our choice of a *quadratic* dissipation

potential  $\mathcal{V}$ . It has to be noted that in [EŠ13] the operators  $A$  and  $B$  are required to satisfy a condition of *Andrews-Ball* type, namely that there exists  $\lambda \geq 0$  such that the operator  $(B + \lambda A) : V \rightarrow V^*$  is monotone:

$$\exists \lambda \geq 0 \quad \forall v, w \in V \quad \langle Bv - Bw, v - w \rangle_V \geq -\lambda \|v - w\|_A^2. \quad (5.16)$$

This exactly corresponds to the uniform Fréchet subdifferentiability condition (5.14).

**Examples for the subgradient estimate in Hypothesis 5.3.** It is not difficult to verify that Hypothesis 5.3 is, e.g., satisfied by the energy in Example 4.16, with the space  $\mathcal{V} = H^1(\Omega)$ ; recall that, in the latter space the closedness from Hyp. 4.12 is also fulfilled.

The following example has a structure similar to Ex. 4.16, but it features a *nonsmooth* double-well term  $W$ , with a kink at  $|u| = 2$ . Again, to make calculations more transparent we neglect the  $z$ -dependence.

**Example 5.10.** Let us consider  $W_2 : \mathbb{R} \rightarrow \mathbb{R}$

$$W_2(u) := \begin{cases} a(1 - u^2)^2 & \text{if } |u| \leq 2, \\ b(1 - u^2)^2 + 9(a - b) & \text{if } |u| > 2, \end{cases} \quad \text{with constants } 0 < a < b. \quad (5.17)$$

It is a  $\Lambda$ -convex function for any  $\Lambda \geq 2b$ , i.e., for all  $u, \tilde{u} \in \mathbb{R}^d$ , for all  $\theta \in [0, 1]$

$$W_2(\theta u + (1 - \theta)\tilde{u}) \leq \theta W_2(u) + (1 - \theta)W_2(\tilde{u}) + \Lambda\theta(1 - \theta)|u - \tilde{u}|^2. \quad (5.18)$$

Its Fréchet subdifferential  $\partial^- W_2 : \mathbb{R} \rightrightarrows \mathbb{R}$  is given by

$$\partial^- W_2(u) = \begin{cases} \{-4au(1 - u^2)\} & \text{if } |u| < 2, \\ \{-4\alpha u(1 - u^2), \alpha \in [a, b]\} & \text{if } |u| = 2, \\ \{-4bu(1 - u^2)\} & \text{if } |u| > 2. \end{cases}$$

and every  $\xi \in \partial^- W_2(u)$  satisfies

$$|\xi|^{4/3} \leq (4\alpha(|u| + |u|^3))^{4/3} \leq 48\alpha|u|^4 \leq \frac{48^{4/3}\alpha^{1/3}}{\min\{a, 1\}} (W_2(u) + 9(b - a) + 2bu^2). \quad (5.19)$$

Let now  $\mathbf{V} := L^4(\Omega)$  and  $\mathbf{W} = L^2(\Omega)$ , with  $\Omega \subset \mathbb{R}^d$  a bounded Lipschitz domain, and let  $\mathcal{W}_2 : \mathbf{V} \rightarrow [0, +\infty)$  be given by  $\mathcal{W}_2(u) := \int_{\Omega} W_2(u) dx$ . Then, (5.19) leads to subdifferential estimate (5.8). From convexity inequality (5.18) we obtain

$$\begin{aligned} \mathcal{W}_2(\theta u + (1 - \theta)\tilde{u}) &\leq \theta \mathcal{W}_2(u) + (1 - \theta)\mathcal{W}_2(\tilde{u}) + \Lambda\theta(1 - \theta)\|u - \tilde{u}\|_{\mathbf{W}}^2 \\ &\leq \theta \mathcal{W}_2(u) + (1 - \theta)\mathcal{W}_2(\tilde{u}) + \Lambda C_{\mathbf{V}\mathbf{W}}\theta(1 - \theta)\|u - \tilde{u}\|_{\mathbf{V}}^{1/4}, \end{aligned} \quad (5.20)$$

where the last estimate follows from the continuous embedding  $\|w\|_{\mathbf{W}} \leq C_{\mathbf{V}\mathbf{W}}\|w\|_{\mathbf{V}}$ . Therefore, the enhanced Fréchet subdifferentiability (5.7) ensues upon choosing  $\mathcal{V}(v) := \frac{1}{4} \int_{\Omega} |v|^4 dx$ , with  $\nu = \frac{1}{4}$ .

Observe that the corresponding energy augmented by a Dirichlet integral, i.e.  $\mathcal{E}(t, u) := \int_{\Omega} \frac{1}{2} |\nabla u|^2 + W_2(u) dx$  complies, in the context of the space  $\mathbf{U} = H_0^1(\Omega)$ , with Hypotheses 5.2 and 5.3 and, in addition, with Hypotheses 2.2, 2.7, and with the closedness à la Minty from Hyp. 4.8.

## 6. APPLICATIONS

In this section we collect some examples of mechanical systems with a mixed rate-dependent/rate-independent character, to which our existence Theorems 4.9, 4.13, 5.4, 5.6 apply.

### 6.1. Semistable energetic solutions to a class of rate-independent processes in visco-elastic solids.

Here we consider a class of rate-independent systems in a visco-elastic body  $\Omega \subset \mathbb{R}^d$ , also subject to inertia, in the frame of the theory of *generalized standard solids*. Their analysis was first addressed in [Rou09]. The variable governed by viscosity is the displacement  $\mathbf{u}$  (in a small-strain regime,  $\varepsilon(\mathbf{u}) := \frac{1}{2}(\nabla \mathbf{u} + \nabla \mathbf{u}^\top)$  denoting the small strain tensor), while the internal variable  $z$ , in general taking values in  $\mathbb{R}^m$ , describes some rate-independent process such as, e.g., plasticity with hardening, damage, or phase transformations in a solid.

**Energetics.** The energy has the form

$$\mathcal{E}(t, \mathbf{u}, z) := \int_{\Omega} \varphi(\varepsilon(\mathbf{u}), z, \nabla z) dx - \int_{\Omega} \mathbf{f} \cdot \mathbf{u} dx \quad (6.1)$$

where  $\mathbf{f}$  is a given external force and the energy density  $\varphi : \mathbb{R}_{\text{sym}}^{d \times d} \times \mathbb{R}^m \times \mathbb{R}^{m \times d} \rightarrow [0, +\infty)$  is generally assumed smooth, although in [Rou09, Rmk. 3.3] it is hinted that an extension to nonsmooth energies could be possible.

The “viscous” dissipation potential  $\mathcal{V}$  is quadratic and has the form

$$\mathcal{V}(\mathbf{v}) := \int_{\Omega} \frac{1}{2} \varepsilon(\mathbf{v}) : \mathbb{D} : \varepsilon(\mathbf{v}) \quad \text{with } \mathbb{D} \in \mathbb{R}_{\text{sym}}^{d \times d \times d \times d} \text{ positive definite,} \quad (6.2)$$

while the “rate-independent” potential  $\mathcal{R}$  is

$$\begin{aligned} \mathcal{R}(\zeta) &= \int_{\Omega} R(\zeta) dx \quad \text{with } R : \mathbb{R}^m \rightarrow [0, +\infty] \text{ convex, positively homogeneous of degree 1, and} \\ &\exists c > 0 \quad \forall \zeta \in \mathbb{R}^m : \quad R(\zeta) \geq c|\zeta|. \end{aligned} \quad (6.3)$$

The kinetic energy is  $\mathcal{K}(\mathbf{v}) = \frac{\varrho}{2} \int_{\Omega} |\mathbf{v}|^2 dx$  with  $\varrho \geq 0$ , but all the results from [Rou09] could be extended to the case of a spatially heterogeneous mass density, cf. (6.6) ahead. Therefore, our own Hypothesis 2.3 is satisfied with the spaces  $\mathbf{V} = H_{\Gamma_D}^1(\Omega; \mathbb{R}^d)$ , with  $\Gamma_D$  a portion of the boundary  $\partial\Omega$  on which zero Dirichlet boundary conditions are imposed, and  $\mathbf{Z} = L^1(\Omega; \mathbb{R}^m)$ , while  $\mathbf{W} = L^2(\Omega; \mathbb{R}^d)$ . Throughout this section, we will use the notation

$$W_{\Gamma_D}^{1,p}(\Omega; \mathbb{R}^d) := \{v \in W^{1,p}(\Omega, \mathbb{R}^d) : v = 0 \text{ on } \Gamma_D\}.$$

**Properties of the energy.** Concerning the energy density  $\varphi$ , ROUBÍČEK provides classes of sufficient conditions ensuring the existence of *semistable energetic* solutions. In fact, in [Rou09] other weak solvability notions are proposed and related existence results are obtained. Nonetheless, in the next lines we will only focus on the energetic concept and highlight the relations between ROUBÍČEK’s assumptions on  $\varphi$  for the existence result [Rou09, Prop. 5.3], and the abstract conditions for our own existence Thms. 4.9 and 5.6 (for  $\mathcal{V}$  quadratic).

First of all, it is required (cf. [Rou09, (12a)]) that  $\varphi(e, z, Z) \geq \tilde{c}(|e|^p + |Z|^q)$  for some  $\tilde{c} > 0$  and  $p, q > 1$ . This sets the spaces  $\mathbf{U}$  and  $\mathbf{X}$  in which the coercivity Hypothesis 2.7 is verified, namely

$$\mathbf{U} = W_{\Gamma_D}^{1,p}(\Omega; \mathbb{R}^d) \cap H_{\Gamma_D}^1(\Omega; \mathbb{R}^d) \quad \text{and} \quad \mathbf{X} = W^{1,q}(\Omega; \mathbb{R}^m).$$

It is not difficult to check that Hyp. 2.2 is also fulfilled.

The other basic condition (cf. [Rou09, (16)]) is that there exists  $\ell \geq 0$  such that the mapping  $(e, z, Z) \mapsto \varphi(e, z, Z) + \ell|e|^2$  is convex, which guarantees the validity of our uniform Fréchet differentiability Hyps. 4.2 & 5.2.

In [Rou09, Prop. 5.3] ROUBÍČEK also proposes three sets of conditions (cf. (66a)/(66b)/(66c) therein) that he alternatively uses to verify the mutual recovery sequence condition from Hyp. 2.5. More in detail,

- (a) either it is assumed ([Rou09, (66a)]) that  $R$  is continuous (and in particular takes values in  $[0, +\infty)$ ), in combination with the condition that  $\partial_e \varphi$  is *p-strongly monotone*, i.e.

$$\begin{aligned} &\exists \alpha > 0 \quad \forall (z, Z) \in \mathbb{R}^m \times \mathbb{R}^{m \times d} \quad \forall e, \tilde{e} \in \mathbb{R}_{\text{sym}}^{d \times d} : \\ &\alpha(|\tilde{e}|^{p-2} \tilde{e} - |e|^{p-2} e) : (\tilde{e} - e) \leq \partial_e \varphi(\tilde{e}, z, Z) - \partial_e \varphi(e, z, Z). \end{aligned} \quad (6.4)$$

Observe that (6.4) implies our Hyp. 4.8, whereas the combination of (6.4) and the continuity of  $R$  allows for the choice of the constant recovery sequence  $\tilde{z}_n = \tilde{z}$  in Hyp. 2.5;

- (b) or, (cf. [Rou09, (66b)]), in addition to (6.4) suitable growth properties of  $(z, Z) \mapsto \varphi(e, z, Z)$  and of  $(z, Z) \mapsto \partial_{(z,Z)} \varphi(e, z, Z)$  are required. This again ensures Hyp. 4.8, while the recovery sequence  $(\tilde{z}_n)_n$  for Hyp. 2.5 is constructed via the so-called *quadratic trick*, cf. e.g. [MR15, Lemma 3.5.3];
- (c) or it is supposed, cf. [Rou09, (66c)], that  $\varphi(e, z, Z) = \varphi_1(e, z, Z) + \varphi_2(z, Z)$  with  $(e, Z) \mapsto \varphi_1(e, z, Z)$  affine and fulfilling suitable growth conditions, and  $\varphi_2$  quadratic. This guarantees Hyps. 2.5 and 4.8.

**Existence of semistable energetic solutions.** All in all, we have verified that the class of material models in [Rou09] complies, in the case of a gradient system, i.e.  $\varrho = 0$ , with all the conditions in Theorem 4.9. Thus our own result applies, yielding the existence of *semistable energetic* solutions.

In the case with inertia, i.e.  $\varrho > 0$ , ROUBÍČEK additionally assumes a growth condition for  $\partial_e \varphi$  (cf. [Rou09, (21)]) that is akin to our own Hypothesis 5.3, although slightly weaker. Indeed, it leads to a weaker formulation



of the momentum equation than ours (5.13) (cf. Thm. 5.6), in which the inertial term is integrated by parts in time.

**6.2. Adhesive contact between visco-elastic materials with inertia.** We consider two visco-elastic bodies  $\Omega_+$  and  $\Omega_-$ , bonded along a prescribed contact surface  $\Gamma_C$ , during a timespan  $[0, T]$ . We will take as state variable the displacement field  $\mathbf{u} : (0, T) \times \Omega \rightarrow \mathbb{R}^d$  (with  $\Omega := \Omega_- \cup \Omega_+$ ), subject to viscosity and inertia, and denote by  $\llbracket \mathbf{u} \rrbracket$  its jump across  $\Gamma_C$ . Along the lines of the theory of generalized standard materials [HN75, Fré02b], the adhesiveness of the bonding is modeled by the *delamination variable*  $z : (0, T) \times \Gamma_C \rightarrow [0, 1]$ , describing the fraction of the effective molecular links between  $\Omega_+$  and  $\Omega_-$ , so that  $z(t, x) = 1$  ( $z(t, x) = 0$ , respectively), means fully intact (broken) bonding at time  $t \in (0, T)$  and at the material point  $x \in \Gamma_C$ . We postulate a *rate-independent* evolution for  $z$ . For a different approach we refer to, e.g., [KP11], where the motion of a (single) visco-elasto-plastic body, also subject to inertia, in contact with an elasto-plastic obstacle, is modeled with the aid of suitable hysteresis operators.

Following [Fré88], we model this phenomenon by the energy

$$\begin{aligned} \mathcal{E}(t, \mathbf{u}, z) := & \int_{\Omega \setminus \Gamma_C} \frac{1}{2} \varepsilon(\mathbf{u}) : \mathbb{C} : \varepsilon(\mathbf{u}) \, dx + \int_{\Gamma_C} \kappa z \|\llbracket \mathbf{u} \rrbracket\|^2 \, d\mathcal{H}^{d-1}(x) \\ & + \int_{\Gamma_C} (J(\llbracket \mathbf{u} \rrbracket) + I_{[0,1]}(z) - a_0 z) \, d\mathcal{H}^{d-1}(x) - \langle \mathbf{f}(t), \mathbf{u} \rangle_{H^1(\Omega; \mathbb{R}^d)}, \end{aligned} \quad (6.5)$$

where  $\mathcal{H}^{d-1}$  is the  $(d-1)$ -dimensional Hausdorff measure,  $\mathbb{C} \in \mathbb{R}^{d \times d \times d \times d}_{\text{sym}}$  the symmetric, positive definite elasticity tensor,  $\kappa$  a fixed positive constant modulating the adhesive contact contribution,  $J : \mathbb{R}^d \rightarrow [0, +\infty]$  a convex functional,  $a_0 \geq 0$  the specific energy stored by delamination, and  $\mathbf{f} \in W^{1,1}(0, T; H^1(\Omega \setminus \Gamma_C; \mathbb{R}^d)^*)$  a given external loading incorporating volume and surface forces acting on the Neumann part  $\Gamma_N$  of the boundary  $\partial\Omega$  (whereas on the Dirichlet part  $\Gamma_D$  null displacement is imposed). Moreover, in (6.5) and also below,  $I_A$  denotes the indicator function of a convex set  $A \subset \mathbb{R}$ , i.e.,  $I_A(\zeta) = 0$  if  $\zeta \in A$  and  $I_A(\zeta) = \infty$  if  $\zeta \in \mathbb{R} \setminus A$ . The dissipation potentials are

$$\mathcal{V}(\mathbf{v}) := \int_{\Omega \setminus \Gamma_C} \frac{1}{2} \varepsilon(\mathbf{v}) : \mathbb{D} : \varepsilon(\mathbf{v}) \, dx, \quad \mathcal{R}(\zeta) := \int_{\Gamma_C} R(\zeta) \, d\mathcal{H}^{d-1}(x)$$

where  $\mathbb{D} \in \mathbb{R}^{d \times d \times d \times d}_{\text{sym}}$  is the symmetric, positive definite viscosity tensor. The density of the 1-positively homogeneous potential  $\mathcal{R}$  is given by  $R(\zeta) = -a_1 \zeta + I_{(-\infty, 0]}(\zeta)$ , with  $a_1 \geq 0$  the specific energy dissipated by delamination. Inertial effects are encompassed in the model through the kinetic energy

$$\mathcal{K}(\mathbf{w}) := \int_{\Omega} \frac{\varrho(x)}{2} |\mathbf{w}|^2 \, dx, \quad (6.6a)$$

with  $\varrho$  a non-negative function in  $L^1(\Omega)$ , with  $\varrho \geq \bar{\varrho} > 0$  a.e. in  $\Omega$ , so that the underlying Hilbert space is

$$\mathbf{W} = \{ \mathbf{w} : \sqrt{\varrho} \mathbf{w} \in L^2(\Omega; \mathbb{R}^d) \}. \quad (6.6b)$$

The analysis of this adhesive contact model in the *fully* rate-independent case, i.e. when the flow rule for the adhesive parameter is coupled to the *static* momentum balance, goes back to [KMR06]. The existence of semistable energetic solutions to the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ , where the other ambient spaces are, naturally,  $\mathbf{V} = H^1_{\Gamma_D}(\Omega \setminus \Gamma_D; \mathbb{R}^d)$ , and  $\mathbf{Z} = L^1(\Gamma_C)$ , stems from the analysis in [RR11], also encompassing the coupling with thermal processes. It is indeed possible to show that the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$  in fact complies with all of the assumptions of Theorem 5.6. Here we will just confine the discussion to Hypothesis 5.3, highlighting a significant point in the analysis of contact systems with inertia.

The Fréchet subdifferential  $\partial_{\mathbf{u}}^- \mathcal{E}$  is given by  $\partial_{\mathbf{u}}^- \mathcal{E}(t, \mathbf{u}, z) = \{ D_{\mathbf{u}} \mathcal{E}(t, \mathbf{u}, z) \}$ , with

$$\langle D_{\mathbf{u}} \mathcal{E}(t, \mathbf{u}, z), \mathbf{v} \rangle_{H^1(\Omega; \mathbb{R}^d)} := \int_{\Omega \setminus \Gamma_C} \varepsilon(\mathbf{u}) : \mathbb{C} : \varepsilon(\mathbf{v}) \, dx + \int_{\Gamma_C} (\kappa z \llbracket \mathbf{u} \rrbracket \llbracket \mathbf{v} \rrbracket + \zeta \llbracket \mathbf{v} \rrbracket) \, d\mathcal{H}^{d-1}(x) - \langle \mathbf{f}(t), \mathbf{v} \rangle_{H^1(\Omega; \mathbb{R}^d)}$$

for all  $\mathbf{v} \in H^1(\Omega; \mathbb{R}^d)$ , where  $\zeta \in L^2(\Gamma_C)$  fulfills  $\zeta(x) \in \partial J(\llbracket \mathbf{u}(x) \rrbracket)$  for a.a.  $x \in \Gamma_C$ . Now, while the  $H^1(\Omega; \mathbb{R}^d)^*$ -norm of all the other contributions to  $D_{\mathbf{u}} \mathcal{E}(t, \mathbf{u}, z)$  is controlled by  $\mathcal{E}$  in the sense of (5.8), for the validity of Hyp. 5.3 the convex subdifferential  $\partial J : \mathbb{R}^d \rightrightarrows \mathbb{R}^d$  cannot be an unbounded operator. Therefore, the choice  $J(\llbracket \mathbf{u} \rrbracket) := I_{[0, +\infty)}(\llbracket \mathbf{u} \rrbracket \cdot \nu)$  ( $\nu$  denoting the unit normal to  $\Gamma_C$ , oriented from  $\Omega_+$  to  $\Omega_-$ ), is not admissible. Observe that such a choice would lead to a flow rule for the delamination variable encompassing the *Signorini contact* conditions, and in fact the existence of solutions to the adhesive contact system with Signorini conditions

and inertia has been a challenging problem in the literature on contact problems for a long time, cf. [RR11, Rmk. 5.3]. Recent progress has been done in [PS09]; see also [Pet12] for results in the numerical analysis of a (spatially one-dimensional) visco-elasto-dynamic contact problem with the Signorini conditions.

Instead, in order to comply with Hyp. 5.3, one can for instance take for  $J$  the Yosida approximation  $J_\epsilon(\mathbf{u}) := -\frac{1}{\epsilon}([\mathbf{u}] \cdot \nu)^-$ , with  $\epsilon > 0$ ,  $(\cdot)^-$  denoting the negative part. This corresponds to the so-called *normal compliance conditions*.

**6.3. Rate-independent damage in visco-elastic solids.** In this section we highlight some possible extensions that are included in our analytical results but not yet covered by the results by ROUBÍČEK. In particular, we will focus on a model for partial damage in visco-elastic solids at small strains, located in a bounded Lipschitz domain  $\Omega \subset \mathbb{R}^d$ , observed over the finite timespan  $[0, T]$ . Let us mention in advance that we will allow for a *nonsmooth* dependence  $u \mapsto \mathcal{E}(t, u, z)$  and for *unidirectionality* in the evolution of  $z$ , enforced by a dissipation potential with values in  $[0, +\infty]$ , combined with a BV-gradient regularization for  $z$ .

The state variables of the model under consideration are the displacement field  $\mathbf{u} : [0, T] \times \Omega \rightarrow \mathbb{R}^d$ , with  $\varepsilon(\mathbf{u}) = \frac{1}{2}(\nabla \mathbf{u} + \nabla \mathbf{u}^\top)$  the small strain tensor, and the damage variable  $z : [0, T] \times \Omega \rightarrow [0, 1]$ . In particular,  $z(t, x) = 1$  corresponds to the case that the material is fully intact at the point  $(t, x) \in [0, T] \times \Omega$ , while  $z(t, x) = 0$  refers to maximal damage at  $(t, x)$ . As is common in the modeling of rate-independent damage without healing, the 1-homogeneous dissipation potential is

$$\mathcal{R}(\zeta) := \int_{\Omega} R(\zeta) \, dx, \quad R(\zeta) := \begin{cases} |\zeta| & \text{if } \zeta \leq 0, \\ +\infty & \text{otherwise.} \end{cases} \quad (6.7)$$

The energy functional  $\mathcal{E} : [0, T] \times \mathbf{V} \times \mathbf{Z} \rightarrow (-\infty, +\infty]$  has the following structure

$$\mathcal{E}(t, \mathbf{u}, z) := \begin{cases} \int_{\Omega} g(z)W(\varepsilon(\mathbf{u})) \, dx - \langle \mathbf{f}(t), \mathbf{u} \rangle_{\mathbf{V}} + \mathcal{G}(z) & \text{if } (\mathbf{u}, z) \in \mathbf{U} \times \mathbf{X}, \\ +\infty & \text{otherwise.} \end{cases} \quad (6.8)$$

Here,  $\mathbf{f} \in C^1([0, T]; \mathbf{V}^*)$  is a given external loading. The space  $\mathbf{U}$ , specified in (6.13) ahead, shall encompass the homogeneous Dirichlet condition imposed on the Dirichlet boundary  $\Gamma_D$ . The energy contributions  $g(\cdot)W(\cdot)$  and  $\mathcal{G}(\cdot)$ , encompassing a gradient term in  $z$ , will have to be chosen such that the associated energy spaces comply with the topological setup (2.1) and (2.5). Here we want to point out that the latter condition includes the case of (the non-reflexive space)  $\mathbf{X} = \text{BV}(\Omega)$ . This in particular allows for a gradient regularization of the form

$$\mathcal{G} : \mathbf{X} \rightarrow [0, +\infty], \quad \mathcal{G}(z) := |Dz|(\Omega) + \int_{\Omega} I_{[0,1]}(z) \, dx, \quad (6.9)$$

with  $|Dz|(\Omega)$  the total variation of  $z$  in  $\Omega$ , and  $I_{[0,1]}$  the indicator function of  $[0, 1]$ . This will be our choice of  $\mathcal{G}$  throughout this section. In particular, by [AFP05, Rem. 3.5, Thm. 3.23], it is ensured that  $\mathcal{G}(z_n) \leq C$  for a sequence  $(z_n)_n$  implies that there exists  $z$  with  $\mathcal{G}(z) \leq C$  and a subsequence such that  $z_n \rightarrow z$  in  $L^1(\Omega)$ , and that  $\mathcal{G}$  is lower semicontinuous with respect to strong convergence in  $L^1(\Omega)$ . Moreover, to ensure that the energy of the system decreases with increasing damage, which is a typical feature of material damage, the function  $g$  is chosen as follows:

$$g \in C^0([0, 1]) \text{ monotonically increasing with } 0 < g_0 < g(0) < g(1) < g_1 < +\infty; \quad (6.10)$$

the fact that  $0 < g_0 < g(0)$  reflects partial damage, as this constraint ensures the coercivity of  $\mathcal{E}(t, \cdot, 0)$  given that  $W$  is chosen suitably. More precisely, for the elastic energy density  $W$  we assume

$$\text{Convexity: } W \in C^0(\mathbb{R}^{d \times d}, \mathbb{R}^+) \text{ is convex,} \quad (6.11a)$$

$$\text{Coercivity \& growth: } \exists C, C' > 0, p \in (1, \infty), \forall e \in \mathbb{R}^{d \times d} : C(|e|^p - 1) \leq W(e) \leq C'(|e|^p + 1). \quad (6.11b)$$

Observe that (6.11a) and (6.11b) imply the following

$$\text{Stress control: } \exists C > 0, p' = p/(p-1), \forall e \in \mathbb{R}^{d \times d} : |\partial_e^{\text{sup}} W(e)|^{p'} \leq C(W(e) + 1), \quad (6.11c)$$

where  $\partial_e W : \mathbb{R}^{d \times d} \rightrightarrows \mathbb{R}^{d \times d}$  is the convex subdifferential, and  $|\partial_e^{\text{sup}} W(e)| := \sup\{|\xi| : \xi \in \partial_e W(e)\}$ . Let us stress that we are not assuming any smoothness condition on  $W$ . As for the viscous dissipation  $\mathcal{V} : \mathbf{V} \rightarrow [0, +\infty)$  we stay general for the moment and only assume that

$$\mathbf{U} \subset \mathbf{V} \text{ continuously, and } \mathcal{V} \text{ complies with Hyp. 2.3.} \quad (6.12)$$

In addition, we are going to consider the case of inertia with the kinetic energy given by (6.6). We are going to use this example to emphasize how the growth assumptions on  $W$  affect the choice of  $\mathcal{V}$ , and which of the existence theorems is applicable in the different cases.

In view of (6.7), (6.9), and (6.11) we have

$$\mathbf{U} := W_{\Gamma_D}^{1,p}(\Omega; \mathbb{R}^d), \quad \mathbf{X} := \text{BV}(\Omega) \cap L^\infty(\Omega), \quad \mathbf{Z} := L^1(\Omega). \quad (6.13)$$

We now summarize the basic properties of  $\mathcal{E}$  and of the corresponding system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ .

**Proposition 6.1** (Properties of  $\mathcal{E}$  and  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ ). *Let the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$  be given by (6.7)–(6.13). Then, the following statements hold:*

- (1) *The energy functional  $\mathcal{E} : [0, T] \times \mathbf{V} \times \mathbf{Z} \rightarrow (-\infty, +\infty]$  has bounded sublevels in  $\mathbf{U} \times \mathbf{X}$ , i.e. also (2.19) from Hyp. 2.7 holds true.*
- (2)  *$\mathcal{E} : [0, T] \times \mathbf{V} \times \mathbf{Z} \rightarrow (-\infty, +\infty]$  is bounded from below and lower semicontinuous w.r.t. weak convergence in  $\mathbf{V}$  for  $\mathbf{u}$  and strong convergence in  $L^q(\Omega)$ , for any  $q \in [1, \infty)$ , for  $z$ , whence (2.6a) & (2.6b) in Hyp. 2.2. Furthermore, also condition (2.6c) on  $\partial_t \mathcal{E}$  is verified.*
- (3) *The mutual recovery sequence condition from Hyp. 2.5 holds true.*

*Proof. Ad (1):* The boundedness of the energy sublevels ensues on the one hand directly from the definition of  $\mathcal{G}$  and on the other hand from coercivity estimate (6.11b) together with Korn's inequality and the bounds on the given data, since

$$\begin{aligned} \mathcal{E}(t, \mathbf{u}, z) &\geq \int_{\Omega} g_0 C |\varepsilon(\mathbf{u} + \mathbf{u}_D)|^p dx - \sup_t \|\mathbf{f}(t)\|_{\mathbf{V}^*} \|\mathbf{u}\|_{\mathbf{V}} \\ &\geq g_0 C 2^{1-p} C_K \|\mathbf{u}\|_{\mathbf{U}}^p - a C \|\mathbf{u}_D(t)\|_{\mathbf{U}}^p - C_{\mathbf{V}\mathbf{U}} \sup_t \|\mathbf{f}(t)\|_{\mathbf{V}^*} \|\mathbf{u}\|_{\mathbf{U}} \\ &\geq g_0 C 2^{-p} C_K \|\mathbf{u}\|_{\mathbf{U}}^p - \tilde{C} = c \|\mathbf{u}\|_{\mathbf{U}}^p - \tilde{C}. \end{aligned} \quad (6.14)$$

*Ad (2):* The boundedness from below is a direct consequence from the above deduced estimate together with the fact that  $\mathcal{G}(z) \geq 0$  for all  $z \in \mathbf{Z}$ . The lower semicontinuity of  $\mathcal{E}(t, \cdot, \cdot)$  w.r.t. weak convergence in  $\mathbf{V}$  and strong convergence in  $L^q(\Omega)$  is ensured by [AFP05, Rem. 3.5, Thm. 3.23], assumption (6.11a), and lower semicontinuity results such as e.g. [FL07, Thm. 7.5]. Using that  $\partial_t \mathcal{E}(t, \mathbf{u}, z) = -\langle \mathbf{f}'(t), \mathbf{u} \rangle_{\mathbf{V}}$  and exploiting that  $\mathbf{f} \in C^1([0, T]; \mathbf{V}^*)$ , we immediately check (2.6c).

*Ad (3):* For the construction of the mutual recovery sequence we refer to [Tho13]. For given  $\hat{z} \in \mathbf{X}$  with  $\hat{z} \leq z$  a.e. in  $\Omega$  this construction is such that  $0 \leq \hat{z}_k \leq z_k$  a.e. in  $\Omega$ ,  $\hat{z}_k \rightarrow \hat{z}$  in  $L^1(\Omega)$ ,  $\limsup_{k \rightarrow \infty} (\mathcal{G}(\hat{z}_k) - \mathcal{G}(z_k)) \leq \mathcal{G}(\hat{z}) - \mathcal{G}(z)$ . Hence, also  $\mathcal{R}(\hat{z}_k - z_k) \rightarrow \mathcal{R}(\hat{z} - z)$ . In order to find that  $\limsup_{k \rightarrow \infty} \int_{\Omega} (g(\hat{z}_k) - g(z_k)) W(\varepsilon(\mathbf{u}_k + \mathbf{u}_D(t))) dx \leq \int_{\Omega} (g(\hat{z}) - g(z)) W(\varepsilon(\mathbf{u} + \mathbf{u}_D(t))) dx$  we observe that the integrand on the left- and on the right-hand side is non-positive. We now define the functional  $\tilde{\mathcal{E}} : \mathbf{Z} \times \mathbf{Z} \times \mathbf{V} \rightarrow [0, +\infty]$ ,  $\tilde{\mathcal{E}}(z, \hat{z}, \mathbf{u}) := \int_{\Omega} (g(z) - g(\hat{z})) W(\varepsilon(\mathbf{u} + \mathbf{u}_D)) + I_{[0, +\infty)}(z - \hat{z}) dx$  and observe that  $\tilde{\mathcal{E}}$  is lower semicontinuous with respect to strong convergence in  $L^q(\Omega)$  and weak convergence in  $\mathbf{V}$ . Hence, we obtain the desired estimate by lower semicontinuity.  $\square$

In order to establish the whole bunch of assumptions of the abstract existence results, it remains to discuss the conditions on  $\partial_u^- \mathcal{E}(t, \cdot, z)$ . For this, we note that  $u \mapsto \mathcal{E}(t, \mathbf{u}, z)$  is convex, hence  $\partial_u^- \mathcal{E}(t, \mathbf{u}, z)$  coincides with the convex subdifferential  $\partial_u \mathcal{E}(t, \mathbf{u}, z)$  and will thus be denoted by  $\partial_u \mathcal{E}(t, \mathbf{u}, z)$  hereafter. It is given by

$$\begin{aligned} \xi \in \partial_u \mathcal{E}(t, \mathbf{u}, z) \subset \mathbf{V}^* \text{ if and only if } \xi &= -\text{div}(g(z)\tilde{\xi}) - \mathbf{f}(t) \text{ with } \tilde{\xi} \in \partial_e W(\varepsilon(\mathbf{u})) \text{ for all } (t, \mathbf{u}, z) \in \text{dom}(\partial_u \mathcal{E}) \\ &\text{with } \text{dom}(\partial_u \mathcal{E}) = \{(t, \mathbf{u}, z) \in [0, T] \times \mathbf{V} \times \mathbf{Z} : \exists \tilde{\xi} \in \partial_e W(\varepsilon(\mathbf{u})) \text{ s.t. } -\text{div}(g(z)\tilde{\xi}) \in \mathbf{V}^*\}. \end{aligned} \quad (6.15)$$

To derive (6.15), we observe that  $\mathcal{E}(t, \mathbf{u}, z) = \mathcal{W}(\mathbf{u}, z) - \langle \mathbf{f}(t), \mathbf{u} \rangle_{\mathbf{V}}$  with the stored elastic energy term  $\mathcal{W}(\mathbf{u}, z) := \int_{\Omega} g(z) W(\varepsilon(\mathbf{u})) dx$ . Since the second contribution to  $\mathcal{E}$  is smooth as a function of  $\mathbf{u}$ , the sum rule applies, yielding that  $\partial_u \mathcal{E}(t, \mathbf{u}, z) = \partial_u \mathcal{W}(\mathbf{u}, z) - \mathbf{f}(t)$ . In turn, we compute the convex subdifferential  $\partial_u \mathcal{W}$  by applying the chain rule from [IT79, Chap. 4, Thm. 2, p. 201], and thus conclude that  $\zeta \in \partial_u \mathcal{W}(\mathbf{u}, z)$  if and only if  $\zeta = -\text{div}(g(z)\tilde{\xi})$ , with  $\tilde{\xi} \in \partial_e W(\varepsilon(\mathbf{u}))$  such that  $-\text{div}(g(z)\tilde{\xi}) \in \mathbf{V}^*$ . Hence, (6.15) ensues. Clearly, by the very definition of convex subdifferential, the Fréchet subdifferentiability conditions are satisfied, cf. Hyp. 4.2 for in the gradient system case, and Hyp. 5.2 for the case with inertia. We now have to verify the closedness

of the subdifferential  $\partial_u \mathcal{E}(t, \cdot, z)$ . Moreover, in the case with inertia also the subgradient estimate proposed in Hyp. 5.3 has to be checked.

To do so, we are going to distinguish between the case that  $\mathcal{V}$  has general superlinear growth and the case that  $\mathcal{V}$  is quadratic. The latter case e.g. covers Kelvin-Voigt rheology, where  $\mathcal{V}$  is given by (6.2), and hence  $\mathbf{V} = H_{\Gamma_D}^1(\Omega; \mathbb{R}^d)$  so that, for  $\mathbf{U} \subset \mathbf{V}$  continuously,  $p \geq 2$  is necessary. One may also consider a weak damping of the form  $\mathcal{V}(\mathbf{u}') := \frac{1}{2} \int_{\Omega} c |\mathbf{u}'|^2 dx$ , i.e. here  $\mathbf{V} = L^2(\Omega; \mathbb{R}^d)$  and  $\mathbf{U} \Subset \mathbf{V}$  for any  $p > 2d/(d+2)$ . In the case that  $\mathcal{V}$  is of general superlinear growth we have in mind that e.g.  $\mathcal{V}(\mathbf{u}') := \|\varepsilon(\mathbf{u}')\|_{L^r(\Omega; \mathbb{R}^{d \times d})}^q$  with  $1 < r \leq p$  and  $q > 1$ , such that  $\mathbf{V} = W^{1,r}(\Omega; \mathbb{R}^d)$  and  $\mathbf{U} \subset \mathbf{V}$  continuously. But also a weak damping can be handled of the form  $\mathcal{V}(\mathbf{u}') := \|\mathbf{u}'\|_{L^r(\Omega; \mathbb{R}^d)}^q$  with  $p > rd/(d+r)$  and  $r, q > 1$ , such that  $\mathbf{V} = L^r(\Omega; \mathbb{R}^d)$  and  $\mathbf{U} \Subset \mathbf{V}$ . Moreover we will treat gradient systems and damped inertial systems separately.

**The case of gradient systems and  $\mathcal{V}$  quadratic.** In this setting (which already falls outside the scope of the results in [Rou09]), as outlined in (4.1), the viscous dissipation potential  $\mathcal{V}(v) := \frac{1}{2}a(v, v)$  is defined via a continuous and coercive bilinear form  $a : \mathbf{V} \times \mathbf{V} \rightarrow \mathbb{R}$ . Therefore, with the induced inner product  $\mathbf{V}$  is a Hilbert space, and we thus have to distinguish the cases  $\mathbf{V} = L^2(\Omega; \mathbb{R}^d)$  and  $\mathbf{V} = H_{\Gamma_D}^1(\Omega; \mathbb{R}^d)$ . Now, the assumptions of Theorem 4.9 have to be checked. In view of Prop. 6.1, it remains to establish the closedness of  $\partial_u \mathcal{E}$ . According to Section 4.1, this can be done by verifying the recovery sequence Hypothesis 4.6 if  $\mathbf{U} \Subset \mathbf{V}$  (i.e. if  $\mathbf{V} = L^2(\Omega; \mathbb{R}^d)$ ), whereas, if only  $\mathbf{U} \subset \mathbf{V}$  continuously (i.e., for  $\mathbf{V} = H_{\Gamma_D}^1(\Omega; \mathbb{R}^d)$ ), the closedness is to be verified via the closedness-argument à la Minty from Hyp. 4.8.

**Proposition 6.2** (Properties of  $\partial_u \mathcal{E}(t, \cdot, z)$ ). *Let the assumptions of Prop. 6.1 hold true and assume that  $\mathcal{V} : \mathbf{V} \rightarrow [0, +\infty]$  is quadratic.*

- (1) *Let  $\mathbf{U} \Subset \mathbf{V}$ . Then  $\mathcal{E} : [0, T] \times \mathbf{V} \times \mathbf{Z} \rightarrow (-\infty, +\infty]$  satisfies the recovery sequence Hypothesis 4.6.*
- (2) *Let  $\mathbf{U} \subset \mathbf{V}$  continuously. Then  $\mathcal{E} : [0, T] \times \mathbf{V} \times \mathbf{Z} \rightarrow (-\infty, +\infty]$  satisfies the closedness à la Minty from Hypothesis 4.8.*

*Proof.* **Ad (1):** For any given  $\tilde{\mathbf{u}} \in D_u$ , the constant sequence  $\tilde{\mathbf{u}}_k = \tilde{\mathbf{u}}$  serves as a recovery sequence in Hyp. 4.6.

**Ad (2):** Consider a sequence  $(\mathbf{t}_k, \mathbf{u}_k, z_k)_k \subset [0, T] \times \mathbf{V} \times \mathbf{Z}$  complying with the assumptions in (4.13a), i.e.  $\mathbf{t}_k \rightarrow \mathbf{t}$  pointwise a.e. in  $(0, T)$ ,  $\mathbf{u}_k \rightharpoonup \mathbf{u}$  in  $L^\infty(0, T; \mathbf{U}) \cap H^1(0, T; \mathbf{V})$ ,  $z_k \overset{*}{\rightharpoonup} z$  in  $L^\infty(0, T; L^q(\Omega))$  and  $z_k(t) \rightarrow z(t)$  in  $L^q(\Omega)$  for any  $1 \leq q < \infty$  and all  $t \in [0, T]$ ,  $\xi_k \rightharpoonup \xi$  in  $L^2(0, T; \mathbf{V}^*)$  with  $\xi_k \in \partial_u \mathcal{E}(\mathbf{t}_k(t), \mathbf{u}_k(t), z_k(t))$  for a.e.  $t \in (0, T)$ , and  $\limsup_k \int_0^T \langle \xi_k, \mathbf{u}_k \rangle_{\mathbf{V}} dt \leq \int_0^T \langle \xi, \mathbf{u} \rangle_{\mathbf{V}} dt$ : We have to verify that also  $\xi(t) \in \partial_u \mathcal{E}(\mathbf{t}(t), \mathbf{u}(t), z(t))$  for a.e.  $t \in (0, T)$ . First of all, from the above lim sup-condition, taking into account the representation formula (6.15) for  $\xi_k$  and the fact that  $\mathbf{f}(\mathbf{t}_k) \rightarrow \mathbf{f}(\mathbf{t})$  in  $L^p(0, T; \mathbf{V}^*)$  for every  $1 \leq p < \infty$ , we deduce that

$$\limsup_{k \rightarrow \infty} \int_0^T \int_{\Omega} g(z_k) \tilde{\xi}_k : \varepsilon(\mathbf{u}_k) dx dt \leq \int_0^T \langle \tilde{\zeta}, \mathbf{u} \rangle_{\mathbf{V}} dx dt \quad (6.16)$$

with  $\tilde{\zeta} = \xi + \mathbf{f}(\mathbf{t}(t)) \in L^2(0, T; \mathbf{V}^*)$ . It can be checked that the integral functionals  $\mathcal{W}_k(\mathbf{u}) := \int_0^T \mathcal{W}(\mathbf{u}(t), z_k(t)) dt$  Mosco-converge to  $\mathcal{W}(\mathbf{u}) := \int_0^T \mathcal{W}(\mathbf{u}(t), z(t)) dt$  w.r.t. the topology of  $L^2(0, T; \mathbf{V})$ . Hence, their convex subdifferentials converge in the sense of graphs to the subdifferential  $\partial \mathcal{W}$  (cf. [Att84]). Therefore, from the lim sup-condition (6.16) we deduce that  $\tilde{\zeta} \in \partial \mathcal{W}(\mathbf{u})$ , i.e.  $\tilde{\zeta}(t) \in \partial_u \mathcal{W}(\mathbf{u}(t), z(t))$  for a.a.  $t \in (0, T)$ . All in all, we conclude that  $\xi(t) \in \partial_u \mathcal{E}(\mathbf{t}(t), \mathbf{u}(t), z(t))$  for a.e.  $t \in (0, T)$ .  $\square$

Now, Theorem 4.9 guarantees the existence of a semistable energetic solution.

**Theorem 6.3.** *Let the assumptions of Prop. 6.2 hold true. Then the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  admits a semistable energetic solution.*

**The gradient system case and  $\mathcal{V}$  of general superlinear growth.** In this frame, the closedness of  $\partial_u \mathcal{E}(t, \cdot, z)$  is to be verified via Hypothesis 4.12. This condition requires the identification of the limit (as an element of the limit subdifferential) via *weak convergence both* of the sequence  $(\mathbf{u}_k)_k$  and the elements  $(\xi_k)_k$  of their corresponding subdifferentials *without any additional identification qualification* (such as the existence of a recovery sequence for the energy functional, cf. Hyp. 4.6, or the closedness à la Minty, cf. Hyp. 4.8). In view of the form (6.8) of the energy functional, such identification can be carried out if  $W$  is of quadratic nature, cf. also Example 4.16. In other words, in case of a viscous dissipation potential of general superlinear growth,

the energy functional has to be of quadratic growth in the respective variable, i.e., here, we may consider

$$W(e) := \frac{1}{2}e:\mathbb{C}:e \quad \text{with } \mathbb{C} \in \mathbb{R}^{d \times d \times d \times d} \text{ symmetric and positively definite.} \quad (6.17)$$

Hence,  $\mathbf{U} = H_{\Gamma_D}^1(\Omega; \mathbb{R}^d)$  and the condition  $\mathbf{U} \subset \mathbf{V}$  restricts  $\mathbf{V}$  to be either a Sobolev space  $W_{\Gamma_D}^{1,q}(\Omega; \mathbb{R}^d)$  with  $1 < q \leq 2$  or a Lebesgue space  $L^q(\Omega; \mathbb{R}^d)$  with  $1 < q \leq 2d/(d-2)$  for  $d > 2$  and  $1 < q < \infty$  for  $d = 2$ . By the representation formula (6.15), we have with the quadratic ansatz (6.17) that, if  $\mathbf{u}_k \rightarrow \mathbf{u}$  in  $\mathbf{V}$ ,  $z_k \rightarrow z$  in  $\mathbf{Z}$ ,  $\xi_k \rightarrow \xi$  in  $\mathbf{V}^*$  with  $\xi_k \in \partial_u \mathcal{E}(t, \mathbf{u}_k, z_k) = \{g(z_k)\mathbb{C} : \varepsilon(\mathbf{u}_k)\}$  for all  $k \in \mathbb{N}$ , the closedness of the subdifferential, i.e., that also  $\xi \in \partial_u \mathcal{E}(t, \mathbf{u}, z) = \{g(z)\mathbb{C} : \varepsilon(\mathbf{u})\}$ . Therefore we now summarize

**Proposition 6.4** (Properties of  $\partial_u \mathcal{E}(t, \cdot, z)$  and  $\partial_t \mathcal{E}(t, \mathbf{u}, z)$ ). *Let the assumptions of Prop. 6.1 hold true and let  $\mathcal{E}(t, \cdot, z)$  be quadratic as in (6.17) as well as  $\mathbf{U} = H_0^1(\Omega; \mathbb{R}^d) \subset \mathbf{V}$ . Then, the system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  satisfies the closedness Hypothesis 4.12.*

Now, we may invoke Theorem 4.13, which ensures the existence of a weak semistable energetic solution.

**Theorem 6.5.** *Let the assumptions of Prop. 6.4 be satisfied. Then the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  admits a weak semistable energetic solution.*

**The case with inertia and  $\mathcal{V}$  of general superlinear growth.** As above, the closedness of  $\partial_u \mathcal{E}(t, \cdot, z)$  has to be verified via Hypothesis 4.12 and hence, also here the energy functional has to be quadratic with  $W$  of the form (6.17) and  $\mathbf{U} = H_{\Gamma_D}^1(\Omega; \mathbb{R}^d)$ . In addition, in the case with inertia also the subgradient estimate due to Hyp. 5.3 has to be shown. More precisely, for  $\xi \in \partial_u \mathcal{E}(t, \mathbf{u}, z)$  it has to be established that  $|\langle \xi, v \rangle_{\mathbf{V}}|^\sigma \leq C(\mathcal{E}(t, \mathbf{u}, z) + \|\mathbf{u}\|_{\mathbf{V}} + 1)\|v\|_{\mathbf{V}}$  for all  $v \in \mathbf{V}$ . However, taking a closer look at the representation formula (6.15) we see that the stress control (6.11c) can only imply the subgradient estimate in terms of the energy if the divergence can be swapped to the test function  $v \in \mathbf{V}$  by integration by parts. But this requires that  $\varepsilon(v) \in L^p(\Omega; \mathbb{R}^{d \times d})$  with  $p \geq 2$  for any  $v \in \mathbf{V}$ , hence  $\mathbf{V} \subset W_{\Gamma_D}^{1,p}(\Omega; \mathbb{R}^d)$  with  $p \geq 2$ . On other hand,  $\mathbf{U} \subset \mathbf{V}$  enforces  $p \leq 2$ . All in all, this leads to  $\mathbf{V} = \mathbf{U} = H_{\Gamma_D}^1(\Omega; \mathbb{R}^d)$ .

**Proposition 6.6** (Properties of  $\partial_u \mathcal{E}(t, \cdot, z)$  and  $\partial_t \mathcal{E}(t, \mathbf{u}, z)$ ). *Let the assumptions of Prop. 6.4 be satisfied and let  $\mathbf{V} = \mathbf{U} = H_0^1(\Omega; \mathbb{R}^d)$ . Then, the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$  satisfies the closedness Hypothesis 4.12 as well as condition (2.6c) and  $\partial_u \mathcal{E}(t, \cdot, z)$  complies with the subgradient estimate (5.8) from Hyp. 5.3.*

Due to these findings the abstract existence Theorem 5.4 yields the existence of a weak semistable energetic solution.

**Theorem 6.7.** *Let the assumptions of Prop. 6.6 hold true. Then the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$  admits a weak semistable energetic solution.*

**The case with inertia,  $\mathcal{V}$  quadratic, and  $\mathbf{V} \Subset \mathbf{W}$ .** Here, the topological setup  $\mathbf{U} \subset \mathbf{V} \Subset \mathbf{W}$  together with (6.13) determines that  $\mathbf{V} := W_{\Gamma_D}^{1,r}(\Omega; \mathbb{R}^d)$  with  $r \in [r^*, p]$  for  $r^* = 2d/(d+2)$  if  $d > 2$ ,  $r^* > 1$  if  $d = 2$ . However, as already seen above, the subgradient estimate due to Hyp. 5.3 enforces that  $\mathbf{V} = \mathbf{U} = H_{\Gamma_D}^1(\Omega; \mathbb{R}^d)$ . Moreover, for this setting, the closedness of  $\partial_u \mathcal{E}(t, \cdot, z)$  is again to be verified via the closedness-argument à la Minty, as already done in Prop. 6.2(2).

**Proposition 6.8** (Properties of  $\partial_u \mathcal{E}(t, \cdot, z)$  and  $\partial_t \mathcal{E}(t, \mathbf{u}, z)$ ). *Let the assumptions of Prop. 6.2 be satisfied and let  $\mathcal{V}$  be quadratic and  $\mathbf{V} = \mathbf{U} \Subset \mathbf{W}$ . Then, the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$  satisfies the closedness Hypothesis 4.8, condition (2.6c) and  $\partial_u \mathcal{E}$  complies with the subgradient estimate (5.8) from Hyp. 5.3.*

This enables us to deduce the existence of semistable energetic solutions via Theorem 5.6.

**Theorem 6.9.** *Let the assumptions of Prop. 6.8 be satisfied. Then the damped inertial system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$  admits a semistable energetic solution.*

**6.4. Rate-dependent damage and rate-independent plasticity with hardening at small strains.** We now focus on a coupled elasto-plastic damage model (at small strains) that was proposed in [AMV14] and first mathematically analyzed in [Cri16, CL16]. While [Cri16] addresses the existence of (*quasistatic evolutions*) for the fully rate-independent evolution of plasticity and damage, [CL16] studies the system regularized by viscosity in the damage variable, and develops the asymptotic analysis for vanishing viscosity. Let us also mention [RSV13, RV16] for the analysis of other systems coupling perfect plasticity and rate-dependent damage, with healing, in the context of the geophysical modeling of lithospheric faults with recurring earthquakes.

In what follows, we aim to revisit the mixed rate-dependent (for the damage variable) / rate-independent (for the plasticity tensor) system studied in [CL16]. There, the authors have proved the existence of a weak notion of solution featuring a formulation of the damage flow rule in terms of a Kuhn-Tucker inequality, in place of a standard subdifferential inclusion, coupled with an energy-dissipation balance. In fact, such a formulation is akin to the weak solvability concept proposed and analyzed in [HK11, HK13] for (rate-dependent) damage coupled with phase separation.

Here we are going to address a variant of this damage/plasticity system, featuring hardening and a gradient regularization for the plastic tensor, and where unidirectionality of damage is neglected. To this regularized system it is possible to apply our abstract approach, leading to the existence of *semistable energetic solutions*, which are in fact stronger than the solutions obtained in [CL16]. In particular, we will solve the damage flow rule pointwise a.e. in the space-time domain, cf. (6.33) below.

**Energy.** We consider damage in a body located in a sufficiently smooth (cf. (6.27) below) bounded domain  $\Omega \subset \mathbb{R}^d$  with  $d \in \{2, 3\}$ . The “viscous” and “rate-independent” variables are, respectively

$$u := \alpha \in [0, 1] \text{ damage parameter; } \quad z := p \in \mathbb{R}_{\text{dev}}^{d \times d} := \{\pi \in \mathbb{R}_{\text{sym}}^{d \times d} : \text{tr}(\pi) = 0\} \text{ plastic tensor,}$$

whereas we will continue to denote by  $\mathbf{u}$  the displacement variable, subject to a time-dependent loading  $\mathbf{u}_D$  on the Dirichlet boundary  $\Gamma_D$ . The overall energy is given by the sum of three contributions

$$\mathcal{E}(t, \alpha, p) := \mathcal{E}_1(\alpha) + \mathcal{E}_2(p) + \mathcal{E}_3(t, \alpha, p). \quad (6.18)$$

We take

$$\mathcal{E}_1 : H^s(\Omega) \rightarrow (-\infty, +\infty], \quad \mathcal{E}_1(\alpha) := a_s(\alpha, \alpha) + \int_{\Omega} I_{[0,1]}(\alpha) + \gamma(\alpha) dx, \quad (6.19)$$

where, along the lines of [KRZ13], and as also done in [CL16], we choose for the gradient regularization of the damage variable the bilinear form

$$a_s(\alpha_1, \alpha_2) = \int_{\Omega} \int_{\Omega} \frac{(\nabla \alpha_1(x) - \nabla \alpha_1(y)) \cdot (\nabla \alpha_2(x) - \nabla \alpha_2(y))}{|x - y|^{d+2(s-1)}} dx dy \quad \text{with } s \in \left(\frac{d}{2}, 2\right) \quad (6.20)$$

associated with the  $s$ -Laplacian operator  $A_s : H^s(\Omega) \rightarrow H^s(\Omega)^*$  imposing homogeneous Neumann boundary conditions (with  $H^s(\Omega)$  we denote the space  $W^{s,2}(\Omega)$ ). In what follows, we will exploit in a crucial way that, since  $s > \frac{d}{2}$ , there holds

$$H^s(\Omega) \Subset C^{0,\delta}(\bar{\Omega}) \quad \text{for some } \delta \in (0, 1]. \quad (6.21)$$

The other contributions to the energy  $\mathcal{E}_1$  are the indicator function of the interval  $[0, 1]$ , enforcing the constraint  $z \in [0, 1]$ , and a smooth, but possibly nonconvex, function  $\gamma \in C^1([0, 1])$ . Unlike in [CL16], where the case of perfect plasticity was considered, we encompass hardening and gradient plasticity in our model through

$$\mathcal{E}_2 : H^1(\Omega; \mathbb{R}_{\text{dev}}^{d \times d}) \rightarrow (0, +\infty), \quad \mathcal{E}_2(p) := \int_{\Omega} \frac{1}{2} p : \mathbb{H} : p + |\nabla p|^2 dx \quad (6.22a)$$

with  $\mathbb{H}$  a symmetric, positive definite operator (of Prager/Ziegler type) on  $\mathbb{R}_{\text{dev}}^{d \times d}$  such that

$$\exists C_{\mathbb{H}} > 0 \quad \forall p \in \mathbb{R}_{\text{dev}}^{d \times d} : \quad p : \mathbb{H} : p \geq C_{\mathbb{H}} |p|^2. \quad (6.22b)$$

Finally, we define  $\mathcal{E}_3 : [0, T] \times C^0(\bar{\Omega}; [0, 1]) \times L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$  by minimizing out the displacement from the elastic energy, namely

$$\begin{aligned} \mathcal{E}_3(t, \alpha, p) &:= \min_{\mathbf{u} \in H_{\Gamma_D}^1(\Omega; \mathbb{R}^d)} \mathcal{J}(t, \alpha, \mathbf{u}, p) \quad \text{with} \\ \mathcal{J}(t, \alpha, \mathbf{u}, p) &:= \int_{\Omega} \frac{1}{2} g(\alpha) (\varepsilon(\mathbf{u} + \mathbf{u}_D) - p) : \mathbb{C} : (\varepsilon(\mathbf{u} + \mathbf{u}_D) - p) dx - \langle \mathbf{f}(t), \mathbf{u} \rangle. \end{aligned} \quad (6.23)$$

Here, following [KRZ13, Sec. 2.4], we assume that

$$\begin{aligned} \mathbb{C} &\in C^{0,\delta}(\overline{\Omega}; \text{Lin}(\mathbb{R}_{\text{sym}}^{d \times d}, \mathbb{R}_{\text{sym}}^{d \times d})) \quad \text{for } \delta \in (0, 1] \text{ as in (6.21),} \\ \exists c_0 > 0 \quad \text{for all } \xi \in \mathbb{R}_{\text{sym}}^{d \times d} \text{ and all } x \in \Omega : \quad \mathbb{C}(x)\xi : \xi \geq c_0|\xi|^2, \end{aligned} \quad (6.24)$$

and that the constitutive function  $g$  fulfills

$$g \in C^1([0, 1]), \quad \text{and } \exists c_1 > 0 \quad \forall z \in [0, 1] : \quad g(z) \geq c_1. \quad (6.25)$$

Along the footsteps of [KRZ13], we assume for the Dirichlet datum  $\mathbf{u}_D$  and the external load  $\mathbf{f}$  that

$$\mathbf{u}_D \in C^{1,1}([0, T]; W^{1,4}(\Omega; \mathbb{R}^d)), \quad \mathbf{f} \in C^{1,1}([0, T]; W^{-1,4}(\Omega; \mathbb{R}^d)). \quad (6.26)$$

Let us mention in advance that the choice of the space  $W^{1,4}(\Omega; \mathbb{R}^d)$  in (6.26) will have a crucial role in the derivation of *higher integrability* estimates for the (unique) minimizer of the energy  $\mathcal{J}(t, \alpha, \cdot, p)$ , cf. Lemma 6.10 below, along with the condition that

$$\Omega \text{ has } C^1\text{-boundary } \partial\Omega, \text{ and Dirichlet boundary } \Gamma_D = \partial\Omega. \quad (6.27)$$

**Dissipation.** For the damage variable, we consider the quadratic dissipation potential

$$\mathcal{V}(\alpha') := \int_{\Omega} \frac{1}{2} |\alpha'|^2 dx \quad \forall \alpha' \in \mathbf{V} := L^2(\Omega). \quad (6.28)$$

Clearly,  $\mathcal{V}$  complies with Hypothesis 2.3. Observe that  $\mathcal{V}$  does not encompass the unidirectionality constraint  $\alpha' \leq 0$ , since, as mentioned in the introduction, in the analysis developed in this paper we cannot allow for  $\mathcal{V}$  taking the value  $+\infty$ . For the plastic tensor, along the footsteps of [AMV14], we include in the 1-homogeneous dissipation potential a dependence on the damage variable, namely we define

$$\begin{aligned} \mathcal{R} : C^0(\overline{\Omega}; [0, 1]) \times \mathbf{Z} &\rightarrow [0, +\infty), \quad \text{with } \mathbf{Z} := L^1(\Omega; \mathbb{R}_{\text{dev}}^{d \times d}), \quad \text{by } \mathcal{R}(\alpha, \pi') := \int_{\Omega} H(\alpha, \pi') dx \quad \text{with} \\ H : [0, 1] \times \mathbb{R}_{\text{dev}}^{d \times d} &\rightarrow [0, +\infty) \text{ continuous, } H(\alpha, \cdot) \text{ convex and 1-positively homogeneous } \forall \alpha \in [0, 1], \\ \exists r, R > 0 \quad \forall (\alpha, \pi') &\in [0, 1] \times \mathbb{R}_{\text{dev}}^{d \times d} : \quad r|\pi'| \leq H(\alpha, \pi') \leq R|\pi'|. \end{aligned} \quad (6.29a)$$

It follows from (6.29a) that for every  $\alpha \in [0, 1]$  the dissipation density  $H(\alpha, \cdot)$  is the support function of a closed convex  $K(\alpha) \subset \mathbb{R}_{\text{dev}}^{d \times d}$ , i.e.

$$H(\alpha, \pi') = \sup_{\sigma \in K(\alpha)} \sigma : \pi'.$$

In the lines of [Cri16, Rmk. 2.1], we further specialize to the case in which

$$K(\alpha) = h(\alpha)K \quad \text{with } \begin{cases} h : [0, 1] \rightarrow [0, +\infty) \text{ continuous, and } \exists c_2 > 0 \quad \forall \alpha \in [0, 1] : h(\alpha) \geq c_2 > 0, \\ K \subset \mathbb{R}_{\text{dev}}^{d \times d} \text{ closed, convex, with } B_r(0) \subset K \subset B_R(0) \text{ for some } 0 < r < R. \end{cases} \quad (6.29b)$$

Hence, we have the following explicit formula for  $H$

$$H(\alpha, \pi') = h(\alpha) \sup_{\sigma \in K} \sigma : \pi' \doteq h(\alpha) |\pi'|_K, \quad (6.30)$$

where we have used the norm  $|\cdot|_K$  induced by  $K$ , equivalent to the Euclidean norm thanks to (6.29b). It is immediate to verify that for every  $\alpha \in C^0(\overline{\Omega}; [0, 1])$  the dissipation potential  $\mathcal{R}(\alpha, \cdot)$  complies with Hyp. 2.3, with a constant uniform w.r.t.  $\alpha$ .

The special form (6.30) of  $H$  naturally leads to the following definition for the total variation functional induced by  $\mathcal{R}$ : we set on an interval  $[s, t] \subset [0, T]$

$$\text{Var}_{\mathcal{R}}(\alpha, p; [s, t]) := \iint_{[s, t] \times \overline{\Omega}} h(\alpha) |p'|_K(dx dr) \quad \text{for every } \alpha \in C^0([0, T]; C^0(\overline{\Omega}; [0, 1])) \text{ and } p \in \text{BV}([0, T]; \mathbf{Z}). \quad (6.31)$$

where the Radon measure  $p' \in \text{M}([0, T] \times \overline{\Omega}; \mathbb{R}_{\text{dev}}^{d \times d})$  is the distributional derivative of  $p$ , and  $|p'|_K$  its total variation with respect to the norm  $|\cdot|_K$ .

**The notion of semistable energetic solution and the time-discrete scheme.** We now revisit the notion of *semistable energetic* solution to the system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  given by (6.18), (6.28), and (6.29), adapting Def. 3.1 in the gradient system case, to the present situation with a *state-dependent* dissipation potential  $\mathcal{R}$ . Hence, the semistability condition reads

$$\mathcal{E}(t, \alpha(t), p(t)) \leq \mathcal{E}(t, \alpha(t), \tilde{p}) + \mathcal{R}(\alpha(t), \tilde{p} - p(t)) \quad \forall \tilde{p} \in L^1(\Omega; \mathbb{R}^{d \times d}) \quad \text{for all } t \in [0, T], \quad (6.32)$$

whereas the energy-dissipation inequality has the same form as (3.4) but features the  $\alpha$ -dependent total variation  $\text{Var}_{\mathcal{R}}(\alpha, p; [0, t])$ . Finally, let us highlight that the subdifferential inclusion for  $\alpha$  reads

$$\begin{aligned} \alpha'(t) + A_s \alpha(t) + \partial I_{[0,1]}(\alpha(t)) + \gamma'(\alpha(t)) \\ + \frac{1}{2} g'(\alpha)(\varepsilon(\mathbf{u}_{\min}(t) + \mathbf{u}_D(t)) - p(t)) : \mathbb{C} : (\varepsilon(\mathbf{u}_{\min}(t) + \mathbf{u}_D(t)) - p(t)) \ni 0 \quad \text{in } L^2(\Omega) \text{ for a.a. } t \in (0, T), \end{aligned} \quad (6.33)$$

where we have used the explicit formula (6.47) below for the Fréchet subdifferential of  $\alpha \mapsto \mathcal{E}(t, \alpha, p)$  (w.r.t. the  $L^2(\Omega)$ -topology), with  $\mathbf{u}_{\min}(t) \in H_{\Gamma_D}^1(\Omega; \mathbb{R}^d)$  the (unique) minimizer of  $\mathcal{J}(t, \alpha(t), \cdot, p(t))$ .

In order to prove the existence of semistable energetic solutions to  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ , we pass to the limit in the time-discretization scheme from Problem 4.1, where the first minimum problem (4.3a) now reads

$$p_\tau^n \in \underset{p \in \mathbf{Z}}{\text{Argmin}} \left( \tau \mathcal{R} \left( \alpha_\tau^{n-1}, \frac{p - p_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, \alpha_\tau^{n-1}, p) \right), \quad (6.34)$$

whereas (4.3b) stays the same. Hence, for the limit passage in the semistability condition it will be necessary to verify the following mutual recovery sequence condition, tailored to the fact that the map  $\alpha \mapsto \mathcal{G}(\alpha, p)$  has sublevels bounded in  $H^s(\Omega)$ , cf. Proposition 6.13 below, and that we will be able to prove for every  $t \in [0, T]$  that  $(\bar{\alpha}_\tau(t) - \underline{\alpha}_\tau(t)) \rightarrow 0$  in  $H^s(\Omega)$ , cf. (4.15c) :

Let  $(t_n, \bar{\alpha}_n, \alpha_n, p_n)_n \subset [0, T] \times H^s(\Omega) \times H^s(\Omega) \times H^1(\Omega; \mathbb{R}^{d \times d})$  fulfill for every  $n \in \mathbb{N}$  the semistability condition

$$\mathcal{E}(t_n, \alpha_n, p_n) \leq \mathcal{E}(t_n, \alpha_n, \tilde{p}) + \mathcal{R}(\bar{\alpha}_n, \tilde{p} - p_n) \quad \forall \tilde{p} \in L^1(\Omega; \mathbb{R}^{d \times d}),$$

and suppose that  $t_n \rightarrow t$ ,  $(\alpha_n, p_n) \rightharpoonup (\alpha, p)$  in  $H^s(\Omega) \times H^1(\Omega; \mathbb{R}^{d \times d})$  with  $\sup_{n \in \mathbb{N}} \mathcal{G}(\alpha_n, p_n) \leq C$ , and that  $(\alpha_n - \bar{\alpha}_n) \rightarrow 0$  in  $H^s(\Omega)$ . Then for every  $\tilde{p} \in L^1(\Omega; \mathbb{R}^{d \times d})$  there exists  $\tilde{p}_n \rightarrow \tilde{p}$  in  $L^1(\Omega; \mathbb{R}^{d \times d})$  such that

$$\lim_{n \rightarrow \infty} (\mathcal{R}(\bar{\alpha}_n, \tilde{p}_n - p_n) + \mathcal{E}(t_n, \alpha_n, \tilde{p}_n) - \mathcal{E}(t_n, \alpha_n, p_n)) \leq \mathcal{R}(\alpha, \tilde{p} - p) + \mathcal{E}(t, \alpha, \tilde{p}) - \mathcal{E}(t, \alpha, p). \quad (6.35)$$

**Properties of the reduced energy  $\mathcal{E}_3$ .** Prior to verifying the hypotheses of Thm. 4.9, we start with the following preliminary results, providing basic properties of the reduced energy  $\mathcal{E}_3$  from (6.23). In Lemmas 6.10–6.12 (whose proofs are indeed drawn from [KRZ13]), we will distinguish between the properties valid for  $p \in L^2(\Omega; \mathbb{R}^{d \times d})$ , and those requiring  $p \in L^4(\Omega; \mathbb{R}^{d \times d})$ .

**Lemma 6.10.** *Assume (6.24)–(6.27). Then, for every  $(t, \alpha, p) \in [0, T] \times C^0(\bar{\Omega}; [0, 1]) \times L^2(\Omega; \mathbb{R}^{d \times d})$  there exists a unique minimizer  $\mathbf{u}_{\min} = \mathbf{u}_{\min}(t, \alpha, p) \in H_{\Gamma_D}^1(\Omega; \mathbb{R}^d)$  and*

$$\begin{aligned} \forall \epsilon \in (0, c_0 c_1) \quad \exists d_1^\epsilon > 0 \quad \forall (t, \alpha, p) \in [0, T] \times C^0(\bar{\Omega}; [0, 1]) \times L^2(\Omega; \mathbb{R}^{d \times d}) : \\ \mathcal{E}_3(t, \alpha, p) \geq \frac{1}{2} (c_0 c_1 - \epsilon) \|\mathbf{u}_{\min}\|_{H^1(\Omega; \mathbb{R}^d)}^2 + \frac{1}{2} \left( c_0 c_1 - \frac{1}{\epsilon} \right) \|p\|_{L^2(\Omega; \mathbb{R}^{d \times d})}^2 \\ - d_1^\epsilon (\|\mathbf{u}_D\|_{L^\infty(0, T; W^{1,4}(\Omega; \mathbb{R}^d))}^2 + \|\mathbf{f}\|_{L^\infty(0, T; W^{-1,4}(\Omega; \mathbb{R}^d))}^2) \end{aligned} \quad (6.36)$$

Moreover, if  $p \in L^4(\Omega; \mathbb{R}^{d \times d})$ , then  $\mathbf{u}_{\min} \in W^{1,4}(\Omega; \mathbb{R}^d)$  and

for every  $M > 0$  there exist  $d_2(M), d_3(M) > 0$  such that

$$\forall (t, \alpha, p) \in [0, T] \times C^0(\bar{\Omega}; [0, 1]) \times L^4(\Omega; \mathbb{R}^{d \times d}), \quad \|\alpha\|_{C^0(\bar{\Omega})} + \|p\|_{L^4(\Omega; \mathbb{R}^{d \times d})} \leq M : \quad (6.37)$$

$$\|\mathbf{u}_{\min}(t, \alpha, p)\|_{W^{1,4}(\Omega; \mathbb{R}^d)} \leq d_2(M) (\|\mathbf{u}_D\|_{L^\infty(0, T; W^{1,4}(\Omega; \mathbb{R}^d))} + \|\mathbf{f}\|_{L^\infty(0, T; W^{-1,4}(\Omega; \mathbb{R}^d))});$$

$$\forall (t_1, \alpha_1, p), (t_2, \alpha_2, p) \in [0, T] \times C^0(\bar{\Omega}; [0, 1]) \times L^4(\Omega; \mathbb{R}^{d \times d}), \quad \|\alpha_1\|_{C^0(\bar{\Omega})} + \|\alpha_2\|_{C^0(\bar{\Omega})} + \|p\|_{L^4(\Omega; \mathbb{R}^{d \times d})} \leq M :$$

$$\|\mathbf{u}_{\min}(t_1, \alpha_1, p) - \mathbf{u}_{\min}(t_2, \alpha_2, p)\|_{W^{1,4}(\Omega; \mathbb{R}^d)}$$

$$\leq d_3(M) (|t_1 - t_2| + \|\alpha_1 - \alpha_2\|_{C^0(\bar{\Omega})}) (\|\mathbf{u}_D\|_{L^\infty(0, T; W^{1,4}(\Omega; \mathbb{R}^d))} + \|\mathbf{f}\|_{L^\infty(0, T; W^{-1,4}(\Omega; \mathbb{R}^d))}). \quad (6.38)$$



*Sketch of the proof.* Estimate (6.36) can be easily checked by elementary calculations, also making use of Korn's inequality (cf. also the calculations for [KRZ13, Lemma 2.4]). The existence of a minimizer standardly follows from the coercivity and lower semicontinuity properties of the functional  $J(t, \alpha, \cdot, p)$ , via the *direct method*; its uniqueness is due to the uniform convexity of  $J(t, \alpha, \cdot, p)$ . Estimates (6.37) and (6.38) rely on higher integrability results for weak solutions to elliptic equations in smooth domains, cf. e.g. [Giu03, Sec. 10.4]: we refer to the proofs of [KRZ13, Lemmas 2.4–2.5, Prop. 2.11] for all details.  $\square$

Let us highlight that (6.38) is a continuous dependence estimate for  $\mathbf{u}_{\min}$  with *fixed*  $p$ . It will play a crucial role for the study of the continuity properties of  $D_\alpha \mathcal{E}_3(\cdot, \cdot, p)$  in our next result.

**Lemma 6.11.** *Assume (6.24)–(6.27). Then:*

(1) *For every  $(\alpha, p) \in C^0(\overline{\Omega}; [0, 1]) \times L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$  the map  $t \mapsto \mathcal{E}_3(t, \alpha, p)$  is in  $C^1([0, T]; \mathbb{R})$  with*

$$\partial_t \mathcal{E}_3(t, \alpha, p) = \int_{\Omega} g(\alpha) \mathbb{C}(\varepsilon(\mathbf{u}_{\min}(t, \alpha, p) + \mathbf{u}_D(t)) - p) : \varepsilon(\mathbf{u}'_D(t)) \, dx - \langle \mathbf{f}'(t), \mathbf{u}_{\min}(t, \alpha, p) \rangle_{H^1(\Omega; \mathbb{R}^d)}, \quad (6.39)$$

*and there exists a constant  $d_4 > 0$  such that for all  $(t, \alpha, p) \in [0, T] \times C^0(\overline{\Omega}); [0, 1] \times L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$*

$$|\partial_t \mathcal{E}_3(t, \alpha, p)| \leq d_4 (\|\mathbf{u}_D\|_{C^1([0, T]; W^{1,4}(\Omega; \mathbb{R}^d))}^2 + \|\mathbf{f}\|_{C^1([0, T]; W_{\Gamma_D}^{-1,4}(\Omega; \mathbb{R}^d))}^2 + \|p\|_{L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})}^2). \quad (6.40)$$

*Moreover, there exist  $d_5 > 0$  depending on  $\|\mathbf{u}_D\|_{C^{1,1}([0, T]; W^{1,4}(\Omega; \mathbb{R}^d))}$ ,  $\|\mathbf{f}\|_{C^{1,1}([0, T]; W_{\Gamma_D}^{-1,4}(\Omega; \mathbb{R}^d))}$ , and such that for all  $(t_i, \alpha_i, p_i) \in [0, T] \times C^0(\overline{\Omega}; [0, 1]) \times L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$ ,  $i = 1, 2$ ,*

$$|\partial_t \mathcal{E}_3(t_1, \alpha_1, p_1) - \partial_t \mathcal{E}_3(t_2, \alpha_2, p_2)| \leq d_5 (|t_1 - t_2| + \|\alpha_1 - \alpha_2\|_{C^0(\overline{\Omega})} + \|p_1 - p_2\|_{L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})}). \quad (6.41)$$

(2) *For every  $t \in [0, T]$  and  $p \in L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$  the functional  $\alpha \mapsto \mathcal{E}_3(t, \alpha, p)$  is Gâteaux-differentiable on  $C^0(\overline{\Omega}; [0, 1])$  with Gâteaux derivative*

$$D_\alpha \mathcal{E}_3(t, \alpha, p)[\beta] = \int_{\Omega} \frac{1}{2} g'(\alpha) (\varepsilon(\mathbf{u}_{\min}(t, \alpha, p) + \mathbf{u}_D) - p) : \mathbb{C} : (\varepsilon(\mathbf{u}_{\min}(t, \alpha, p) + \mathbf{u}_D) - p) \beta \, dx \quad \text{for all } \beta \in C^0(\overline{\Omega}). \quad (6.42)$$

*If  $p \in L^4(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$ , then  $D_\alpha \mathcal{E}_3(t, \alpha, p)$  can be identified with an element in  $L^2(\Omega)$  and, moreover, there exists a constant  $d_6 > 0$ , depending on  $\|\mathbf{u}_D\|_{C^{1,1}([0, T]; W^{1,4}(\Omega; \mathbb{R}^d))}$ ,  $\|\mathbf{f}\|_{C^{1,1}([0, T]; W_{\Gamma_D}^{-1,4}(\Omega; \mathbb{R}^d))}$ , and  $p \in L^4(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$ , such that for every  $t_1, t_2 \in [0, T]$  and  $\alpha_1, \alpha_2 \in C^0(\overline{\Omega})$*

$$\|D_\alpha \mathcal{E}_3(t_1, \alpha_1, p) - D_\alpha \mathcal{E}_3(t_2, \alpha_2, p)\|_{L^2(\Omega)} \leq d_6 (|t_1 - t_2| + \|\alpha_1 - \alpha_2\|_{C^0(\overline{\Omega})}). \quad (6.43)$$

*Sketch of the proof.* We refer to [KRZ13, Lemma 2.6] for the proof of (6.39)–(6.41). In order to deduce formula (6.42), observe that

$$D_\alpha \mathcal{E}_3(t, \alpha, p)[\beta] = \partial_\alpha J(t, \alpha, u_{\min}(t, \alpha, p), p)[\beta] + \langle D_u J(t, \alpha, u_{\min}(t, \alpha, p), p), D_\alpha u_{\min}(t, \alpha, p)[\beta] \rangle_{H_{\Gamma_D}^1(\Omega; \mathbb{R}^d)}.$$

Here,  $\langle D_u J(t, \alpha, u_{\min}(t, \alpha, p), p), D_\alpha u_{\min}(t, \alpha, p)[\beta] \rangle_{H_{\Gamma_D}^1(\Omega; \mathbb{R}^d)} = 0$  in view of the Euler-Lagrange equation (here tested by  $D_\alpha u_{\min}(t, \alpha, p)[\beta]$ ) for the minimum problem (6.23). Straightforward calculation of the term  $\partial_\alpha J(t, \alpha, u_{\min}(t, \alpha, p), p)[\beta]$  results in (6.42). Moreover, in view of (6.37), one sees that  $D_\alpha \mathcal{E}_3(t, \alpha, p) \in L^2(\Omega)$  as soon as  $p \in L^4(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$ . The proof of (6.43) can be obtained by suitably adapting the calculations for [KRZ13, Lemma 2.8].  $\square$

Finally, we examine the continuity properties of  $\mathcal{E}_3$ .

**Lemma 6.12.** *Assume (6.24)–(6.27). Then,*

- (1) *for every  $p \in L^4(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$  the functional  $(t, \alpha) \mapsto \mathcal{E}_3(t, \alpha, p)$  is in  $C^{1,1}([0, T] \times C^0(\overline{\Omega}; [0, 1]))$ ;*  
(2) *for every sequence  $(t_n, \alpha_n, p_n) \subset [0, T] \times C^0(\overline{\Omega}; [0, 1]) \times L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$*

$$\left\{ \begin{array}{l} t_n \rightarrow t, \\ \alpha_n \rightarrow \alpha \quad \text{in } C^0(\overline{\Omega}), \\ p_n \rightarrow p \quad \text{in } L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d}) \end{array} \right\} \Rightarrow \mathcal{E}_3(t_n, \alpha_n, p_n) \rightarrow \mathcal{E}_3(t, \alpha, p). \quad (6.44)$$

*Sketch of the proof.* The Fréchet differentiability, with Lipschitz continuous differential, of  $\mathcal{E}_3(\cdot, \cdot, p)$  is a consequence of Lemma 6.11. To show (6.44), we first observe that, given a sequence  $(t_n, \alpha_n, p_n)_n$  as in (6.44), the sequence  $\mathbf{u}_n := \mathbf{u}_{\min}(t_n, \alpha_n, p_n)$ , bounded in  $H^1(\Omega; \mathbb{R}^d)$  by (6.36), up to a (not relabeled) subsequence weakly converges to some limit  $\mathbf{u}$  in  $H_{\Gamma_D}^1(\Omega; \mathbb{R}^d)$ . With standard lower semicontinuity arguments we find

$$\liminf_{n \rightarrow \infty} \mathcal{E}_3(t_n, \alpha_n, p_n) = \liminf_{n \rightarrow \infty} \mathcal{J}(t_n, \alpha_n, \mathbf{u}_n, p_n) \geq \mathcal{J}(t, \alpha, \mathbf{u}, p) \geq \mathcal{E}_3(t, \alpha, p).$$

We now show that  $\mathbf{u}$  is the unique minimizer of  $\mathcal{J}(t, \alpha, \cdot, p)$  by observing that

$$\mathcal{J}(t, \alpha, \mathbf{u}, p) \leq \limsup_{n \rightarrow \infty} \mathcal{J}(t_n, \alpha_n, \mathbf{u}_n, p_n) \stackrel{(1)}{\leq} \limsup_{n \rightarrow \infty} \mathcal{J}(t_n, \alpha_n, \mathbf{v}, p_n) \stackrel{(2)}{=} \mathcal{J}(t, \alpha, \mathbf{v}, p) \quad \forall \mathbf{v} \in H_{\Gamma_D}^1(\Omega; \mathbb{R}^d), \quad (6.45)$$

where (1) is due to  $\mathbf{u}_n \in \text{Argmin} \mathcal{J}(t_n, \alpha_n, \cdot, p_n)$ , and (2) is guaranteed by the convergence properties of  $(t_n, \alpha_n, p_n)_n$ . Choosing  $\mathbf{v} = \mathbf{u}$  in (6.45), we find that  $\lim_{n \rightarrow \infty} \mathcal{J}(t_n, \alpha_n, \mathbf{u}_n, p_n) = \mathcal{J}(t, \alpha, \mathbf{u}, p)$ , whence (6.44).  $\square$

**Properties of the energy  $\mathcal{E}$ .** We are now in the position to check that the energy functional  $\mathcal{E}$  complies with the conditions of Thm. 4.9.

**Proposition 6.13.** *Assume (6.24)–(6.27). Let  $\mathcal{E}$  be defined by (6.18), (6.19), (6.22), and (6.23) and, in addition, suppose that the constants  $C_{\mathbb{H}}$ ,  $c_0$ , and  $c_1$  from (6.22b), (6.24), and (6.25), respectively, fulfill*

$$c_0 c_1 > \frac{1}{C_{\mathbb{H}} + c_0 c_1}. \quad (6.46)$$

Then,

- (1)  $\mathcal{E}$  complies with Hypotheses 2.2 and 2.7, with the spaces  $\mathbf{U} = H^s(\Omega)$  and  $\mathbf{X} = H^1(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$ .
- (2) The mutual recovery sequence condition (6.35) is satisfied.
- (3) The Fréchet subdifferential  $\partial_{\alpha}^{-} \mathcal{E}(t, \alpha, p) : L^2(\Omega) \rightrightarrows L^2(\Omega)$  is given at any point  $(t, \alpha, p) \in \text{dom}(\partial_{\alpha}^{-} \mathcal{E})$  by

$$\partial_{\alpha}^{-} \mathcal{E}(t, \alpha, p) = A_s \alpha + \partial I_{[0,1]}(\alpha) + \gamma'(\alpha) + \frac{1}{2} g'(\alpha) (\varepsilon(\mathbf{u}_{\min}(t, \alpha, p) + \mathbf{u}_D) - p) : \mathbb{C} : (\varepsilon(\mathbf{u}_{\min}(t, \alpha, p) + \mathbf{u}_D) - p), \quad (6.47)$$

and the uniform Fréchet subdifferentiability condition from Hyp. 4.2 is satisfied.

- (4)  $\mathcal{E}$  complies with Hyp. 4.6.

*Proof. Ad (1):* Taking into account that the function  $\gamma$  is bounded from below on  $[0, 1]$ , the positive definiteness of the hardening tensor  $\mathbb{H}$ , and the previously proved (6.36), we find that  $\mathcal{E}$  is bounded from below. Indeed, from the hardening contribution one gains the positive term  $\frac{C_{\mathbb{H}}}{2} \|p\|_{L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})}^2$ , and it is sufficient to choose the constant  $\epsilon \in (0, c_0 c_1)$  in (6.36) such that  $C_{\mathbb{H}} + c_0 c_1 - \frac{1}{\epsilon} > 0$ . This is possible thanks to the compatibility condition (6.46). In this way, we also straightforwardly verify the coercivity property (2.19).

To check (2.6a) in Hyp. 2.7, let us consider a sequence  $(\alpha_n, p_n) \rightharpoonup (\alpha, p)$  in  $L^2(\Omega) \times L^1(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$  such that  $\liminf_{n \rightarrow \infty} \mathcal{E}(t, \alpha_n, p_n) < +\infty$ : up to a subsequence we may suppose that  $\sup_n \mathcal{E}(t, \alpha_n, p_n) < +\infty$ . Hence, by (2.19), we find that  $(\alpha_n, p_n) \rightharpoonup (\alpha, p)$  in  $H^s(\Omega) \times H^1(\Omega; \mathbb{R}_{\text{dev}}^{d \times d}) \Subset C^0(\bar{\Omega}) \times L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$ . Thus, by (6.44) we conclude that  $\mathcal{E}_3(t, \alpha_n, p_n) \rightarrow \mathcal{E}_3(t, \alpha, p)$ . The lower semicontinuity inequality for  $\mathcal{E}_1$  and  $\mathcal{E}_2$  is standard.

Clearly, for every  $(t, \alpha, p) \in [0, T] \times C^0(\bar{\Omega}; [0, 1]) \times L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$  we have  $\partial_t \mathcal{E}(t, \alpha, p) = \partial_t \mathcal{E}_3(t, \alpha, p)$ . Hence, (2.6c) is a consequence of (6.40), which gives a control of the power in terms of the energy, and of (6.41), yielding the upper semicontinuity of the power. This concludes the verification of Hypothesis 2.2.

**Ad (2):** Let  $(t_n, \alpha_n, p_n)_n \subset [0, T] \times L^2(\Omega) \times L^1(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$  be a sequence in the conditions of Hyp. 2.5. Thanks to Hyp. 2.7 and by the compact embeddings  $H^s(\Omega) \Subset C^0(\bar{\Omega})$  and  $H^1(\Omega; \mathbb{R}_{\text{dev}}^{d \times d}) \Subset L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$  we have  $(\alpha_n, p_n) \rightarrow (\alpha, p)$  in  $C^0(\bar{\Omega}) \times L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$ . Hence, for (6.35) we may choose the constant recovery sequence  $\tilde{p}_n := \tilde{p}$ , taking into account that

$$\begin{cases} \mathcal{R}(\bar{\alpha}_n, \tilde{p} - p_n) \rightarrow \mathcal{R}(\alpha, \tilde{p} - p) & \text{as } \bar{\alpha}_n \rightarrow \alpha \text{ in } C^0(\bar{\Omega}) \text{ and } p_n \rightarrow p \text{ in } L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d}), \\ \mathcal{E}(t_n, \alpha_n, \tilde{p}) \rightarrow \mathcal{E}(t, \alpha, p) & \text{by Lemma 6.12.} \end{cases}$$

**Ad (3):** The representation formula for  $\partial_{\alpha}^{-} \mathcal{E}$  ensues from the sum rule for the Fréchet subdifferential, taking into account that  $\mathcal{E}(t, \cdot, p)$  is given by the sum of either Gâteaux differentiable or convex terms. Next, we

observe that for every  $t \in [0, T]$ ,  $\alpha_1, \alpha_2 \in C^0(\bar{\Omega})$ , and  $p \in L^4(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$  and for every  $\xi_i \in \partial_{\alpha}^- \mathcal{E}(t, \alpha_i, p)$ ,  $i = 1, 2$ :

$$\begin{aligned} \int_{\Omega} (\xi_1 - \xi_2) (\alpha_1 - \alpha_2) dx &\geq a_s(\alpha_1 - \alpha_2, \alpha_1 - \alpha_2) + \int_{\Omega} (\gamma'(\alpha_1) - \gamma'(\alpha_2)) (\alpha_1 - \alpha_2) dx \\ &\quad + \int_{\Omega} (D_{\alpha} \mathcal{E}_3(t, \alpha_1, p) - D_{\alpha} \mathcal{E}_3(t, \alpha_2, p)) (\alpha_1 - \alpha_2) dx \\ &\geq \|\alpha_1 - \alpha_2\|_{H^s(\Omega)}^2 - \|\alpha_1 - \alpha_2\|_{L^2(\Omega)}^2 - C \|\alpha_1 - \alpha_2\|_{L^2(\Omega)}^2 - d_6 \|\alpha_1 - \alpha_2\|_{C^0(\Omega)} \|\alpha_1 - \alpha_2\|_{L^2(\Omega)}, \end{aligned}$$

where the first inequality is due to the monotonicity of  $\partial I_{[0,1]}$ , and the second one follows from adding and subtracting  $\|\alpha_1 - \alpha_2\|_{L^2(\Omega)}^2$ , since  $\sqrt{\|\alpha\|_{L^2(\Omega)}^2 + a_s(\alpha, \alpha)}$  gives the  $H^s(\Omega)$ -norm, from using that  $\gamma$  is Lipschitz continuous on  $[0, 1]$ , and from estimate (6.43). Using now the continuous embedding  $H^s(\Omega) \Subset C^0(\bar{\Omega})$  and Young's inequality to estimate the very last term in the above formula, we conclude that

$$\begin{aligned} \exists d_7, > 0 \forall (t, p) \in [0, T] \times L^4(\Omega; \mathbb{R}_{\text{dev}}^{d \times d}) \forall \alpha_1, \alpha_2 \in H^s(\Omega) \forall \xi_i \in \partial_{\alpha}^- \mathcal{E}(t, \alpha_i, p), i = 1, 2 : \\ \int_{\Omega} (\xi_1 - \xi_2) (\alpha_1 - \alpha_2) dx \geq \frac{1}{2} \|\alpha_1 - \alpha_2\|_{H^s(\Omega)}^2 - d_7 \|\alpha_1 - \alpha_2\|_{L^2(\Omega)}^2. \end{aligned} \quad (6.48)$$

It is standard to deduce from (6.48) that  $\mathcal{E}$  complies with the  $\lambda$ -convexity (4.7), whence Hyp. 4.2.

**Ad (4):** Hypothesis 4.6 can be verified with the choice of the constant recovery sequence  $\tilde{\alpha}_n := \tilde{\alpha}$ , taking into account that, for a sequence  $(t_n, \alpha_n, p_n)_n$  as in Hyp. 4.6,

$$\begin{aligned} \limsup_{n \rightarrow \infty} (\mathcal{E}(t_n, \tilde{\alpha}, p_n) - \mathcal{E}(t_n, \alpha_n, p_n)) &= \limsup_{n \rightarrow \infty} (\mathcal{E}_1(\tilde{\alpha}) + \mathcal{E}_3(t_n, \tilde{\alpha}, p_n) - \mathcal{E}_1(\alpha_n) - \mathcal{E}_3(t_n, \alpha, p_n)) \\ &\leq \mathcal{E}_1(\tilde{\alpha}) + \mathcal{E}_3(t, \tilde{\alpha}, p) - \mathcal{E}_1(\alpha) - \mathcal{E}_3(t, \alpha, p) = \mathcal{E}(t, \tilde{\alpha}, p) - \mathcal{E}(t, \alpha, p) \end{aligned}$$

thanks to (6.44) and the lower semicontinuity properties of  $\mathcal{E}_1$  and  $\mathcal{E}_3$ .  $\square$

Hence, from Theorem 4.9 we conclude

**Theorem 6.14.** *Assume (6.24)–(6.27), and (6.46). Then, for every  $(\alpha_0, p_0) \in H^s(\Omega) \cap H^1(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$  there exists an enhanced semistable energetic solution  $(\alpha, p)$ , with*

$$\alpha \in L^\infty(0, T; H^s(\Omega)) \cap H^1(0, T; L^2(\Omega)) \quad \text{and} \quad p \in L^\infty(0, T; H^1(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})) \cap \text{BV}([0, T]; L^1(\Omega; \mathbb{R}_{\text{dev}}^{d \times d}))$$

to the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  given by (6.18), (6.28), and (6.29) such that  $\alpha(0) = \alpha_0$  and  $p(0) = p_0$ .

## 7. PROOFS OF THEOREMS 4.9, 4.13, 5.4, AND 5.6

In the proof of our existence results for the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  we will reverse the order in Section 4 and indeed start with the case of a general dissipation potential  $\mathcal{V}$  in Sec. 7.1. In this general frame, on the one hand, we will employ *variational* techniques tailored to the proof of the discrete energy-dissipation inequality. On the other hand, in the limit passage to the time-continuous limit we will resort to Young measure arguments that will play a major role also in the particular case of quadratic  $\mathcal{V}$ , cf. Sec. 7.2. Finally, in Sec. 7.3 we will address the proofs of Theorems 5.4 ( $\mathcal{V}$  general) and 5.6 ( $\mathcal{V}$  quadratic).

**7.1. Proof of Theorem 4.13.** For the time-discrete analysis of the rate-dependent differential inclusion (3.2), we will adapt the arguments from the proof of [MRS13b, Thm. 4.4], see also [RS06]. More precisely, we will start by providing some “stationary estimates” for the solutions of the time-incremental minimum problem (4.22a), yielding the discrete values  $(u_\tau^n)_{n=1}^{N_\tau}$ , at each fixed time-step. In particular, in Lemma 7.1 below, we will give the crucial time-discrete energy-dissipation inequality (7.5). We shall combine it with the estimates associated with the minimum problem (4.22b), yielding the discrete values  $(z_\tau^n)_{n=1}^{N_\tau}$ , to derive the a priori estimates on the approximate solutions in Proposition 7.2. Also relying on Young measure compactness arguments, in Prop. 7.3 we derive a series of convergences (along suitable subsequences) of the discrete solutions to a pair  $(u, z)$ , which we will then show to be a *weak semistable energetic solution* to the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$ .

Preliminarily, we fix some properties of the minimization problem

$$\mathcal{J}_r(\bar{t}, \bar{u}, \bar{z}) := \inf_{u \in \mathbf{D}_u} \left( r \mathcal{V} \left( \frac{u - \bar{u}}{r} \right) + \mathcal{E}(\bar{t} + r, u, \bar{z}) \right), \quad \text{for given } (\bar{t}, \bar{u}, \bar{z}) \in [0, T] \times \mathbf{D}_u \times \mathbf{D}_z \text{ and } 0 < r < T - \bar{t}, \quad (7.1)$$

in Lemma 7.1 below, which we recall here from [MRS13b] for the reader's convenience. Observe that it is in the proof of this Lemma, in particular in the derivation of the energy-dissipation inequality (7.5), that condition (4.21) on  $\mathcal{V}$  comes into play, cf. the proof of [MRS13b, Lemma 6.1].

**Lemma 7.1.** [MRS13b, Lemma 6.1] *Under Hypotheses 2.2, 2.3, 2.7, condition (4.21) on  $\mathcal{V}$ , and Hyp. 4.12, for every  $(\bar{t}, \bar{u}, \bar{z}) \in [0, T] \times \mathbf{D}_u \times \mathbf{D}_z$  and  $0 < r < \min\{\tau_o, T - \bar{t}\}$  the set*

$$\mathbf{M}_r(\bar{t}, \bar{u}, \bar{z}) := \underset{u \in \mathbf{D}_u}{\text{Argmin}} \left( r\mathcal{V} \left( \frac{u - \bar{u}}{r} \right) + \mathcal{E}(\bar{t} + r, u, \bar{z}) \right) \neq \emptyset,$$

and there exists a measurable selection  $r \in (0, T - \bar{t}) \mapsto u_r \in \mathbf{M}_r(\bar{t}, \bar{u}, \bar{z})$  fulfilling for every  $r \in (0, T - \bar{t})$  the Euler-Lagrange equation

$$\partial\mathcal{V} \left( \frac{u_r - \bar{u}}{r} \right) + \partial_u^- \mathcal{E}(\bar{t} + r, u_r, \bar{z}) \ni 0 \quad \text{in } \mathbf{V}^*, \quad (7.2)$$

as well as

$$\exists C > 0 \quad \forall r \in (0, T - \bar{t}) : \quad \mathcal{G}(u_r, \bar{z}) \leq C\mathcal{G}(\bar{u}, \bar{z}), \quad (7.3)$$

$$\lim_{r \downarrow 0} \|u_r - \bar{u}\|_{\mathbf{V}} = 0, \quad \lim_{r \downarrow 0} \mathcal{J}_r(\bar{t}, \bar{u}, \bar{z}) = \mathcal{E}(\bar{t}, \bar{u}, \bar{z}), \quad (7.4)$$

and complying with the energy-dissipation inequality

$$r_0\mathcal{V} \left( \frac{u_{r_0} - \bar{u}}{r_0} \right) + \int_0^{r_0} \mathcal{V}^*(-\xi_r) dr + \mathcal{E}(\bar{t} + r_0, u_{r_0}, \bar{z}) \leq \mathcal{E}(\bar{t}, \bar{u}, \bar{z}) + \int_0^{r_0} \partial_t \mathcal{E}(\bar{t} + r, u_r, \bar{z}) dr \quad (7.5)$$

for any  $r_0 \in (0, T - \bar{t})$ , with  $\xi_r$  any selection in  $\partial_u^- \mathcal{E}(\bar{t} + r, u_r, \bar{z}) \cap (-\partial\mathcal{V}(\frac{u_r - \bar{u}}{r}))$ .

While referring to [MRS13b] for the proof, let us only comment here on the Euler-Lagrange equation: it is an immediate consequence of [MRS13b, Prop. 4.2], which can be applied thanks to Hypothesis 4.12.

Exploiting (7.5), we will now derive the discrete version (7.8) below of the energy-dissipation inequality (3.4), whence all of the a priori estimates on the approximate solutions arising from the time-discrete scheme in Problem 4.11. The Euler-Lagrange equation (7.2) rewrites for the variational interpolant  $\tilde{u}_\tau$  from (4.23) as

$$\partial\mathcal{V} \left( \frac{\tilde{u}_\tau(t) - u_\tau(t)}{t - \underline{t}_\tau(t)} \right) + \partial_u^- \mathcal{E}(t, \tilde{u}_\tau(t), \underline{z}_\tau(t)) \ni 0 \quad \text{in } \mathbf{V}^* \text{ for a.a. } t \in (0, T).$$

In what follows, we will denote by  $\tilde{\xi}_\tau : (0, T) \rightarrow \mathbf{V}^*$  a measurable map fulfilling

$$\tilde{\xi}_\tau(t) \in \partial_u^- \mathcal{E}(t, \tilde{u}_\tau(t), \underline{z}_\tau(t)), \quad -\tilde{\xi}_\tau(t) \in \partial\mathcal{V} \left( \frac{\tilde{u}_\tau(t) - u_\tau(t)}{t - \underline{t}_\tau(t)} \right) \quad \text{for a.a. } t \in (0, T). \quad (7.6)$$

For later use, we also point out that the minimum problem (4.22a) yields

$$\bar{\xi}_\tau(t) \in \partial_u^- \mathcal{E}(t, \bar{u}_\tau(t), \underline{z}_\tau(t)), \quad -\bar{\xi}_\tau(t) \in \partial\mathcal{V}(u'_\tau(t)) \quad \text{for a.a. } t \in (0, T), \quad (7.7)$$

with  $\bar{\xi}_\tau : (0, T) \rightarrow \mathbf{V}^*$  the piecewise constant interpolant of the elements  $(\xi_\tau^n)_{n=1}^{N_\tau}$  fulfilling the Euler-Lagrange equation for (4.22a).

**Proposition 7.2.** *Assume Hypotheses 2.2, 2.3, 2.7, and condition (4.21) on  $\mathcal{V}$ .*

*Then, the interpolants  $(\bar{u}_\tau, \underline{u}_\tau, u_\tau, \tilde{u}_\tau, \bar{z}_\tau, \underline{z}_\tau, z_\tau)$  and  $(\tilde{\xi}_\tau)_\tau$  of the discrete solutions to Problem 4.11 comply with the discrete energy-dissipation inequality*

$$\begin{aligned} & \int_{\bar{\tau}_\tau(s)}^{\bar{\tau}_\tau(t)} \mathcal{V}(u'_\tau(r)) + \mathcal{V}^*(-\tilde{\xi}_\tau(r)) dr + \int_{\bar{\tau}_\tau(s)}^{\bar{\tau}_\tau(t)} \mathcal{R}(z'_\tau(r)) dr + \mathcal{E}(\bar{\tau}_\tau(t), \bar{u}_\tau(t), \bar{z}_\tau(t)) \\ & \leq \mathcal{E}(\bar{\tau}_\tau(s), \bar{u}_\tau(s), \bar{z}_\tau(s)) + \int_{\bar{\tau}_\tau(s)}^{\bar{\tau}_\tau(t)} \partial_t \mathcal{E}(r, \tilde{u}_\tau(r), \underline{z}_\tau(r)) dr, \end{aligned} \quad (7.8)$$

and the discrete semistability condition

$$\mathcal{E}(\bar{\tau}_\tau(t), \bar{u}_\tau(t), \bar{z}_\tau(t)) \leq \mathcal{E}(\bar{\tau}_\tau(t), \bar{u}_\tau(t), \tilde{z}) + \mathcal{R}(\tilde{z} - \bar{z}_\tau(t)) \quad \text{for all } \tilde{z} \in \mathbf{Z} \text{ and for all } t \in (0, T]. \quad (7.9)$$

Moreover, there exists  $C > 0$  such that for every  $0 < \tau < \tau_o$  the following estimates hold

$$\begin{aligned} \sup_{t \in (0, T)} \left( \mathfrak{G}(\bar{u}_\tau(t), \bar{z}_\tau(t)) + \tau_o \mathcal{V} \left( \frac{\bar{u}_\tau(t) - u_0}{\tau_o} \right) + \mathfrak{R}(\bar{z}_\tau(t) - z_0) \right) &\leq C, \\ \sup_{t \in (0, T)} \left( \mathfrak{G}(\tilde{u}_\tau(t), \tilde{z}_\tau(t)) + \tau_o \mathcal{V} \left( \frac{\tilde{u}_\tau(t) - u_0}{\tau_o} \right) + \mathfrak{R}(\tilde{z}_\tau(t) - z_0) \right) &\leq C, \end{aligned} \quad (7.10a)$$

and analogously for  $\sup_{t \in (0, T)} \mathfrak{G}(\bar{u}_\tau(t), \tilde{z}_\tau(t)) + \tau_o \mathcal{V}((\bar{u}_\tau(t) - u_0)/\tau_o)$ ;

$$\sup_{t \in (0, T)} |\partial_t \mathcal{E}(t, \tilde{u}_\tau(t), \tilde{z}_\tau(t))| \leq C; \quad (7.10b)$$

$$\int_0^T \mathcal{V}(u'_\tau(r)) dr \leq C, \quad \int_0^T \mathcal{V}^*(-\tilde{\xi}_\tau(r)) dr \leq C, \quad \int_0^T \mathfrak{R}(z'_\tau(r)) dr \leq C; \quad (7.10c)$$

the families  $(u'_\tau)_\tau \subset L^1(0, T; \mathbf{V})$  and  $(\tilde{\xi}_\tau)_\tau \subset L^1(0, T; \mathbf{V}^*)$  are equi-integrable, and

$$\sup_{t \in (0, T)} \|\bar{u}_\tau(t) - \underline{u}_\tau(t)\|_{\mathbf{V}} + \sup_{t \in (0, T)} \|u_\tau(t) - \underline{u}_\tau(t)\|_{\mathbf{V}} + \sup_{t \in (0, T)} \|\tilde{u}_\tau(t) - \underline{u}_\tau(t)\|_{\mathbf{V}} = o(1) \text{ as } \tau \downarrow 0. \quad (7.10d)$$

*Proof.* The discrete semistability (7.9) is a direct consequence of the minimum problem (4.22b) and of the triangle inequality satisfied by  $\mathfrak{R}$ . The proof of the remaining items in the statement follows the lines of [MRS13b, Prop. 6.3]. Therefore we shall just outline its main steps, referring to [MRS13b] for all details.

In order to derive (7.8), let us first of all fix two nodes  $t_{n-1}, t_n \in \mathcal{P}_\tau$  and write (7.5) with  $\bar{t} = t_{n-1}$ ,  $\bar{u} = u_\tau^{n-1}$ ,  $\bar{z} = z_\tau^{n-1}$ ,  $r_0 = t - t_{n-1}$ ,  $u_{r_0} = \tilde{u}_\tau(t)$ ,  $u_r = \tilde{u}_\tau(r)$ , and  $\xi_r = \tilde{\xi}_\tau(r)$ , for  $r \in (t_{n-1}, t)$ . Thus we obtain with a change of variables in the second integral on the left-hand side

$$\begin{aligned} (t - t_{n-1}) \mathcal{V} \left( \frac{\tilde{u}_\tau(t) - \underline{u}_\tau(t)}{t - t_{n-1}} \right) + \int_{t_{n-1}}^t \mathcal{V}^*(-\tilde{\xi}_\tau(r)) dr + \mathcal{E}(t, \tilde{u}_\tau(t), \tilde{z}_\tau(t)) \\ \leq \mathcal{E}(t_{n-1}, \underline{u}_\tau(t), \tilde{z}_\tau(t)) + \int_{t_{n-1}}^t \partial_t \mathcal{E}(r, \tilde{u}_\tau(r), \tilde{z}_\tau(r)) dr, \end{aligned} \quad (7.11)$$

whence, for  $t = t_n$ ,

$$\int_{t_{n-1}}^{t_n} \mathcal{V}(u'_\tau(r)) + \mathcal{V}^*(-\tilde{\xi}_\tau(r)) dr + \mathcal{E}(t_n, u_\tau^n, z_\tau^{n-1}) \leq \mathcal{E}(t_{n-1}, u_\tau^{n-1}, z_\tau^{n-1}) + \int_{t_{n-1}}^{t_n} \partial_t \mathcal{E}(r, \tilde{u}_\tau(r), \tilde{z}_\tau(r)) dr. \quad (7.12)$$

On the other hand, we deduce from (4.22b) that

$$\tau \mathcal{R} \left( \frac{z_\tau^n - z_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, u_\tau^n, z_\tau^n) \leq \mathcal{E}(t_n, u_\tau^n, z_\tau^{n-1}).$$

Adding this to (7.12), due to the cancelation of the term  $\mathcal{E}(t_n, u_\tau^n, z_\tau^{n-1})$  we obtain

$$\begin{aligned} \int_{t_{n-1}}^{t_n} \mathcal{V}(u'_\tau(r)) + \mathcal{V}^*(-\tilde{\xi}_\tau(r)) dr + \tau \mathcal{R} \left( \frac{z_\tau^n - z_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, u_\tau^n, z_\tau^n) \\ \leq \mathcal{E}(t_{n-1}, u_\tau^{n-1}, z_\tau^{n-1}) + \int_{t_{n-1}}^{t_n} \partial_t \mathcal{E}(r, \tilde{u}_\tau(r), \tilde{z}_\tau(r)) dr \end{aligned}$$

which gives (7.8) upon summing up over the index  $n$ .

Using the power control condition from (2.6c) and arguing in the very same way as in the proof of [MRS13b, Prop. 6.3], one infers that

$$\sup_{t \in (0, T)} \left( \mathfrak{G}(\bar{u}_\tau(t), \bar{z}_\tau(t)) + \int_0^{\bar{t}_\tau(t)} \mathcal{V}(u'_\tau(s)) ds \right) \leq C. \quad (7.13)$$

In order to conclude the first of (7.10), we proceed in the following way: For  $t_0 \in (0, T)$  fixed,  $1 \leq n_0 \leq N_\tau$  fulfill  $\bar{t}_\tau(t_0) = n_0 \tau$ ; suppose that  $n_0 \tau \leq \tau_o$ , with  $\tau_o > 0$  from Hyp. 2.7. Then,

$$\int_0^{\bar{t}_\tau(t_0)} \mathcal{V}(u'_\tau(s)) ds = n_0 \tau \sum_{n=1}^{n_0} \frac{1}{n_0} \mathcal{V} \left( \frac{u_\tau^n - u_\tau^{n-1}}{\tau} \right) \stackrel{(1)}{\geq} \bar{t}_\tau(t_0) \mathcal{V} \left( \sum_{n=1}^{n_0} \frac{u_\tau^n - u_\tau^{n-1}}{\bar{t}_\tau(t_0)} \right) \stackrel{(2)}{\geq} \tau_o \mathcal{V} \left( \frac{\bar{u}_\tau(t_0) - u_0}{\tau_o} \right),$$

where (1) follows from the convexity of  $\mathcal{V}$ , and (2) from the fact that for every  $v \in \mathbf{V}$  the map  $\tau \mapsto \mathcal{V}\left(\frac{v}{\tau}\right)$  is nonincreasing. Then, from (7.13) we conclude that  $\tau_o \mathcal{V}\left(\frac{\bar{u}_\tau(t_0) - u_0}{\tau_o}\right) \leq C$  for  $t_0 \in (0, T)$  sufficiently small such that  $\bar{\tau}_\tau(t_0) \leq \tau_o$ . Then, the estimate

$$\sup_{t \in (0, T)} \tau_o \mathcal{V}\left(\frac{\bar{u}_\tau(t) - u_0}{\tau_o}\right) \leq C \quad (7.14)$$

can be concluded with a standard argument, dividing  $[0, T]$  into subintervals of length less or equal than  $\tau_o$ . The calculations for the estimate of the  $\mathcal{R}$ -dissipation are simpler by 1-homogeneity, and we thus conclude the first of (7.10a). The second of (7.10a) analogously ensues from (7.11). Let us only finally comment on the bound for  $\sup_{t \in [0, T]} \mathcal{G}(\bar{u}_\tau(t), \bar{z}_\tau(t))$  in (7.10a): From (4.22a) we deduce for every  $t \in [0, T]$

$$\mathcal{E}(\bar{\tau}_\tau(t), \bar{u}_\tau(t), \bar{z}_\tau(t)) \leq \mathcal{E}(\bar{\tau}_\tau(t), \underline{u}_\tau(t), \underline{z}_\tau(t)) = \mathcal{E}(\underline{\tau}_\tau(t), \underline{u}_\tau(t), \underline{z}_\tau(t)) + \int_{\underline{\tau}_\tau(t)}^{\bar{\tau}_\tau(t)} \partial_t \mathcal{E}(s, \underline{u}_\tau(t), \underline{z}_\tau(t)) ds \leq C,$$

where the last estimate ensues from the bound for  $\sup_{t \in [0, T]} \mathcal{G}(\bar{u}_\tau(t), \bar{z}_\tau(t))$ , and from (2.6c).  $\square$

The next result shall also involve the Young measure limit of the family  $(\tilde{\xi}_\tau)_\tau$ . We refer to the Appendix for a self-contained exposition of the Young measure compactness result underlying the proof of Proposition 7.3.

**Proposition 7.3.** *Assume Hypotheses 2.2, 2.3, 2.7, and the closedness (4.24a) from Hypothesis 4.12.*

*Then, for every null sequence  $(\tau_k)_k$  of time-steps there exist a (not relabeled) subsequence, a pair  $(u, z)$  as in (3.1), a function  $\mathcal{E} \in \text{BV}([0, T])$ , and a time-dependent Young measure  $\boldsymbol{\mu} = (\mu_t)_{t \in (0, T)} \in \mathcal{Y}(0, T; \mathbf{V}^*)$  such that convergences (4.15) and (4.25) hold as  $k \rightarrow \infty$  and, in addition*

$$\begin{cases} \mathcal{E}(t) = \lim_{k \rightarrow \infty} \mathcal{E}(\bar{\tau}_{\tau_k}(t), \bar{u}_{\tau_k}(t), \bar{z}_{\tau_k}(t)) & \text{for all } t \in [0, T], \\ \mathcal{E}(t) \leq \liminf_{k \rightarrow \infty} \mathcal{E}(\bar{\tau}_{\tau_k}(t), \bar{u}_{\tau_k}(t), \bar{z}_{\tau_k}(t)) & \text{for almost all } t \in (0, T), \\ \mathcal{E}(t) \geq \mathcal{E}(t, u(t), z(t)) & \text{for all } t \in [0, T], \end{cases} \quad (7.15)$$

while  $\boldsymbol{\mu}$  is the limit Young measure of  $(\tilde{\xi}_{\tau_k})_k$ , whence

$$\tilde{\xi}_{\tau_k} \rightharpoonup \xi \quad \text{in } L^1(0, T; \mathbf{V}^*) \quad \text{with } \xi(t) = \int_{\mathbf{V}^*} \zeta d\mu_t(\zeta) \quad \text{for a.a. } t \in (0, T). \quad (7.16)$$

Moreover, the following energy-dissipation inequality holds for all  $0 \leq s \leq t \leq T$

$$\begin{aligned} & \int_s^t \mathcal{V}(u'(r)) dr + \int_s^t \int_{\mathbf{V}^*} \mathcal{V}^*(-\zeta) d\mu_r(\zeta) dr + \text{Var}_{\mathcal{R}}(z; [s, t]) + \mathcal{E}(t) \\ & \leq \mathcal{E}(s) + \int_s^t \partial_t \mathcal{E}(r, u(r), z(r)) dr. \end{aligned} \quad (7.17)$$

*Proof.* Convergences (4.15a)–(4.15c) ensue from estimates (7.10) via standard compactness arguments, cf. [Sim87], also taking into account the coercivity from Hyp. 2.7. The pointwise  $\sigma_{\mathbf{X}}$ -convergence for  $(\bar{z}_{\tau_k})_k$  (which then implies (4.15g) thanks to (2.5b)), is a consequence of the bounds in  $L^\infty(0, T; \mathbf{X}) \cap \text{BV}([0, T]; \mathbf{Z})$  via a Helly-type argument (cf. e.g. [MT04, Thm. 6.1]), which also gives

$$\text{Var}_{\mathcal{R}}(z; [s, t]) \leq \liminf_{k \rightarrow \infty} \int_s^t \mathcal{R}(z'_{\tau_k}(r)) dr,$$

whence  $z \in \text{BV}([0, T]; \mathbf{Z})$ . Finally, the ‘stability’ property (4.15f) ensues from a standard argument, cf. e.g. the proof of [RTP15, Thm. 4.1]. With (4.15f) at hand, it is then easy to deduce convergences (4.15d) and (4.15e) for  $(z_{\tau_k})_k$  as well. We refer to the proof of [MRS13b, Prop. 6.4] for all details on the first and third inequalities in (7.15) and on (7.16). The limit passage in (7.8), leading to (7.17), also follows from the arguments for [MRS13b, Prop. 6.4], combined with the observation that

$$\liminf_{k \rightarrow \infty} \int_{\bar{\tau}_{\tau_k}(s)}^{\bar{\tau}_{\tau_k}(t)} \mathcal{R}(z'_{\tau_k}(r)) dr \geq \text{Var}_{\mathcal{R}}(z; [s, t])$$

for every  $0 \leq s \leq t \leq T$ , thanks to (4.15g). Finally, let us comment on the second of (7.15). It ensues from the minimum problem (4.22b), yielding

$$\mathcal{E}(\bar{\tau}_{\tau_k}(t), \bar{u}_{\tau_k}(t), \bar{z}_{\tau_k}(t)) \leq \mathcal{E}(\bar{\tau}_{\tau_k}(t), \bar{u}_{\tau_k}(t), z_{\tau_k}(t)) \quad \text{for all } t \in [0, T].$$

For later use, observe also that (2.6c) yields

$$\limsup_{\tau_k \downarrow 0} \partial_t \mathcal{E}(t, \tilde{u}_{\tau_k}(t), \tilde{z}_{\tau_k}(t)) \leq \partial_t \mathcal{E}(t, u(t), z(t)) \quad \text{for a.a. } t \in (0, T). \quad (7.18)$$

□

We are now in the position to establish the **proof of Theorem 4.13**. In order to obtain the energy-dissipation inequality (3.4) we need to gain further insight into the Young measure energy-dissipation inequality (7.17), and in particular into the properties of the limit  $\boldsymbol{\mu} = (\mu_t)_{t \in (0, T)}$  of  $(\tilde{\xi}_{\tau_k})$ . It follows from Theorem A.2 ahead that for almost all  $t \in (0, T)$  the measure  $\mu_t$  is concentrated on the limit points of  $(\tilde{\xi}_{\tau_k}(t))_k$  with respect to the weak topology of  $\mathbf{V}^*$ . Now, due to the first of (7.6),  $\tilde{\xi}_{\tau_k}(t) \in \partial_u^- \mathcal{E}(t, \tilde{u}_{\tau_k}(t), \tilde{z}_{\tau_k}(t))$  for almost all  $t \in (0, T)$ . Combining the pointwise convergences (4.15c) and (4.15g) for  $(\tilde{u}_{\tau_k}(t))_k$  and  $(\tilde{z}_{\tau_k}(t))_k$  with the closedness condition (4.24a), we ultimately conclude that

$$\text{the support of } \mu_t \text{ is a subset of } \partial_u^- \mathcal{E}(t, u(t), z(t)) \text{ for almost all } t \in (0, T). \quad (7.19)$$

Since the latter set is closed and convex, we conclude that

$$\xi(t) = \int_{\mathbf{V}^*} \zeta d\mu_t(\zeta) \in \partial_u^- \mathcal{E}(t, u(t), z(t)) \quad \text{for a.a. } t \in (0, T). \quad (7.20)$$

Then, by Jensen's inequality we have

$$\int_s^t \int_{\mathbf{V}^*} \mathcal{V}^*(-\zeta) d\mu_r(\zeta) dr \geq \int_s^t \mathcal{V}^*(-\xi(r)) dr \quad \text{for every } 0 \leq s \leq t \leq T. \quad (7.21)$$

Hence, in view of (7.21) and (7.18) we are in the position to deduce from (7.17), written on the interval  $(0, t)$ , the energy-dissipation inequality (3.4).

Furthermore, observe that, when it holds, the enhanced closedness/continuity condition (4.24b) guarantees

$$\lim_{k \rightarrow \infty} \mathcal{E}(\bar{\tau}_{\tau_k}(t), \bar{u}_{\tau_k}(t), \bar{z}_{\tau_k}(t)) = \mathcal{E}(t) = \mathcal{E}(t, u(t), z(t)) \quad \text{for almost all } t \in (0, T). \quad (7.22)$$

To check this, from estimate (7.10c) and Fatou's Lemma we deduce that  $\liminf_{k \rightarrow \infty} \|u'_{\tau_k}(t)\|_{\mathbf{V}} < +\infty$  for almost all  $t \in (0, T)$ . Since  $\partial \mathcal{V} : \mathbf{V} \rightrightarrows \mathbf{V}^*$  is a bounded operator, we then have

$$\liminf_{k \rightarrow \infty} \|\bar{\xi}_{\tau_k}(t)\|_{\mathbf{V}^*} < +\infty \quad \text{for a.a. } t \in (0, T), \quad (7.23)$$

where  $t \mapsto \bar{\xi}_{\tau_k}(t)$  is a selection in  $(-\partial \mathcal{V}(u'_{\tau_k}(t))) \cap \partial^- \mathcal{E}(t, \bar{u}_{\tau_k}(t), \bar{z}_{\tau_k}(t))$  fulfilling the Euler Lagrange equation (7.7). We are thus in the position to apply condition (4.24b) (also in view of (7.10a)), thus concluding that

$$\liminf_{k \rightarrow \infty} \mathcal{E}(\bar{\tau}_{\tau_k}(t), \bar{u}_{\tau_k}(t), \bar{z}_{\tau_k}(t)) = \mathcal{E}(t, u(t), z(t)) \quad \text{for a.a. } t \in (0, T). \quad (7.24)$$

But then, by the second and the third of (7.15), we infer  $\mathcal{E}(t, u(t), z(t)) \leq \mathcal{E}(t) \leq \mathcal{E}(t, u(t), z(t))$  for almost all  $t \in (0, T)$ , whence (7.22). Therefore, from (7.17) we gather the enhanced energy-dissipation inequality (4.20).

Finally, with a standard procedure in the analysis of rate-independent processes we pass to the limit in the discrete semistability (7.9) by resorting to the mutual recovery sequence condition (2.17). More detailed, for all  $t \in (0, T)$  we apply Hypothesis 2.5 to the sequences  $(\bar{u}_{\tau_k}(t), \bar{z}_{\tau_k}(t))_k$ , converging to  $(u(t), z(t))$  as in (4.15c) and (4.15g), and thus for every  $\tilde{z} \in \mathbf{Z}$  we construct a sequence  $\tilde{z}_k \rightarrow \tilde{z}$  in  $\mathbf{Z}$  such that (2.17) holds. Exploiting (2.17), we thus conclude that the semistability inequality (3.3) holds with  $\tilde{z}$ . This concludes the proof. ■

**7.2. Proof of Theorem 4.9.** Preliminarily, observe that from (4.3b) we conclude that

$$0 \in \partial^- \left( \tau \mathcal{V} \left( \frac{\cdot - u_{\tau}^{n-1}}{\tau} \right) + \mathcal{E}(t_n, \cdot, z_{\tau}^n) \right) (u_{\tau}^n) = \partial \mathcal{V} \left( \frac{u_{\tau}^n - u_{\tau}^{n-1}}{\tau} \right) + \partial_u^- \mathcal{E}(t_n, u_{\tau}^n, z_{\tau}^n), \quad (7.25)$$

where the last equality follows from the sum rule for the Fréchet subdifferential, since  $\mathcal{V}$  is Fréchet differentiable. Hence we obtain the Euler-Lagrange equation

$$\partial \mathcal{V} \left( \frac{u_{\tau}^n - u_{\tau}^{n-1}}{\tau} \right) + \xi_{\tau}^n \ni 0, \quad \xi_{\tau}^n \in \partial_u^- \mathcal{E}(t_n, u_{\tau}^n, z_{\tau}^n) \quad \text{for all } n = 1, \dots, N_{\tau}. \quad (7.26)$$

We denote by  $\bar{\xi}_{\tau}$  the piecewise constant interpolant of the elements  $(\xi_{\tau}^n)_{n=1}^{N_{\tau}} \subset \mathbf{V}^*$ , so that (7.26) rewrites as

$$\partial \mathcal{V}(u'_{\tau}(t)) + \bar{\xi}_{\tau}(t) \ni 0, \quad \bar{\xi}_{\tau}(t) \in \partial_u^- \mathcal{E}(\bar{\tau}_{\tau}(t), \bar{u}_{\tau}(t), \bar{z}_{\tau}(t)) \quad \text{for a.a. } t \in (0, T). \quad (7.27)$$

We start with the counterpart to Proposition 7.2. Observe that the result below holds for *any* dissipation potential  $\mathcal{V}$ , not necessarily quadratic. Instead, the *uniform Fréchet subdifferentiability* from Hypothesis 4.2 plays here a key role.

**Proposition 7.4.** *Assume Hypotheses 2.2, 2.3, 2.7, and 4.2. Let  $(\bar{u}_\tau, \underline{u}_\tau, u_\tau, \bar{z}_\tau, z_\tau, z_\tau)_\tau$  and  $(\bar{\xi}_\tau)_\tau$  be the approximate solutions constructed from the time-discretization scheme in Problem 4.1.*

*Then, there exists  $C > 0$  such that for every  $0 < \tau < \tau_o$  the following estimates hold*

$$\sup_{t \in (0, T)} \left( \mathcal{G}(\bar{u}_\tau(t), \bar{z}_\tau(t)) + \tau_o \mathcal{V} \left( \frac{\bar{u}_\tau(t) - u_0}{\tau_o} \right) + \mathcal{R}(\bar{z}_\tau(t) - z_0) \right) \leq C, \quad (7.28a)$$

$$\sup_{t \in (0, T)} |\partial_t \mathcal{E}(t, \underline{u}_\tau(t), z_\tau(t))| \leq C,$$

$$\int_0^T \mathcal{V}(u'_\tau(r)) \, dr \leq C, \quad \int_0^T \mathcal{R}(z'_\tau(r)) \, dr \leq C, \quad (7.28b)$$

$$\int_0^T \mathcal{V}^*(-\bar{\xi}_\tau(r)) \, dr \leq C, \quad (7.28c)$$

and (7.10d) holds.

Furthermore, the interpolants comply with the discrete semistability condition

$$\mathcal{E}(\bar{\tau}_\tau(t), \underline{u}_\tau(t), \bar{z}_\tau(t)) \leq \mathcal{E}(\bar{\tau}_\tau(t), \underline{u}_\tau(t), \hat{z}) + \mathcal{R}(\hat{z} - \bar{z}_\tau(t)) \quad \text{for all } \hat{z} \in \mathbf{Z} \text{ and for all } t \in (0, T], \quad (7.29)$$

and with the discrete energy-dissipation inequality

$$\int_{\bar{\tau}_\tau(s)}^{\bar{\tau}_\tau(t)} \mathcal{V}(u'_\tau(r)) + \mathcal{V}^*(-\bar{\xi}_\tau(r)) \, dr + \int_{\bar{\tau}_\tau(s)}^{\bar{\tau}_\tau(t)} \mathcal{R}(z'_\tau(r)) \, dr + \mathcal{E}(\bar{\tau}_\tau(t), \bar{u}_\tau(t), \bar{z}_\tau(t))$$

$$\leq \mathcal{E}(\bar{\tau}_\tau(s), \bar{u}_\tau(s), \bar{z}_\tau(s)) + \int_{\bar{\tau}_\tau(s)}^{\bar{\tau}_\tau(t)} \partial_t \mathcal{E}(r, \underline{u}_\tau(r), z_\tau(r)) \, dr + \bar{\Lambda} \int_{\bar{\tau}_\tau(s)}^{\bar{\tau}_\tau(t)} \|\bar{u}_\tau(r) - \underline{u}_\tau(r)\|_{\mathbf{V}} \|u'_\tau(r)\|_{\mathbf{V}} \, dr, \quad (7.30)$$

for a suitable  $\bar{\Lambda} > 0$ .

Observe that, for  $\mathcal{V}$  quadratic, estimates (7.28b) and (7.28c) in particular yield

$$\|u'_\tau(r)\|_{L^2(0, T; \mathbf{V})} \leq C, \quad \|\bar{\xi}_\tau(r)\|_{L^2(0, T; \mathbf{V}^*)} \leq C. \quad (7.31)$$

*Proof.* While the discrete semistability (7.29) is again a direct consequence of the minimum problem (4.3a), the discrete energy-dissipation inequality (7.30) follows from a combination of the time-discrete scheme (4.3) with the uniform Fréchet subdifferentiability condition (4.6) on energy sublevels. Hence, it can be derived only after obtaining the energy estimate (7.28a). Therefore, we need to derive an intermediate energy-dissipation inequality, i.e. (7.34) below.

Let us fix  $n \in \{1, \dots, N_\tau\}$ . From the minimum problem (4.3a) we gather

$$\tau \mathcal{R} \left( \frac{z_\tau^n - z_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, u_\tau^{n-1}, z_\tau^n) \leq \mathcal{E}(t_n, u_\tau^{n-1}, z_\tau^{n-1}), \quad (7.32)$$

while (4.3b) yields

$$\tau \mathcal{V} \left( \frac{u_\tau^n - u_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, u_\tau^n, z_\tau^n) \leq \mathcal{E}(t_n, u_\tau^{n-1}, z_\tau^n). \quad (7.33)$$

Adding (7.32) and (7.33) and summing over the index  $n$  leads to the intermediate inequality for all  $0 \leq s \leq t \leq T$

$$\int_{\bar{\tau}_\tau(s)}^{\bar{\tau}_\tau(t)} (\mathcal{V}(u'_\tau(r)) + \mathcal{R}(z'_\tau(r))) \, dr + \mathcal{E}(\bar{\tau}_\tau(t), \bar{u}_\tau(t), \bar{z}_\tau(t)) \leq \mathcal{E}(\bar{\tau}_\tau(s), \bar{u}_\tau(s), \bar{z}_\tau(s)) + \int_{\bar{\tau}_\tau(s)}^{\bar{\tau}_\tau(t)} \partial_t \mathcal{E}(r, \underline{u}_\tau(r), z_\tau(r)) \, dr \quad (7.34)$$

whereby estimates (7.28a) and (7.28b) ensue from the very same arguments as for the proof of Proposition 7.2. Moreover, we recover the stability estimates (7.10d) as a consequence of the dissipation bound in (7.28b).

We now obtain (7.30) by rephrasing the discrete Euler-Lagrange equation (7.26) as

$$\mathcal{V} \left( \frac{u_\tau^n - u_\tau^{n-1}}{\tau} \right) + \mathcal{V}^*(-\xi_\tau^n) = \left\langle -\xi_\tau^n, \frac{u_\tau^n - u_\tau^{n-1}}{\tau} \right\rangle_{\mathbf{V}}. \quad (7.35)$$



On the other hand, from  $\xi_\tau^n \in \partial_u^- \mathcal{E}(t_n, u_\tau^n, z_\tau^n)$  and (4.6) we obtain

$$-\langle \xi_\tau^n, u_\tau^n - u_\tau^{n-1} \rangle_{\mathbf{V}} \leq \mathcal{E}(t_n, u_\tau^{n-1}, z_\tau^n) - \mathcal{E}(t_n, u_\tau^n, z_\tau^n) + \bar{\Lambda} \|u_\tau^n - u_\tau^{n-1}\|_{\mathbf{V}}^2,$$

with  $\bar{\Lambda} = \Lambda_{\bar{C}} > 0$  and the energy sublevel  $\bar{C} := \max\{C, \tilde{C}\}$ , where the constant  $C$  is from (7.28a), which yields that  $(u_\tau^n, z_\tau^n) \in \mathcal{S}_C$ , while  $\tilde{C}$  is from (7.36) below. Indeed, we have

$$\mathcal{E}(t_n, u_\tau^{n-1}, z_\tau^n) \stackrel{(7.32)}{\leq} \mathcal{E}(t_n, u_\tau^{n-1}, z_\tau^{n-1}) = \mathcal{E}(t_{n-1}, u_\tau^{n-1}, z_\tau^{n-1}) + \int_{t_{n-1}}^{t_n} \partial_t \mathcal{E}(s, u_\tau^{n-1}, z_\tau^{n-1}) ds \stackrel{(7.28a)}{\leq} \tilde{C}, \quad (7.36)$$

which gives  $(u_\tau^{n-1}, z_\tau^n) \in \mathcal{S}_{\tilde{C}}$ .

Combining (7.35) and (7.36) we arrive at

$$\tau \mathcal{V} \left( \frac{u_\tau^n - u_\tau^{n-1}}{\tau} \right) + \tau \mathcal{V}^*(-\xi_\tau^n) \leq \mathcal{E}(t_n, u_\tau^{n-1}, z_\tau^n) - \mathcal{E}(t_n, u_\tau^n, z_\tau^n) + \bar{\Lambda} \|u_\tau^n - u_\tau^{n-1}\|_{\mathbf{V}}^2,$$

which we add to (7.32), thus obtaining, with the cancelation of two terms and using that  $\mathcal{E}(t_n, u_\tau^{n-1}, z_\tau^n) = \mathcal{E}(t_{n-1}, u_\tau^{n-1}, z_\tau^n) + \int_{t_{n-1}}^{t_n} \partial_t \mathcal{E}(s, u_\tau^{n-1}, z_\tau^n) ds$ ,

$$\begin{aligned} & \tau \mathcal{V} \left( \frac{u_\tau^n - u_\tau^{n-1}}{\tau} \right) + \tau \mathcal{V}^*(-\xi_\tau^n) + \tau \mathcal{R} \left( \frac{z_\tau^n - z_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, u_\tau^n, z_\tau^n) \\ & \leq \mathcal{E}(t_{n-1}, u_\tau^{n-1}, z_\tau^{n-1}) + \int_{t_{n-1}}^{t_n} \partial_t \mathcal{E}(s, u_\tau^{n-1}, z_\tau^{n-1}) ds + \bar{\Lambda} \|u_\tau^n - u_\tau^{n-1}\|_{\mathbf{V}}^2. \end{aligned} \quad (7.37)$$

Adding up over the index  $n$  leads to (7.30).

Observe that the last term on the right-hand side of (7.30) can be estimated as

$$\bar{\Lambda} \int_{\bar{\tau}_\tau(s)}^{\bar{\tau}_\tau(t)} \|\bar{u}_\tau(r) - \underline{u}_\tau(r)\|_{\mathbf{V}} \|u'_\tau(r)\|_{\mathbf{V}} dr \leq C\tau \int_0^T \|u'_\tau(r)\|_{\mathbf{V}} dr \leq C\tau, \quad (7.38)$$

where for the last estimate we have relied on the previously obtained ‘stability’ estimates (7.10d), as well as the bound (7.28b). Thus, from (7.30) we deduce (7.28c).  $\square$

The proof of the forthcoming Proposition 7.5 relies on the same Young measure argument as Prop. 7.3, hence we omit the details. Let us only mention that, in order to conclude the energy-dissipation inequality (7.40) below we pass to the limit in (7.30), relying on the bound on (7.38) to conclude that the last term on the right-hand side of (7.30) tends to zero. We also use (7.21) from the proof of Theorem 4.13.

**Proposition 7.5.** *Assume Hypotheses 2.2, 2.3, 2.7, and 4.2. Let  $\mathcal{V}$  be quadratic, cf. (4.1).*

*Then, for every null sequence  $(\tau_k)_k$  of time-steps there exist a (not relabeled) subsequence, a pair  $(u, z)$  as in (3.1) and a function  $\mathcal{E} \in \text{BV}([0, T])$ , such that convergences (4.15) hold as  $k \rightarrow \infty$  as well as convergence (4.16) and the energy convergence (7.15). There also exists a Young measure  $\boldsymbol{\mu} = (\mu_t)_{t \in (0, T)}$ , limit of  $(\bar{\xi}_{\tau_k})_k$ , such that*

$$\bar{\xi}_{\tau_k} \rightharpoonup \xi \quad \text{in } L^2(0, T; \mathbf{V}^*) \quad \text{with } \xi(t) = \int_{\mathbf{V}^*} \zeta d\mu_t(\zeta) \quad \text{for a.a. } t \in (0, T). \quad (7.39)$$

*Moreover, the following energy-dissipation inequality holds for all  $0 \leq s \leq t \leq T$*

$$\int_s^t \mathcal{V}(u'(r)) dr + \int_s^t \mathcal{V}^*(-\xi(r)) dr + \text{Var}_{\mathcal{R}}(z; [s, t]) + \mathcal{E}(t) \leq \mathcal{E}(s) + \int_s^t \partial_t \mathcal{E}(t, r, u(r)) dr. \quad (7.40)$$

We are now in the position to carry out the **proof of Theorem 4.9**. The energy-dissipation inequality (3.4) on  $(0, t)$  follows from (7.40), taking into account that  $\mathcal{E}(t) \geq \mathcal{E}(t, u(t), z(t))$  for all  $t \in (0, T]$ , and that  $\mathcal{E}(0) = \mathcal{E}(0, u_0, z_0)$  by (7.15). It remains to prove the subdifferential inclusion (3.2). For this, we distinguish two cases:

**Case (1): Hyp. 4.6 holds:** It follows from  $\mathbf{U} \Subset \mathbf{V}$  that

$$\bar{u}_{\tau_k}, \underline{u}_{\tau_k}, u_{\tau_k} \rightarrow u \quad \text{in } L^\infty(0, T; \mathbf{V}). \quad (7.41)$$

Hence, we have convergence (4.18) and, in view of Lemma 4.7, the energy convergence (4.19) holds. Moreover, Lemma 4.7 guarantees that for almost all  $t \in (0, T)$  the set of the limit points of the sequence  $(\bar{\xi}_{\tau_k}(t))_k$ , hence the support of the measure  $\mu_t$ , is contained in  $\partial_u^- \mathcal{E}(t, u(t), z(t))$ . Then, with the same argument as in the proof of Theorem 4.13, cf. (7.20), we find that  $\xi(t) \in \partial_u^- \mathcal{E}(t, u(t), z(t))$  for a.a.

$t \in (0, T)$ . On the other hand, since  $\mathcal{V}$  is quadratic, passing to the limit as  $k \rightarrow \infty$  in (7.27) we conclude that  $-\xi(t) \in \partial\mathcal{V}(u'(t)) = \{Au'(t)\}$  for almost all  $t \in (0, T)$ , and (3.2) ensues.

**Case (2): Hyp. 4.8 holds:** Passing to the limit in (7.27), exploiting the fact that  $\mathcal{V}$  is quadratic, we get  $-\xi(t) \in \partial\mathcal{V}(u'(t)) = \{Au'(t)\}$  for almost all  $t \in (0, T)$ . We now use Hypothesis 4.8 to show that

$$\xi(t) \in \partial_u^- \mathcal{E}(t, u(t), z(t)) \quad \text{for a.a. } t \in (0, T). \quad (7.42)$$

With this aim, we test (7.27) by  $\bar{u}_{\tau_k}$ , integrate in time, and find for every  $t \in [0, T]$  that

$$\begin{aligned} \limsup_{k \rightarrow \infty} \int_0^{\bar{t}_{\tau_k}(t)} \langle \bar{\xi}_{\tau_k}, \bar{u}_{\tau_k} \rangle_{\mathbf{V}} ds &= - \liminf_{k \rightarrow \infty} \int_0^{\bar{t}_{\tau_k}(t)} \langle Au'_{\tau_k}, \bar{u}_{\tau_k} \rangle_{\mathbf{V}} ds \\ &\leq - \liminf_{k \rightarrow \infty} \frac{1}{2} a(\bar{u}_{\tau_k}(t), \bar{u}_{\tau_k}(t)) + \frac{1}{2} a(u_0, u_0) \\ &\leq -\frac{1}{2} a(u(t), u(t)) + \frac{1}{2} a(u_0, u_0) = - \int_0^t \langle Au', u \rangle_{\mathbf{V}} ds = \int_0^t \langle \xi, u \rangle_{\mathbf{V}} ds. \end{aligned} \quad (7.43)$$

Then, in view of the closedness property (4.13a) we conclude (7.42). If (4.13b) holds in addition, we infer the energy convergence (4.19).

Clearly, under the energy convergence (4.19) it is possible to conclude that  $(u, z)$  fulfills the energy-dissipation inequality (3.4) for all  $t \in (0, T]$  and almost all  $s \in (0, t)$ . For the limit passage in the discrete semistability (7.29) we resort to Hypothesis 2.5.

Observe that, under Hypothesis 4.6 or under the enhanced (4.13b), we also have  $\mathcal{E}(\bar{t}_{\tau_k}(t), \underline{u}_{\tau_k}(t), \bar{z}_{\tau_k}(t)) \rightarrow \mathcal{E}(t, u(t), z(t))$  as  $k \rightarrow \infty$  for almost all  $t \in (0, T)$ . Then, arguing as in (7.36) we infer that

$$\limsup_{k \rightarrow \infty} \mathcal{E}(\bar{t}_{\tau_k}(t), \underline{u}_{\tau_k}(t), \bar{z}_{\tau_k}(t)) \leq \mathcal{E}(t, u(t), z(t)),$$

whence  $\mathcal{E}(\bar{t}_{\tau_k}(t), \underline{u}_{\tau_k}(t), \bar{z}_{\tau_k}(t)) \rightarrow \mathcal{E}(t, u(t), z(t))$  for almost all  $t \in (0, T)$ . Therefore, it is sufficient to exploit the weaker variant of Hypothesis 2.5 in Remark 2.6. This concludes the proof in order to obtain the validity semistability condition (3.3) at almost all  $t \in (0, T)$ .  $\blacksquare$

We now perform the proof of **Lemma 4.7**: to prove that  $\xi \in \partial_u^- \mathcal{E}(t, u, z)$ , we will show that

$$\mathcal{E}(t, v, z) - \mathcal{E}(t, u, z) \geq \langle \xi, v - u \rangle_{\mathbf{V}} - \Lambda \|v - u\|_{\mathbf{V}}^2 \quad \text{for every } v \in \mathbf{D}_u \quad (7.44)$$

With this aim, we use Hypothesis 4.6 and for every fixed  $v \in \mathbf{D}_u$  we consider a recovery sequence  $v_n \rightarrow v$  in  $\mathbf{V}$  such that (4.11) holds. By Hyp. 4.2, there exists  $\Lambda = \Lambda_C > 0$  such that for every  $n \in \mathbb{N}$

$$\mathcal{E}(t_n, v_n, z_n) - \mathcal{E}(t_n, u_n, z_n) \geq \langle \xi_n, v_n - u_n \rangle_{\mathbf{V}} - \Lambda \|v_n - u_n\|_{\mathbf{V}}^2. \quad (7.45)$$

We pass to the limit in (7.45) combining the convergences in (4.12) with (4.11), and conclude (7.44).

To prove the energy convergence, it is sufficient to check  $\limsup_n \mathcal{E}(t_n, u_n, z_n) \leq \mathcal{E}(t, u, z)$ . This can be deduced from choosing  $v = u$  in (4.6), whence

$$\mathcal{E}(t, u, z) \geq \limsup_{n \rightarrow \infty} \mathcal{E}(t_n, u_n, z_n) + \lim_{n \rightarrow \infty} \left( \langle \xi_n, u - u_n \rangle_{\mathbf{V}} - \Lambda \|u - u_n\|_{\mathbf{V}}^2 \right) = \limsup_{n \rightarrow \infty} \mathcal{E}(t_n, u_n, z_n). \quad \blacksquare$$

We close this Section with some comments on the alternating time-incremental minimization schemes for the gradient system  $(\mathbf{V}, \mathbf{Z}, \mathcal{V}, \mathcal{R}, \mathcal{E})$  from Problems 4.1 and 4.11.

**Remark 7.6.** Alternate minimization algorithms can be efficiently used to construct discrete solutions to systems of evolutionary equations whose underlying energy functionals are *separately convex* w.r.t. their variables. Indeed, decoupling the minimization problems typically allows one to exploit these convexity properties, not only for proving the existence of critical points, but also for deriving discrete energy inequalities and a priori estimates.

This strategy was used in the pioneering paper [Rou09], to construct discrete solutions in coupled rate-dependent/rate-independent systems in continuum mechanical models for solids; it has therefore been a natural choice in the present ‘abstract’ context as well. In fact, it has proved to be particularly eligible for the proof of Theorem 4.13. Indeed, for the derivation of the discrete energy-dissipation inequality (7.8) it has been crucial to use the decoupled minimum problem (4.22a) giving rise to the discrete solutions  $(u_\tau^n)$ . As a matter of fact, the argument for Lemma 7.1, which is at the basis of (7.8), does rely on the variational structure of (4.22a).

Let us also mention that a very refined implementation of the alternate minimization algorithms has been recently used in [KN15], in the context of a rate-independent model for fracture and damage driven by a (separately convex) phase-field energy of Ambrosio-Tortorelli type [AT90]. By cleverly modifying the standard ‘one-step’ method, the authors have constructed solutions to the time-continuous rate-independent system akin to the parameterized (Balanced Viscosity) solutions from [MRS13a].

**7.3. Proof of Theorems 5.4 and 5.6.** The following result collects all the a priori estimates on the approximate solutions constructed from the time-discretization scheme in Problem 5.1. In Proposition 7.7 below we will derive the a priori estimates under the uniform subdifferentiability Hypothesis 5.2, whereas in Remark 7.8 we will shortly describe how it is possible to deduce them under the *weaker* uniform subdifferentiability (5.14) in the case of quadratic  $\mathcal{V}$ .

**Proposition 7.7.** *Assume Hypotheses 2.2, 2.3, 2.7, as well as Hypotheses 5.2 and 5.3.*

*Let  $(\bar{u}_\tau, \underline{u}_\tau, u_\tau, \bar{z}_\tau, \underline{z}_\tau, z_\tau)_\tau$  be the approximate solutions constructed from the time-discretization scheme in Problem 5.1, with  $(v_\tau)_\tau$  from (5.5) and  $(\bar{\xi}_\tau)_\tau$  complying with (5.6).*

*Then, the interpolants satisfy the discrete semistability condition (7.29) and the discrete energy-dissipation inequality*

$$\begin{aligned} & \frac{1}{2} \|u'_\tau(t)\|_{\mathbf{W}}^2 + \int_{\bar{t}_\tau(s)}^{\bar{t}_\tau(t)} \mathcal{V}(u'_\tau(r)) + \mathcal{V}^*(-\bar{\xi}_\tau(r) - v'_\tau(r)) \, dr + \int_{\bar{t}_\tau(s)}^{\bar{t}_\tau(t)} \mathcal{R}(z'_\tau(r)) \, dr + \mathcal{E}(\bar{t}_\tau(t), \bar{u}_\tau(t), \bar{z}_\tau(t)) \\ & \leq \frac{1}{2} \|u'_\tau(s)\|_{\mathbf{W}}^2 + \mathcal{E}(\bar{t}_\tau(s), \bar{u}_\tau(s), \bar{z}_\tau(s)) + \int_{\bar{t}_\tau(s)}^{\bar{t}_\tau(t)} \partial_t \mathcal{E}(r, \underline{u}_\tau(r), \underline{z}_\tau(r)) \, dr + \Lambda \tau^\nu \int_{\bar{t}_\tau(s)}^{\bar{t}_\tau(t)} \|u'_\tau(r)\|_{\mathbf{W}} \mathcal{V}(u'_\tau(r))^\nu \, dr, \end{aligned} \quad (7.46)$$

*with  $\Lambda > 0$  as in (5.7). Furthermore, there exists  $C > 0$  such that for every  $0 < \tau < \tau_0$  there hold the stability property (7.10d), the energy-dissipation estimates (7.28a)–(7.28b), and*

$$\int_0^T \mathcal{V}^*(-\bar{\xi}_\tau(r) - v'_\tau(r)) \, dr \leq C, \quad (7.47a)$$

$$\|u'_\tau\|_{L^\infty(0,T;\mathbf{W})} \leq C \quad (7.47b)$$

*Hence,*

$$\|\bar{\xi}_\tau\|_{L^\infty(0,T;\mathbf{V}^*)} \leq C, \quad (7.47c)$$

$$(v'_\tau)_\tau \text{ is equi-integrable in } L^1(0,T;\mathbf{V}^*), \quad (7.47d)$$

*and, thus,*

$$\sup_{t \in [0,T]} \|v_\tau(t) - u'_\tau(t)\|_{\mathbf{V}^*} \rightarrow 0 \quad \text{as } \tau \downarrow 0. \quad (7.47e)$$

*Proof.* Preliminarily, we observe that the Euler-Lagrange equation (5.4) follows from the same argument as in (7.25) in the case of quadratic  $\mathcal{V}$ , taking into account the sum rule

$$\begin{aligned} & \partial^- \left( \frac{1}{\tau^2} \|\cdot - 2u_\tau^{n-1} + u_\tau^{n-2}\|_{\mathbf{W}}^2 + \tau \mathcal{V} \left( \frac{\cdot - u_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, \cdot, z_\tau^n) \right) (u_\tau^n) \\ & = \frac{u_\tau^n - 2u_\tau^{n-1} + u_\tau^{n-2}}{\tau^2} + \partial \mathcal{V} \left( \frac{u_\tau^n - u_\tau^{n-1}}{\tau} \right) + \partial_u^- \mathcal{E}(t_n, u_\tau^n, z_\tau^n) \end{aligned}$$

due to the Fréchet differentiability of  $\mathcal{V}$  and of  $\|\cdot\|_{\mathbf{W}}^2$ .

For  $\mathcal{V}$  general, we apply [MRS13b, Prop. 4.2] to  $\mathcal{V}$  and to  $\mathcal{E}_n := \mathcal{E}(t_n, \cdot, z_\tau^n) + \frac{1}{\tau^2} \|\cdot - 2u_\tau^{n-1} + u_\tau^{n-2}\|_{\mathbf{W}}^2$ , and then observe that the Fréchet subdifferential of  $\mathcal{E}_n$  at  $u_\tau^n$  decomposes into the sum of the Fréchet subdifferential of  $\mathcal{E}$  and the singleton  $\{\frac{1}{\tau^2}(u_\tau^n - 2u_\tau^{n-1} + u_\tau^{n-2})\}$ .

We now start with the proof of (7.46). We rephrase the Euler-Lagrange equation (5.4) as (cf. (7.35))

$$\begin{aligned} & \mathcal{V}\left(\frac{u_\tau^n - u_\tau^{n-1}}{\tau}\right) + \mathcal{V}^*\left(-\xi_\tau^n - \frac{u_\tau^n - 2u_\tau^{n-1} + u_\tau^{n-2}}{\tau^2}\right) \\ &= \left\langle -\xi_\tau^n - \frac{u_\tau^n - 2u_\tau^{n-1} + u_\tau^{n-2}}{\tau^2}, \frac{u_\tau^n - u_\tau^{n-1}}{\tau} \right\rangle_{\mathbf{V}} \\ &\leq -\left\langle \xi_\tau^n, \frac{u_\tau^n - u_\tau^{n-1}}{\tau} \right\rangle_{\mathbf{V}} + \frac{1}{2\tau} \frac{\|u_\tau^{n-1} - u_\tau^{n-2}\|_{\mathbf{W}}^2}{\tau^2} - \frac{1}{2\tau} \frac{\|u_\tau^n - u_\tau^{n-1}\|_{\mathbf{W}}^2}{\tau^2}. \end{aligned} \quad (7.48)$$

On the other hand, from  $\xi_\tau^n \in \partial_u^- \mathcal{E}(t_n, u_\tau^n, z_\tau^n)$  and the Fréchet subdifferentiability condition (5.7) we obtain

$$\langle \xi_\tau^n, u_\tau^n - u_\tau^{n-1} \rangle_{\mathbf{V}} \leq \mathcal{E}(t_n, u_\tau^n, z_\tau^n) - \mathcal{E}(t_n, u_\tau^{n-1}, z_\tau^n) + \Lambda \|u_\tau^n - u_\tau^{n-1}\|_{\mathbf{W}} \mathcal{V}(u_\tau^n - u_\tau^{n-1})^\nu. \quad (7.49)$$

Combining (7.48) and (7.49) with estimate (7.32) which derives from the minimum problem (5.3a), we get

$$\begin{aligned} & \frac{1}{2} \frac{\|u_\tau^n - u_\tau^{n-1}\|_{\mathbf{W}}^2}{\tau^2} + \tau \mathcal{V}\left(\frac{u_\tau^n - u_\tau^{n-1}}{\tau}\right) + \tau \mathcal{V}^*\left(-\xi_\tau^n - \frac{u_\tau^n - 2u_\tau^{n-1} + u_\tau^{n-2}}{\tau^2}\right) + \mathcal{E}(t_n, u_\tau^n, z_\tau^n) + \mathcal{R}(z_\tau^n - z_\tau^{n-1}) \\ &\leq \frac{1}{2} \frac{\|u_\tau^{n-1} - u_\tau^{n-2}\|_{\mathbf{W}}^2}{\tau^2} + \mathcal{E}(t_{n-1}, u_\tau^{n-1}, z_\tau^{n-1}) + \int_{t_{n-1}}^{t_n} \partial_t \mathcal{E}(s, u_\tau^{n-1}, z_\tau^{n-1}) ds + \Lambda \|u_\tau^n - u_\tau^{n-1}\|_{\mathbf{W}} \mathcal{V}(u_\tau^n - u_\tau^{n-1})^\nu. \end{aligned} \quad (7.50)$$

Next, we use that

$$\begin{aligned} \Lambda \tau \left\| \frac{u_\tau^n - u_\tau^{n-1}}{\tau} \right\|_{\mathbf{W}} \mathcal{V}(u_\tau^n - u_\tau^{n-1})^\nu &\stackrel{(1)}{\leq} \Lambda \tau \left\| \frac{u_\tau^n - u_\tau^{n-1}}{\tau} \right\|_{\mathbf{W}} \tau^\nu \mathcal{V}\left(\frac{u_\tau^n - u_\tau^{n-1}}{\tau}\right)^\nu \\ &= \Lambda \tau^\nu \int_{\underline{t}_\tau(t)}^{\bar{t}_\tau(t)} \|u'_\tau(r)\|_{\mathbf{W}} \mathcal{V}(u'_\tau(r))^\nu dr, \end{aligned} \quad (7.51)$$

where (1) follows from the fact that  $\mathcal{V}(u_\tau^n - u_\tau^{n-1}) \leq \tau \mathcal{V}\left(\frac{u_\tau^n - u_\tau^{n-1}}{\tau}\right)$  by the convexity of  $\mathcal{V}$  and the fact that  $\mathcal{V}(0) = 0$ . Combining (7.50) and (7.51) and adding up over the index  $n$  leads to (7.46).

We are now in the position to deduce estimates (7.28a)–(7.28b) and (7.47). Indeed, we estimate the last term on the right-hand side of (7.46) by

$$\begin{aligned} & \Lambda \tau^\nu \int_{\underline{t}_\tau(t)}^{\bar{t}_\tau(t)} \|u'_\tau(r)\|_{\mathbf{W}} \mathcal{V}(u'_\tau(r))^\nu dr \\ &\stackrel{(2)}{\leq} \frac{1}{2} \int_{\underline{t}_\tau(t)}^{\bar{t}_\tau(t)} \mathcal{V}(u'_\tau(r)) dr + C \tau^{\nu/(1-\nu)} \int_{\underline{t}_\tau(t)}^{\bar{t}_\tau(t)} \|u'_\tau(r)\|_{\mathbf{W}}^{1/(1-\nu)} dr, \\ &\stackrel{(3)}{\leq} \frac{1}{2} \int_{\underline{t}_\tau(t)}^{\bar{t}_\tau(t)} \mathcal{V}(u'_\tau(r)) dr + C \tau^{\nu/(1-\nu)} \int_{\underline{t}_\tau(t)}^{\bar{t}_\tau(t)} (\|u'_\tau(r)\|_{\mathbf{W}}^2 + 1) dr, \end{aligned}$$

while (2) is due to Young's inequality and for (3) we have used that  $\nu \leq \frac{1}{2}$ . Therefore, from (7.46), we gather

$$\begin{aligned} & \frac{1}{2} \|u'_\tau(t)\|_{\mathbf{W}}^2 + \int_0^{\bar{t}_\tau(t)} \frac{1}{2} \mathcal{V}(u'_\tau(r)) + \mathcal{V}^*(-\bar{\xi}_\tau(r) - v'_\tau(r)) dr + \int_0^{\bar{t}_\tau(t)} \mathcal{R}(z'_\tau(r)) dr + \mathcal{E}(\bar{t}_\tau(t), \bar{u}_\tau(t), \bar{z}_\tau(t)) \\ &\leq \frac{1}{2} \|v_0\|_{\mathbf{W}}^2 + \mathcal{E}(0, u_0, z_0) + \int_0^{\bar{t}_\tau(t)} \partial_t \mathcal{E}(r, \underline{u}_\tau(r), \underline{z}_\tau(r)) dr + C \tau^{\nu/(1-\nu)} \int_0^{\bar{t}_\tau(t)} \|u'_\tau(r)\|_{\mathbf{W}}^2 dr + C \tau^{\nu/(1-\nu)}. \end{aligned} \quad (7.52)$$

and we conclude by estimating the power term on the right-hand side via (2.6c), as in the proof of Proposition 7.2, and by applying the Gronwall Lemma. Therefore, estimates (7.28a)–(7.28b) and (7.47a)–(7.47b) ensue.

Finally, (7.47c) is a consequence of the bound for the energy (7.28a) combined with Hypothesis 5.3. Then, (7.47d) follows from (7.47a), taking into account that  $\mathcal{V}^*$  has superlinear growth. The stability estimate (7.47e) is an immediate consequence of (7.47d), taking into account that

$$\|v_\tau(t) - u'_\tau(t)\|_{\mathbf{V}^*} \leq \int_{\underline{t}_\tau(t)}^{\bar{t}_\tau(t)} \|v'_\tau(s)\|_{\mathbf{V}^*} ds \quad \text{for every } t \in [0, T].$$

□

**Remark 7.8.** First of all, let us mention that for quadratic  $\mathcal{V}$  (cf. (4.1)), and under the Fréchet subdifferentiability condition (5.14), the time-incremental minimum problem for  $u$  in (5.3b) has a unique solution  $u_\tau^n$ . To see this, we consider two solutions  $u^1$  and  $u^2$  to (5.3b). We subtract the Euler-Lagrange equation for  $u^2$  from the one for  $u^1$ , and test the relation thus obtained by  $u^1 - u^2$ . We thus conclude

$$\frac{1}{\tau^2} \|u^1 - u^2\|_{\mathbf{W}}^2 + \frac{1}{\tau} a(u^1 - u^2, u^1 - u^2) + \langle \xi^1 - \xi^2, u^1 - u^2 \rangle_{\mathbf{V}} = 0,$$

with  $\xi^i \in \partial_u^- \mathcal{E}(t_n, u^i, z_\tau^n)$  fulfilling the Euler-Lagrange equation for  $i = 1, 2$ . Now, from (5.14) we gather that

$$\langle \xi^1 - \xi^2, u^1 - u^2 \rangle_{\mathbf{V}} \geq -\Lambda \|u^1 - u^2\|_{\mathbf{V}}^2.$$

Combining this with the above relation, using that  $a(\cdot, \cdot)$  is coercive with respect to the norm of  $\mathbf{V}$ , and choosing  $\tau > 0$  sufficiently small, we ultimately deduce that  $\|u^1 - u^2\|_{\mathbf{W}}^2 + c \|u^1 - u^2\|_{\mathbf{V}}^2 \leq 0$ , whence  $u^1 = u^2$ .

Moreover, observe that the very same arguments as in the proof of Prop. 7.7 yield that, in addition to the discrete semistability (7.29), the approximate solutions fulfill the discrete energy-dissipation inequality

$$\begin{aligned} & \frac{1}{2} \|u'_\tau(t)\|_{\mathbf{W}}^2 + \int_{\bar{\tau}_\tau(s)}^{\bar{\tau}_\tau(t)} \mathcal{V}(u'_\tau(r)) + \mathcal{V}^*(-\bar{\xi}_\tau(r) - v'_\tau(r)) dr + \int_{\bar{\tau}_\tau(s)}^{\bar{\tau}_\tau(t)} \mathcal{R}(z'_\tau(r)) dr + \mathcal{E}(\bar{\tau}_\tau(t), \bar{u}_\tau(t), \bar{z}_\tau(t)) \\ & \leq \frac{1}{2} \|u'_\tau(s)\|_{\mathbf{W}}^2 + \mathcal{E}(\bar{\tau}_\tau(s), \bar{u}_\tau(s), \bar{z}_\tau(s)) + \int_{\bar{\tau}_\tau(s)}^{\bar{\tau}_\tau(t)} \partial_t \mathcal{E}(r, \underline{u}_\tau(r), \underline{z}_\tau(r)) dr + \Lambda \tau^{1/2} \int_{\bar{\tau}_\tau(s)}^{\bar{\tau}_\tau(t)} \|u'_\tau(r)\|_{\mathbf{V}}^2 dr, \end{aligned} \quad (7.53)$$

whereby estimates (7.47) ensue. In particular, since  $\mathcal{V}$  is quadratic, due to the quadratic growth of  $\mathcal{V}^*$  we have

$$\|v'_\tau\|_{L^2(0, T; \mathbf{V}^*)} \leq C. \quad (7.54)$$

We are now in the position to carry out the **proof of Theorem 5.4**. Let  $(\tau_k)_k$  be a null sequence. With standard weak and strong compactness arguments (cf., e.g., [Sim87]), taking into account estimates (7.28) and Hyp. 2.7, we find a quadruple  $(u, z, \xi, v)$  with  $(u, z)$  as in (3.1),  $\xi \in L^\infty(0, T; \mathbf{V}^*)$ , and  $v \in L^1(0, T; \mathbf{V}^*)$  such that convergences (4.15) hold, and, in addition,

$$u_{\tau_k} \xrightarrow{*} u \quad \text{in } W^{1, \infty}(0, T; \mathbf{W}), \quad (7.55)$$

$$u_{\tau_k} \rightarrow u \quad \text{in } C^0([0, T]; \mathbf{W}), \quad (7.56)$$

$$\bar{u}_{\tau_k}(t) \rightarrow u(t) \quad \text{in } \mathbf{W} \quad \text{for every } t \in [0, T], \quad (7.57)$$

$$\bar{\xi}_{\tau_k} \xrightarrow{*} \xi \quad \text{in } L^\infty(0, T; \mathbf{V}^*), \quad (7.58)$$

$$v_{\tau_k} \xrightarrow{*} v \quad \text{in } L^\infty(0, T; \mathbf{W}) \cap W^{1, 1}(0, T; \mathbf{V}^*), \quad (7.59)$$

$$v_{\tau_k}(t) \rightharpoonup v(t) \quad \text{in } \mathbf{W} \quad \text{for every } t \in [0, T], \quad (7.60)$$

where (7.59) follows from combining the uniform integrability (7.47d) with the  $L^\infty(0, T; \mathbf{W})$ -bound for  $u'_{\tau_k}$ , inherited by  $v_{\tau_k}$ , whereas (7.60) ensues from the fact that the latter sequence is bounded in  $L^\infty(0, T; \mathbf{W}) \cap W^{1, 1}(0, T; \mathbf{V}^*)$  via an infinite-dimensional version of the Helly theorem, cf. e.g. [MT04, Thm. 6.1]. With (7.47e) we easily find that  $v = u'$ , hence  $v' = u''$ , which gives  $u \in W^{2, 1}(0, T; \mathbf{V}^*)$ . Also taking into account (7.60), we infer the pointwise convergence for  $u'_{\tau_k}(t)$  in the second of convergences (5.9), which are therefore proved.

Next, we prove that

$$\xi(t) \in \partial_u^- \mathcal{E}(t, u(t), z(t)) \quad \text{for a.a. } t \in (0, T). \quad (7.61)$$

We distinguish two cases:

- (1) Under the weak closedness condition from Hypothesis 4.12, we employ the very same Young measure argument as in the proof of Theorem 4.13, cf. (7.20), and conclude (7.61).
- (2) In the case  $\mathbf{U} \subseteq \mathbf{V}$  with Hypothesis 4.6, to prove (7.61) we again resort to the Young measure argument from the proof of Theorem 4.9: In order to conclude that the set of the weak limit points of  $(\xi_{\tau_k}(t))_k$  w.r.t. the topology of  $\mathbf{V}^*$  is contained in  $\partial_u^- \mathcal{E}(t, u(t), z(t))$ , we use the closedness property (4.12) from Lemma 4.7. A close perusal of its proof shows that it extends to the case in which the *uniform Fréchet subdifferentiability* holds in the form of Hypothesis 4.3. In fact, the remainder term in (7.45) now reads  $-\Lambda \mathcal{V}(v_n - u_n)^\nu \|v_n - u_n\|_{\mathbf{W}}$ , and it goes to zero as  $n \rightarrow \infty$  since  $\mathbf{V} \subset \mathbf{W}$  continuously, and  $\mathcal{V}$  is continuous on  $\mathbf{V}$ . Likewise, it also extends to the case when the Fréchet subdifferentiability Hypothesis 4.2 is replaced by its weaker variant (5.14).

We now address the limit passage in the discrete energy-dissipation inequality (7.53): It follows from (7.58)–(7.59) that

$$\liminf_{k \rightarrow \infty} \int_{\bar{\tau}_k(s)}^{\bar{\tau}_k(t)} \mathcal{V}^*(-\bar{\xi}_{\tau_k}(r) - v'_{\tau_k}(r)) dr \geq \int_s^t \mathcal{V}^*(-\xi(r) - u''(r)) dr$$

whereas for the remainder term on the right-hand side of (7.53) we have

$$\Lambda \tau^{1/2} \int_{\bar{\tau}_k(s)}^{\bar{\tau}_k(t)} \|u'_{\tau_k}(r)\|_{\mathbf{W}} \mathcal{V}(u'_{\tau_k}(r))^{1/2} dr \leq \Lambda \tau^{1/2} \left( \int_{\bar{\tau}_k(s)}^{\bar{\tau}_k(t)} \|u'_{\tau_k}(r)\|_{\mathbf{W}}^2 dr \right) \left( \int_{\bar{\tau}_k(s)}^{\bar{\tau}_k(t)} \mathcal{V}(u'_{\tau_k}(r)) dr \right) \rightarrow 0 \text{ as } k \rightarrow \infty$$

in view of estimates (7.28b) and (7.47b). Therefore, also exploiting the previously obtained convergences, we conclude the energy-dissipation inequality (3.4) on every interval  $(0, t)$ .

The proof of the energy convergence (4.19) under (4.24b), or under Hyp. 4.6 (and of (4.18) in the case  $\mathbf{U} \Subset \mathbf{V}$ ) follows the very same lines as the arguments developed for Thms. 4.9 & 4.13.

Arguing as in the proof of Theorem 4.13, we use Hypothesis 2.5 (possibly in its weaker form from Remark 2.6, under the enhanced closedness condition (4.24b), or under Hyp. 4.6, which gives the closedness (4.12)), to pass to the limit in the discrete semistability (7.29) and conclude that  $(u, z)$  fulfill the semistability (3.3).

Thus, we conclude that  $(u, z)$  is a weak semistable energetic solution to the system  $(\mathbf{V}, \mathbf{W}, \mathbf{Z}, \mathcal{V}, \mathcal{K}, \mathcal{R}, \mathcal{E})$ . ■

**Remark 7.9.** Even under the conditions guaranteeing the energy convergence (4.19), the proof of the energy-dissipation inequality in its enhanced closedness form remains open. Indeed, due to the *weak* character of the (albeit pointwise) convergence for  $(u'_{\tau_k}(t))_k$  (cf. the second of (5.9)), it is not clear how to pass to the limit in the discrete energy-dissipation inequality (7.46) on a generic interval  $(s, t)$ .

Finally, we present the proof of **Theorem 5.6**, just dwelling on its differences from the proof of Thm. 5.4. Let  $(\tau_k)_k$  be a null sequence. The very same compactness arguments as in the proof of Thm. 5.4 yield that there exists a quadruple  $(u, z, \xi, v)$  with  $(u, z)$  as in (3.1), such that convergences (4.15) and (7.55)–(7.58) hold. Moreover,

$$v_{\tau_k} \rightharpoonup v = u' \quad \text{in } H^1(0, T; \mathbf{V}^*), \quad (7.62)$$

and, in view of the estimates for  $(u'_{\tau_k})_k$  in  $L^2(0, T; \mathbf{V}) \cap L^\infty(0, T; \mathbf{V}^*) \cap H^1(0, T; \mathbf{V}^*)$  and of the compact embedding  $\mathbf{V} \Subset \mathbf{W}$ , we may also suppose, up to a further extraction, that

$$\begin{aligned} u'_{\tau_k} &\rightarrow u' \quad \text{in } L^p(0, T; \mathbf{W}) \quad \text{for all } 1 \leq p < \infty, \quad \text{whence} \\ u'_{\tau_k}(t) &\rightarrow u'(t) \quad \text{in } \mathbf{W} \quad \text{for almost all } t \in (0, T). \end{aligned} \quad (7.63)$$

Exploiting convergences (7.55)–(7.58) as well as (7.62), and the fact that  $\partial \mathcal{V}$  is a linear and bounded operator  $A$ , we first pass to the limit in (5.6) and conclude that  $(u, z)$  fulfill

$$u''(t) + Au'(t) + \xi(t) = 0 \quad \text{for a.a. } t \in (0, T), \quad (7.64)$$

with  $\xi \in L^\infty(0, T; \mathbf{V}^*)$  from (7.58). In order to show that  $\xi(t) \in \partial_u^- \mathcal{E}(t, u(t), z(t))$  for almost all  $t \in (0, T)$ , we resort to Hypothesis 4.8. Hence, we test the discrete momentum equation (5.6) by  $\bar{u}_{\tau_k}$  and integrate in time, thus obtaining

$$\int_0^{\bar{\tau}_k(t)} \langle \bar{\xi}_{\tau_k}(s), \bar{u}_{\tau_k}(s) \rangle_{\mathbf{V}} ds = - \int_0^{\bar{\tau}_k(t)} (v'_{\tau_k}(s), \bar{u}_{\tau_k}(s))_{\mathbf{W}} ds - \int_0^{\bar{\tau}_k(t)} \langle Au'_{\tau_k}(s), \bar{u}_{\tau_k}(s) \rangle ds \doteq I_1 + I_2.$$

Now, the discrete integration-by-part formula yields

$$\begin{aligned} I_1 &= \int_0^{\bar{\tau}_k(t)} (u'_{\tau_k}(s), u'_{\tau_k}(s - \tau_k))_{\mathbf{W}} ds - (\bar{u}_{\tau_k}(t), u'_{\tau_k}(t))_{\mathbf{W}} + (u_0, \dot{u}_\tau^0)_{\mathbf{W}} \\ &\xrightarrow{k \rightarrow \infty} \int_0^t (u'(s), u'(s))_{\mathbf{W}} ds - (u'(t), u(t))_{\mathbf{W}} + (u_0, v_0)_{\mathbf{W}} = - \int_0^t (u''(s), u(s))_{\mathbf{W}} ds, \end{aligned}$$

with  $\dot{u}_\tau^0 := \frac{u_0 - u_\tau^{-1}}{\tau}$  and  $u_\tau^{-1}$  from (5.2), where the limit passage ensues from convergences (7.63). With the very same calculations as for (7.43), we find that  $\limsup_{k \rightarrow \infty} I_2 \leq - \int_0^t \langle Au', u \rangle_{\mathbf{V}} ds$ . Taking into account (7.64), we then conclude that  $\limsup_{k \rightarrow \infty} \int_0^T \langle \bar{\xi}_{\tau_k}(t), \bar{u}_{\tau_k}(t) \rangle_{\mathbf{V}} dt \leq \int_0^T \langle \xi(t), u(t) \rangle_{\mathbf{V}} dt$ , whence the identification of  $\xi(t)$  as an element in  $\partial_u^- \mathcal{E}(t, u(t), z(t))$  by Hypothesis 4.8.

The enhanced energy inequality (5.12) under the stronger requirement (4.13b) from Hyp. 4.8 can be proved by passing to the limit in (7.53), taking into account the gained pointwise energy convergence, as well as the second of convergences (7.63). This concludes the proof.  $\blacksquare$

## APPENDIX A. YOUNG MEASURE TOOLS

For the reader's convenience, we collect here the definition of parameterized (or Young) measure with values in infinite-dimensional spaces, and the Young measure compactness result from [MRS13b] (cf. also [RS06, Ste08]), which extends to the frame of the weak topology classical results within Young measure theory (see e.g. [Bal84, Thm. 1], [Val90, Thm. 16]).

**Notation.** Given an interval  $I \subset \mathbb{R}$ , we denote by  $\mathcal{L}_I$  the  $\sigma$ -algebra of the Lebesgue measurable subsets of  $I$  and, given a reflexive Banach space  $B$ , by  $\mathcal{B}(B)$  its Borel  $\sigma$ -algebra. We use the symbol  $\otimes$  for product  $\sigma$ -algebrae. We recall that a  $\mathcal{L}_I \otimes \mathcal{B}(B)$ -measurable function  $h : I \times B \rightarrow (-\infty, +\infty]$  is a *normal integrand* if for a.a.  $t \in (0, T)$  the map  $x \mapsto h_t(x) = h(t, x)$  is lower semicontinuous on  $B$ .

We consider the space  $B$  endowed with the *weak* topology, and say that a  $\mathcal{L}_{(0,T)} \otimes \mathcal{B}(B)$ -measurable functional  $\mathcal{H} : (0, T) \times B \rightarrow (-\infty, +\infty]$  is a *weakly-normal integrand* if for a.a.  $t \in (0, T)$  the map

$$w \mapsto \mathcal{H}(t, w) \text{ is sequentially lower semicontinuous on } B \text{ w.r.t. the weak topology.} \quad (\text{A.1})$$

We denote by  $\mathcal{M}(0, T; \mathcal{V})$  the set of all  $\mathcal{L}_{(0,T)}$ -measurable functions  $y : (0, T) \rightarrow B$ . A sequence  $(w_n) \subset \mathcal{M}(0, T; B)$  is said to be *weakly-tight* if there exists a weakly-normal integrand  $\mathcal{H} : (0, T) \times B \rightarrow (-\infty, +\infty]$  such that the map

$$w \mapsto \mathcal{H}(t, w) \text{ has compact sublevels w.r.t. the weak topology of } B, \text{ and}$$

$$\sup_n \int_0^T \mathcal{H}(t, w_n(t)) dt < \infty.$$

**Definition A.1 ((Time-dependent) Young measures).** A Young measure in the space  $B$  is a family  $\mu := \{\mu_t\}_{t \in (0, T)}$  of Borel probability measures on  $B$  such that the map on  $(0, T)$

$$t \mapsto \mu_t(A) \text{ is } \mathcal{L}_{(0,T)}\text{-measurable for all } A \in \mathcal{B}(B). \quad (\text{A.2})$$

We denote by  $\mathcal{Y}(0, T; B)$  the set of all Young measures in  $B$ .

**Theorem A.2.** [MRS13b, Theorems A.2, A.3] Let  $\{\mathcal{H}_n\}$ ,  $\mathcal{H} : (0, T) \times B \rightarrow (-\infty, +\infty]$  be weakly-normal integrands such that for all  $w \in B$  and for a.a.  $t \in (0, T)$

$$\mathcal{H}(t, w) \leq \inf \left\{ \liminf_{n \rightarrow \infty} \mathcal{H}_n(t, w_n) : w_n \rightharpoonup w \text{ in } B \right\}. \quad (\text{A.3})$$

Let  $(w_n) \subset \mathcal{M}(0, T; B)$  be a weakly-tight sequence. Then, there exist a subsequence  $(w_{n_k})$  and a Young measure  $\mu = \{\mu_t\}_{t \in (0, T)}$  such that for a.a.  $t \in (0, T)$

$$\mu_t \text{ is concentrated on the set } L(t) := \bigcap_{p=1}^{\infty} \overline{\{w_{n_k}(t) : k \geq p\}}^w \quad (\text{A.4})$$

of the limit points of the sequence  $(w_{n_k}(t))$  with respect to the weak topology of  $B$  and, if the sequence  $t \mapsto \mathcal{H}_{n_k}^-(t, w_{n_k}(t))$  is uniformly integrable, there holds

$$\liminf_{k \rightarrow \infty} \int_0^T \mathcal{H}_{n_k}(t, w_{n_k}(t)) dt \geq \int_0^T \int_{\mathcal{V}} \mathcal{H}(t, w) d\mu_t(w) dt. \quad (\text{A.5})$$

If in addition  $(w_n) \subset L^p(0, T; \mathcal{V})$  is bounded and, in the case  $p = 1$ , also equi-integrable, then, up to extracting a further subsequence from  $(w_{n_k})$  we have that, setting

$$w(t) := \int_B w d\mu_t(w) \quad \text{for a.a. } t \in (0, T),$$

there holds

$$w_{n_k} \rightharpoonup w \quad \text{in } L^p(0, T; B), \quad (\text{A.6})$$

with  $\rightharpoonup$  replaced by  $\overset{*}{\rightharpoonup}$  if  $p = \infty$ .

**Acknowledgment.** R.R. has been partially supported by a MIUR-PRIN'10-11 grant for the project “Calculus of Variations”. M.T. has been partially supported by the Deutsche Forschungsgemeinschaft in the Priority Program 1748 “Reliable simulation techniques in solid mechanics. Development of non-standard discretization methods, mechanical and mathematical analysis” within the project “Finite element approximation of functions of bounded variation and application to models of damage, fracture, and plasticity”. R.R. and M.T. are also grateful for the support from the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). This research was carried out during several visits of R.R. at the Weierstrass Institute, supported by the European Research Council through the ERC Advanced Grant *Analysis of Multiscale Systems Driven by Functionals (267802)*, and of M.T. at the University of Brescia; the kind hospitality at both institutions is gratefully acknowledged.

The authors are grateful to the two anonymous referees of this paper for their careful reading and for plenty of valuable suggestions.

## REFERENCES

- [AFP05] L. Ambrosio, N. Fusco, and D. Pallara. *Functions of Bounded Variation and Free Discontinuity Problems*. Oxford University Press, 2005.
- [AGS08] L. Ambrosio, N. Gigli, and G. Savaré. *Gradient Flows in Metric Spaces and in the Space of Probability Measures*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, second edition, 2008.
- [Amb95] L. Ambrosio. Minimizing movements. *Rend. Accad. Naz. Sci. XL Mem. Mat. Appl. (5)*, 19:191–246, 1995.
- [AMV14] R. Alessi, J.-J. Marigo, and S. Vidoli. Gradient damage models coupled with plasticity and nucleation of cohesive cracks. *Arch. Ration. Mech. Anal.*, 214(2):575–615, 2014.
- [AT90] L. Ambrosio and V.M. Tortorelli. Approximation of functionals depending on jumps by elliptic functionals via  $\Gamma$ -convergence. *Comm. Pure Appl. Math.*, 43(8):999–1036, 1990.
- [Att84] H. Attouch. *Variational Convergence for Functions and Operators*. Applicable Mathematics Series. Pitman (Advanced Publishing Program), Boston, MA, 1984.
- [Bal84] E. J. Balder. A general approach to lower semicontinuity and lower closure in optimal control theory. *SIAM J. Control Optim.*, 22(4):570–598, 1984.
- [CL16] V. Crismale and G. Lazzaroni. Viscous approximation of quasistatic evolutions for a coupled elastoplastic-damage model. *Calc. Var. Partial Differential Equations*, 55, 2016. doi: 10.1007/s00526-015-0947-6.
- [Cri16] V. Crismale. Globally stable quasistatic evolution for a coupled elastoplastic-damage model. *ESAIM Control Optim. Calc. Var.*, 22:883–912, 2016.
- [DGMT80] E. De Giorgi, A. Marino, and M. Tosques. Problems of evolution in metric spaces and maximal decreasing curve. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8)*, 68(3):180–187, 1980.
- [DMFT05] G. Dal Maso, G. Francfort, and R. Toader. Quasistatic crack growth in nonlinear elasticity. *Arch. Rat. Mech. Anal.*, 176:165–225, 2005.
- [DPZ13] M.H. Duong, M.A. Peletier, and J. Zimmer. GENERIC formalism of a Vlasov-Fokker-Planck equation and connection to large-deviation principles. *Nonlinearity*, 26(11):2951–2971, 2013.
- [DU77] J. Diestel and J. J. Uhl, Jr. *Vector Measures*. American Mathematical Society, Providence, R.I., 1977. With a foreword by B. J. Pettis, Mathematical Surveys, No. 15.
- [EŠ11] E. Emmrich and D. Šiška. Full discretisation of second-order nonlinear evolution equations: strong convergence and applications. *Comput. Methods Appl. Math.*, 11(4):441–459, 2011.
- [EŠ13] E. Emmrich and D. Šiška. Evolution equations of second order with nonconvex potential and linear damping: existence via convergence of a full discretization. *J. Differential Equations*, 255(10):3719–3746, 2013.
- [ET74] I. Ekeland and R. Temam. *Analyse Convexe et Problèmes Variationnels*. Dunod; Gauthier-Villars, Paris-Brussels-Montreal, Que., 1974. Collection Études Mathématiques.
- [ET11] E. Emmrich and M. Thälhammer. Doubly nonlinear evolution equations of second order: existence and fully discrete approximation. *J. Differential Equations*, 251(1):82–118, 2011.
- [FL07] I. Fonseca and G. Leoni. *Modern Methods in the Calculus of Variations:  $L^p$  Spaces*. Springer, New York, 2007.
- [Fré88] M. Frémond. Contact with adhesion. In J.J. Moreau, P.D. Panagiotopoulos, and G. Strang, editors, *Topics in Non-smooth Mechanics*, pages 157–186. Birkhäuser, 1988.
- [Fré02a] M. Frémond. *Non-Smooth Thermomechanics*. Springer-Verlag, Berlin, 2002.
- [Fré02b] M. Frémond. *Non-Smooth Thermomechanics*. Springer-Verlag Berlin Heidelberg, 2002.
- [Giu03] E. Giusti. *Direct Methods in the Calculus of Variations*. World Scientific Publishing Co., Inc., River Edge, NJ, 2003.
- [HK11] C. Heinemann and C. Kraus. Existence of weak solutions for Cahn-Hilliard systems coupled with elasticity and damage. *Adv. Math. Sci. Appl.*, 21(2):321–359, 2011.
- [HK13] C. Heinemann and C. Kraus. Existence results for diffuse interface models describing phase separation and damage. *European J. Appl. Math.*, 24(2):179–211, 2013.
- [HN75] B. Halphen and Q.S. Nguyen. Sur les matériaux standards généralisés. *J. Mécanique*, 14:39–63, 1975.
- [IT79] A. D. Ioffe and V. M. Tihomirov. *Theory of extremal problems*, volume 6 of *Studies in Mathematics and its Applications*. North-Holland Publishing Co., Amsterdam, 1979. Translated from the Russian by Karol Makowski.



- [KMR06] M. Kočvara, A. Mielke, and T. Roubíček. A rate-independent approach to the delamination problem. *Math. Mech. Solids*, 11:423–447, 2006.
- [KN15] D. Knees and M. Negri. Convergence of alternate minimization schemes for phase field fracture and damage. 2015. Preprint. Available at [cvgmt.sns.it](http://cvgmt.sns.it).
- [KP11] P. Krejčí and A. Petrov. Existence and uniqueness results for a class of dynamic elasto-plastic contact problems. *J. Math. Anal. Appl.*, 408:125–139, 2011.
- [KRZ13] D. Knees, R. Rossi, and C. Zanini. A vanishing viscosity approach to a rate-independent damage model. *Math. Models Methods Appl. Sci.*, 23(4):565–616, 2013.
- [LS65] J.-L. Lions and W.A. Strauss. Some non-linear evolution equations. *Bull. Soc. Math. France*, 93:43–96, 1965.
- [Mie05] A. Mielke. Evolution in rate-independent systems (Ch.6). In C.M. Dafermos and E. Feireisl, editors, *Handbook of Differential Equations, Evolutionary Equations, vol. 2*, pages 461–559. Elsevier B.V., Amsterdam, 2005.
- [Mie11] A. Mielke. Differential, energetic and metric formulations for rate-independent processes. In *Nonlinear PDE's and applications*, volume 2028 of *Lecture Notes in Mathematics*, pages xiv+224. Springer, Heidelberg; Fondazione C.I.M.E., Florence, 2011. Notes of the C.I.M.E. Summer School held in Cetraro, June 23–28, 2008. Edited by L. Ambrosio and G. Savaré, Centro Internazionale Matematico Estivo (C.I.M.E.) Summer Schools.
- [Mie16] A. Mielke. On evolutionary  $\Gamma$ -convergence for gradient systems. In *Macroscopic and Large Scale Phenomena: Coarse Graining, Mean Field Limits and Ergodicity*, Lecture Notes in Appl. Math. Mech., pages 187–249. Springer, 2016. Edited by A. Muntean, J. Rademacher, A. Zagaris; proceedings of the Summer School in Twente University, June 2012.
- [MM05] A. Mainik and A. Mielke. Existence results for energetic models for rate-independent systems. *Calc. Var. PDEs*, 22:73–99, 2005.
- [MR07] A. Mielke and R. Rossi. Existence and uniqueness results for a class of rate-independent hysteresis problems. *Math. Models Methods Appl. Sci.*, 17(1):81–123, 2007.
- [MR15] A. Mielke and T. Roubíček. *Rate-Independent Systems: Theory and Application*, volume 193 of *Applied Mathematical Sciences*. Springer, 2015.
- [MRS08] A. Mielke, T. Roubíček, and U. Stefanelli.  $\Gamma$ -limits and relaxations for rate-independent evolutionary problems. *Calc. Var. Partial Differential Equations*, 31:387–416, 2008.
- [MRS13a] A. Mielke, R. Rossi, and G. Savaré. Balanced viscosity (BV) solutions to infinite-dimensional rate-independent systems. *J. Eur. Math. Soc.*, to appear. *WIAS-Preprint 1845*, 2013.
- [MRS13b] A. Mielke, R. Rossi, and G. Savaré. Nonsmooth analysis of doubly nonlinear evolution equations. *Calc. Var. Partial Differential Equations*, 46(1-2):253–310, 2013.
- [MT04] A. Mielke and F. Theil. On rate-independent hysteresis models. *NoDEA Nonlinear Differential Equations Appl.*, 11(2):151–189, 2004.
- [Pet12] A. Petrov. On the numerical approximation of a viscoelastodynamic problem with unilateral constraints. *Comput. Methods Appl. Mech. Engrg.*, 205–208:162–168, 2012.
- [PS09] A. Petrov and M. Schatzman. Mathematical results on existence for viscoelastodynamic problems with unilateral constraints. *SIAM J. Math. Anal.*, 40:1882–1904, 2008/09.
- [RMS08] R. Rossi, A. Mielke, and G. Savaré. A metric approach to a class of doubly nonlinear evolution equations and applications. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 7(1):97–169, 2008.
- [Rou05] T. Roubíček. *Nonlinear Partial Differential Equations with Applications*. Birkhäuser, 2005.
- [Rou09] T. Roubíček. Rate-independent processes in viscous solids at small strains. *Math. Methods Appl. Sci.*, 32(7):825–862, 2009.
- [Rou10] T. Roubíček. Thermodynamics of rate-independent processes in viscous solids at small strains. *SIAM J. Math. Anal.*, 42(1):256–297, 2010.
- [RR11] R. Rossi and T. Roubíček. Thermodynamics and analysis of rate-independent adhesive contact at small strains. *Nonlinear Anal.*, 74(10):3159–3190, 2011.
- [RS06] R. Rossi and G. Savaré. Gradient flows of non convex functionals in Hilbert spaces and applications. *ESAIM Control Optim. Calc. Var.*, 12(3):564–614, 2006.
- [RSV13] T. Roubíček, O. Souček, and R. Vodička. A model of rupturing lithospheric faults with reoccurring earthquakes. *SIAM J. Appl. Math.*, 73(4):1460–1488, 2013.
- [RT15] T. Roubíček and G. Tomassetti. Thermomechanics of damageable materials under diffusion: modeling and analysis. *Zeit. Angew. Math. Phys.*, 66:3535–3572, 2015.
- [RT16] R. Rossi and M. Thomas. Coupling rate-independent and rate-dependent processes: Evolutionary  $\Gamma$ -convergence. *In preparation*, 2016.
- [RTP15] T. Roubíček, M. Thomas, and C.G. Panagiotopoulos. Stress-driven local-solution approach to quasistatic brittle delamination. *Nonlinear Anal. Real World Appl.*, 22:645–663, 2015.
- [RV16] T. Roubíček and J. Valdman. Perfect plasticity with damage and healing at small strains, its modeling, analysis, and computer implementation. *SIAM J. Appl. Math.*, 76(1):314–340, 2016.
- [Sim87] J. Simon. Compact sets in the space  $L^p(0, T; B)$ . *Ann. Mat. Pura Appl.*, 146:65–96, 1987.
- [Ste08] Ulisse Stefanelli. The Brezis-Ekeland principle for doubly nonlinear equations. *SIAM J. Control Optim.*, 47(3):1615–1642, 2008.
- [Tho13] M. Thomas. Quasistatic damage evolution with spatial BV-regularization. *Discrete Contin. Dyn. Syst. Ser. S*, 6(1):235–255, 2013.

- [TM10] M. Thomas and A. Mielke. Damage of nonlinearly elastic materials at small strain: existence and regularity results. *Zeit. Angew. Math. Mech.*, 90(2):88–112, 2010.
- [Val90] M. Valadier. Young measures. In *Methods of nonconvex analysis (Varenna, 1989)*, pages 152–188. Springer, Berlin, 1990.

R. ROSSI, DEPARTMENT DIMI, UNIVERSITÀ DEGLI STUDI DI BRESCIA, VIA VALOTTI 9, 25133 BRESCIA - ITALY

*E-mail address:* `riccarda.rossi@unibs.it`

M. THOMAS, WEIERSTRASS INSTITUTE FOR APPLIED ANALYSIS AND STOCHASTICS, MOHRENSTR. 39, 10117 BERLIN - GERMANY

*E-mail address:* `marita.thomas@wias-berlin.de`