

# Interplay of viscosity and dry friction in rate-independent evolutions with nonconvex energies

Riccarda Rossi  
(Università di Brescia)

**joint work (in progress) with**  
Alexander Mielke (WIAS & Humboldt-Universität – Berlin)  
Giuseppe Savaré (Università di Pavia)

WIAS, Berlin, 22.04.2009

## Rate-independent evolutions in the applications

1. quasistatic propagation of fracture [Bourdin, Cagnetti, Chambolle, Dal Maso, Francfort, Giacomini, Knees, Larsen, Lazzaroni, Marigo, Mielke, Negri, Ortner, Ponsiglione, Toader, Zanini .....] ]
2. quasistatic phase transformations in shape memory alloys (SMA) [Auricchio, Levitas, Mainik, Mielke, Theil, Roubíček, Stefanelli....]
3. elastoplasticity: linearized & finite-strain [Dal Maso, DeSimone, Fiaschi, Francfort, Mora, Morini, Mielke, Mainik, Roubíček...]
4. damage [Francfort, Garroni, Larsen, Mielke, Roubíček, Thomas...]
5. delamination [Kočvara, Mielke, Roubíček, Scardia, Zanini...]
6. ferromagnetism, ferroelectricity, superconductivity [Mielke, Schmid, Timofte...]
7. shape evolution of debonding membranes [Bucur, Buttazzo..]

### In these applications

- ▶ Typical **energies** are **nonsmooth & nonconvex**
- ▶ **Ambient spaces** may lack **a natural linear structure** (e.g. in crack propagation)

# Energetic formulation for rate-independent evolutions

## Weak formulations ("derivative-free")

Based on

- ▶ **energetic balance** (energy identity)
- ▶ **stability** conditions
- ▶ possibly enforcing **irreversibility**

# Energetic formulation for rate-independent evolutions

## Abstract approach by Mielke

- ♣ Ambient space  $\mathcal{U}$  topological space
- ♣ Dissipation:  $\mathcal{D} : \mathcal{U} \times \mathcal{U} \rightarrow [0, +\infty]$  pseudo-distance
- ♣ Energy:  $\mathcal{E} : (0, T) \times \mathcal{U} \rightarrow (-\infty, +\infty]$

# Energetic formulation for rate-independent evolutions

## Abstract approach by Mielke

- ♣ Ambient space  $\mathcal{U}$  topological space
- ♣ Dissipation:  $\mathcal{D} : \mathcal{U} \times \mathcal{U} \rightarrow [0, +\infty]$  pseudo-distance
- ♣ Energy:  $\mathcal{E} : (0, T) \times \mathcal{U} \rightarrow (-\infty, +\infty]$

## Energetic formulation [Mielke-Theil'99,'04], [Mielke-Theil-Levitas'02],

[Mainik-Mielke'05]

**Global energetic** solutions  $u : [0, T] \rightarrow \mathcal{U}$ : **global stability condition** & **energy balance**

$$\mathcal{E}(t, u(t)) - \mathcal{E}(t, z) \leq \mathcal{D}(u(t), z) \quad \forall z \in \mathcal{U},$$

$$\mathcal{E}(t, u(t)) + \text{Diss}_{\mathcal{D}}(u, [0, t]) = \mathcal{E}(0, u(0)) + \int_0^t \partial_t \mathcal{E}(r, u(r)) \, dr.$$

$\text{Diss}_{\mathcal{D}}$  being the global dissipation functional associated with  $\mathcal{D}$

## The convex case

In [Mielke-Theil'04]: if

- ▶ ambient space  $\mathcal{U}$  is a **reflexive** Banach space  $\mathcal{B}$
- ▶  $\mathcal{E}(t, \cdot)$  (uniformly) **convex** & **smooth**
- ▶  $\mathcal{D}$  induced by  $\Psi_1 : \mathcal{B} \rightarrow [0, +\infty)$  **convex & 1-positively homogeneous**  
( $\Psi_1 \sim \|\cdot\|$ )

## The convex case

In [Mielke-Theil'04]: if

- ▶ ambient space  $\mathcal{U}$  is a **reflexive** Banach space  $\mathcal{B}$
- ▶  $\mathcal{E}(t, \cdot)$  (uniformly) **convex** & **smooth**
- ▶  $\mathcal{D}$  induced by  $\Psi_1 : \mathcal{B} \rightarrow [0, +\infty)$  **convex & 1-positively homogeneous**  
( $\Psi_1 \sim \|\cdot\|$ )

then

- ▶  $u \in AC([0, T]; \mathcal{B})$  (even Lipschitz in time)
- ▶ the energetic formulation is **equivalent** to the **doubly nonlinear** equation

$$\partial\Psi_1(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 \quad \text{in } \mathcal{B}' \quad t \in (0, T)$$

**(subdifferential formulation)**

with  $\partial\mathcal{E}(t, \cdot)$  **convex subdifferential** of  $\mathcal{E}(t, \cdot)$  w.r.t.  $u$

$$\xi \in \partial\mathcal{E}(t, u) \Leftrightarrow \mathcal{E}(t, w) - \mathcal{E}(t, u) \geq \langle \xi, w - u \rangle \text{ for all } w \in \mathcal{B}$$

## The convex case

In [Mielke-Theil'04]: if

- ▶ ambient space  $\mathcal{U}$  is a **reflexive** Banach space  $\mathcal{B}$
- ▶  $\mathcal{E}(t, \cdot)$  (uniformly) **convex** & **smooth**
- ▶  $\mathcal{D}$  induced by  $\Psi_1 : \mathcal{B} \rightarrow [0, +\infty)$  **convex & 1-positively homogeneous**  
( $\Psi_1 \sim \|\cdot\|$ )



## The convex case

In [Mielke-Theil'04]: if

- ▶ ambient space  $\mathcal{U}$  is a **reflexive** Banach space  $\mathcal{B}$
- ▶  $\mathcal{E}(t, \cdot)$  (uniformly) **convex & smooth**
- ▶  $\mathcal{D}$  induced by  $\Psi_1 : \mathcal{B} \rightarrow [0, +\infty)$  **convex & 1-positively homogeneous**  
( $\Psi_1 \sim \|\cdot\|$ )

then

- ▶  $u \in AC([0, T]; \mathcal{B})$  (even Lipschitz in time)
- ▶ the energetic formulation is **equivalent** to the **doubly nonlinear** equation

$$\partial\Psi_1(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 \quad \text{in } \mathcal{B}' \quad t \in (0, T)$$

**(subdifferential formulation)**

with  $\partial\mathcal{E}(t, \cdot)$  **convex subdifferential** of  $\mathcal{E}(t, \cdot)$  w.r.t.  $u$

$$\xi \in \partial\mathcal{E}(t, u) \Leftrightarrow \mathcal{E}(t, w) - \mathcal{E}(t, u) \geq \langle \xi, w - u \rangle \text{ for all } w \in \mathcal{B}$$

## The nonconvex case: towards local stability

If  $\mathcal{E}(t, \cdot)$  is nonconvex

- ▶  $\Psi_1$ , 1-homogeneous, has a **linear growth** at  $\infty$   $\rightsquigarrow$

$$\partial\Psi_1(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 \quad \text{in } \mathcal{B}' \quad t \in (0, T),$$

$\rightsquigarrow$  we only expect  $u \in BV(0, T; \mathcal{B})$  ( $u$  may **jump!!!**)

## The nonconvex case: towards local stability

If  $\mathcal{E}(t, \cdot)$  is nonconvex

- ▶  $\Psi_1$ , 1-homogeneous, has a **linear growth** at  $\infty \rightsquigarrow$

$$\partial\Psi_1(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 \quad \text{in } \mathcal{B}' \quad t \in (0, T),$$

$\rightsquigarrow$  we only expect  $u \in BV(0, T; \mathcal{B})$  ( $u$  may **jump!!!**)

### Problem with the Global Energetic formulation

**Global stability** forces global energetic solutions to **jump too early** and overcome **too large energy barriers** to avoid energy losses

# Bad Vs. Good jumps

## The simplest nonconvex case

$$\begin{cases} B = \mathbb{R}, & \Psi_1(v) = |v| \quad \forall v \in \mathbb{R} \\ \mathcal{E}(t, u) = \mathcal{W}(u) - \ell(t)u & \forall (t, u) \in [0, T] \times \mathbb{R} \end{cases}$$

- ▶  $\mathcal{W}$  **double well** potential
- ▶  $\ell \in C^1([0, T]) \sim$  external loading

$$\text{Sign}(u'(t)) + \mathcal{W}'(u(t)) \ni \ell(t), \quad t \in (0, T)$$

## Bad Vs. Good jumps


### The simplest nonconvex case

$$\begin{cases} B = \mathbb{R}, & \Psi_1(v) = |v| \quad \forall v \in \mathbb{R} \\ \mathcal{E}(t, u) = \mathcal{W}(u) - \ell(t)u & \forall (t, u) \in [0, T] \times \mathbb{R} \end{cases}$$

- ▶  $\mathcal{W}$  **double well** potential
- ▶  $\ell \in C^1([0, T]) \sim$  external loading

$$\text{Sign}(u'(t)) + \mathcal{W}'(u(t)) \ni \ell(t), \quad t \in (0, T)$$

Convexification  $\mathcal{W}^{**}$  of  $\mathcal{W}$



## Bad Vs. Good jumps


### The simplest nonconvex case

$$\begin{cases} B = \mathbb{R}, & \Psi_1(v) = |v| \quad \forall v \in \mathbb{R} \\ \mathcal{E}(t, u) = \mathcal{W}(u) - \ell(t)u & \forall (t, u) \in [0, T] \times \mathbb{R} \end{cases}$$

- ▶  $\mathcal{W}$  **double well** potential
- ▶  $\ell \in C^1([0, T]) \sim$  external loading

$$\text{Sign}(u'(t)) + \mathcal{W}'(u(t)) \ni \ell(t), \quad t \in (0, T)$$

Global solutions are given by  $u(t) = (D\mathcal{W}^{**})^{-1}(\ell(t) - 1)$ : jumping too early!



## Bad Vs. Good jumps


The simplest nonconvex case

$$\begin{cases} B = \mathbb{R}, & \Psi_1(v) = |v| \quad \forall v \in \mathbb{R} \\ \mathcal{E}(t, u) = \mathcal{W}(u) - \ell(t)u & \forall (t, u) \in [0, T] \times \mathbb{R} \end{cases}$$

- ▶  $\mathcal{W}$  **double well** potential
- ▶  $\ell \in C^1([0, T]) \sim$  external loading

$$\text{Sign}(u'(t)) + \mathcal{W}'(u(t)) \ni \ell(t), \quad t \in (0, T)$$

We aim to model the "right" hysteresis dynamics



## The vanishing viscosity approach

### Aim:

Formulation for rate-independent problems

- ▶ **modelling** only **"natural" jumps** (due to  $u \in BV(0, T; \mathcal{B})$ ),
- ▶ leading to **solutions jumping later** than global energetic solutions



# The vanishing viscosity approach

## Aim:

Formulation for rate-independent problems

- ▶ **modelling** only **"natural" jumps** (due to  $u \in BV(0, T; \mathcal{B})$ ),
- ▶ leading to **solutions jumping later** than global energetic solutions

## The vanishing viscosity method:

select rate-independent evolutions arising in the limit of viscous regularizations. In:

- ▶ plasticity with softening: [Dal Maso-DeSimone-Mora-Morini'08], nonconvex elastoplasticity: [Fiaschi'09].
- ▶ crack propagation: [Toader-Zanini'06], [Knees-Mielke-Zanini'07, '08], [Cagnetti'07].
- ▶ general rate-independent evolution with discontinuous inputs: [Krejčí-Liero'07].

# The vanishing viscosity approach

## Aim:

Formulation for rate-independent problems

- ▶ **modelling** only **"natural" jumps** (due to  $u \in BV(0, T; \mathcal{B})$ ),
- ▶ leading to **solutions jumping later** than global energetic solutions

## The vanishing viscosity method:

select rate-independent evolutions arising in the limit of viscous regularizations. In:

- ▶ plasticity with softening: [Dal Maso-DeSimone-Mora-Morini'08], nonconvex elastoplasticity: [Fiaschi'09].
- ▶ crack propagation: [Toader-Zanini'06], [Knees-Mielke-Zanini'07, '08], [Cagnetti'07].
- ▶ general rate-independent evolution with discontinuous inputs: [Krejčí-Liero'07].

The **vanishing viscosity** approach leads to formulations oriented towards **local stability** [Dal Maso-Toader'02], [Negri-Ortner'07], (crack propagation), [Stefanelli'08]..

# The vanishing viscosity approach

## Problem

In the vanishing viscosity limit:

- ▶ local stability
- ▶ energy inequality

may not be enough for controlling jumps.  $\zeta$  Which further conditions better describe them?

# The vanishing viscosity approach

## Problem

In the vanishing viscosity limit:

- ▶ local stability
- ▶ energy inequality

may not be enough for controlling jumps.  $\zeta$  Which further conditions better describe them?

## An abstract approach

Approximation by vanishing viscosity of

- ▶ doubly nonlinear rate-independent equations (**subdifferential formulation**)
- ▶ in an abstract Banach setting, with general **nonconvex** energy functionals

extending **previous analysis by Efendiev-Mielke** [Efendiev-Mielke'06]

# The technique by Efendiev-Mielke

## A reparametrization technique

- ▶ We are modelling systems with two **time scales**:
  - ▶ a scale **intrinsic** to the system, **fast time scale**
  - ▶ the **slow time scale** of the **external loading**  $\sim \partial_t \mathcal{E}$  (**dominating scale**)  $\rightsquigarrow$  viscous dissipation is negligible!

# The technique by Efendiev-Mielke

## A reparametrization technique

- ▶ We are modelling systems with two **time scales**:
  - ▶ a scale **intrinsic** to the system, **fast time scale**
  - ▶ the **slow time scale** of the **external loading**  $\sim \partial_t \mathcal{E}$  (**dominating scale**)  $\rightsquigarrow$  viscous dissipation is negligible!
- ▶ Jumps in the vanishing viscosity limit correspond to **viscous transitions** between stable states

# The technique by Efendiev-Mielke

## A reparametrization technique

- ▶ We are modelling systems with two **time scales**:
  - ▶ a scale **intrinsic** to the system, **fast time scale**
  - ▶ the **slow time scale** of the **external loading**  $\sim \partial_t \mathcal{E}$  (**dominating scale**)  $\rightsquigarrow$  viscous dissipation is negligible!
- ▶ Jumps in the vanishing viscosity limit correspond to **viscous transitions** between stable states
- ▶ To capture the viscous transition path: **NOT SHRINK** jumps at a point, look at curves with their **arc length parametrization**
- ▶ Asymptotic analysis of (reparametrized) trajectories in an extended phase space

# Nonconvex energies

[Mielke-R.-Savaré, in progress] In infinite dimensions:



## Nonconvex energies

[Mielke-R.-Savaré, in progress] In infinite dimensions:

Approximate

$$\partial\Psi(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 \quad \text{in } \mathcal{B}' \quad t \in (0, T), \quad (\text{DNE})$$

$\Psi$  **1-positively homogeneous**

## Nonconvex energies

[Mielke-R.-Savaré, in progress] In infinite dimensions:

with, as  $\varepsilon \searrow 0$

$$\partial\Psi_\varepsilon(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 \quad \text{in } \mathcal{B}' \quad t \in (0, T), \quad (\text{DNE})$$

$\Psi_\varepsilon$  with superlinear growth

## Nonconvex energies

[Mielke-R.-Savaré, in progress] In infinite dimensions:

with, as  $\varepsilon \searrow 0$

$$\partial\Psi_\varepsilon(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 \quad \text{in } \mathcal{B}' \quad t \in (0, T), \quad (\text{DNE})$$

$\Psi_\varepsilon$  with superlinear growth

Nonconvexity & nonsmoothness of  $\mathcal{E}(t, \cdot)$  affect

both the viscous and the rate-independent equation

## Two issues

## Nonconvex energies

[Mielke-R.-Savaré, in progress] In infinite dimensions:

with, as  $\varepsilon \searrow 0$

$$\partial\Psi_\varepsilon(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 \quad \text{in } \mathcal{B}' \quad t \in (0, T), \quad (\text{DNE})$$

$\Psi_\varepsilon$  with superlinear growth

Nonconvexity & nonsmoothness of  $\mathcal{E}(t, \cdot)$  affect

both the viscous and the rate-independent equation

### Two issues

- ▶  $u \mapsto \mathcal{E}(t, u)$  is nonconvex  $\Rightarrow$  choice of a suitable subdifferential notion  $\partial\mathcal{E}$

## Nonconvex energies

[Mielke-R.-Savaré, in progress] In infinite dimensions:

with, as  $\varepsilon \searrow 0$

$$\partial\Psi_\varepsilon(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 \quad \text{in } \mathcal{B}' \quad t \in (0, T), \quad (\text{DNE})$$

$\Psi_\varepsilon$  with superlinear growth

Nonconvexity & nonsmoothness of  $\mathcal{E}(t, \cdot)$  affect

both the viscous and the rate-independent equation

### Two issues

- ▶  $u \mapsto \mathcal{E}(t, u)$  is nonconvex  $\Rightarrow$  choice of a suitable subdifferential notion  $\partial\mathcal{E}$
- ▶ jumps in the rate-independent evolution  $\Rightarrow$  to be modelled via vanishing viscosity

# Outline

- ♣ existence results for "viscous" doubly nonlinear equations with nonconvex energies
- ♣ two ideas on the vanishing viscosity analysis

# Outline

- ♣ existence results for "viscous" doubly nonlinear equations with nonconvex energies
- ♣ two ideas on the vanishing viscosity analysis
- ♣ parametrized rate-independent evolutions:
  - ▶ **local stability**
  - ▶ **jumps**  $\leftrightarrow$  **viscous transitions** between metastable states

## Which subdifferential notion? Heuristics

$$\begin{cases} \partial\Psi_\varepsilon(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 & \text{in } \mathcal{B}' \quad t \in (0, T), \\ u(0) = u_0 \end{cases} \quad (\text{DNE})$$

$\Psi_\varepsilon$  with superlinear growth



## Which subdifferential notion? Heuristics

$$\begin{cases} \partial\Psi_\varepsilon(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 & \text{in } \mathcal{B}' \quad t \in (0, T), \\ u(0) = u_0 \end{cases} \quad (\text{DNE})$$

$\Psi_\varepsilon$  with superlinear growth

### Step 1: approximation by time discretization

- ▶ Fix time step  $\tau > 0 \rightsquigarrow$  partition  $0 < t_1 < \dots < t_n = n\tau < \dots < t_N = T$  of  $(0, T)$
- ▶ **Discrete solutions:** the recursive minimization problem

Find  $U_\tau^0, U_\tau^1, \dots, U_\tau^N \in \mathcal{B}$ :

$$U_\tau^n \in \operatorname{Argmin}_{U \in \mathcal{B}} \left\{ \tau \Psi_\varepsilon \left( \frac{U - U_\tau^{n-1}}{\tau} \right) + \mathcal{E}(t_n, U) \right\}, \quad U_\tau^0 := u_0$$

has a solution if  $\mathcal{E}(t, \cdot)$  is **coercive** (e.g.,  $\mathcal{E}(t, \cdot) + \|\cdot\|_{\mathcal{B}}^2$  has compact sublevels)

- ▶ **Discrete Euler-Lagrange equation:**  $\partial_{\mathcal{F}} \mathcal{E}(t, \cdot)$  the **Fréchet subdifferential** of  $\mathcal{E}(t, \cdot)$

$$\partial\Psi_\varepsilon \left( \frac{U_\tau^n - U_\tau^{n-1}}{\tau} \right) + \partial_{\mathcal{F}} \mathcal{E}(t_n, U_\tau^n) \ni 0,$$

# The Fréchet subdifferential

Idea: "localize" the convex subdifferential

# The Fréchet subdifferential

Idea: "localize" the convex subdifferential

## The convex subdifferential

Given  $u \in D(\mathcal{E}(t, \cdot))$ ,

$$\xi \in \partial \mathcal{E}(t, u) \Leftrightarrow \mathcal{E}(t, w) - \mathcal{E}(t, u) \geq \langle \xi, w - u \rangle \text{ for all } w \in \mathcal{B}$$

# The Fréchet subdifferential

Idea: "localize" the convex subdifferential

## The Fréchet subdifferential

Given  $u \in D(\mathcal{E}(t, \cdot))$ ,

$$\xi \in \partial_F \mathcal{E}(t, u) \Leftrightarrow \liminf_{w \rightarrow u} \frac{\mathcal{E}(t, w) - \mathcal{E}(t, u) \geq \langle \xi, w - u \rangle}{\|w - u\|_{\mathcal{B}}} \geq 0$$

# The Fréchet subdifferential

Idea: "localize" the convex subdifferential

## The Fréchet subdifferential

Given  $u \in D(\mathcal{E}(t, \cdot))$ ,

$$\xi \in \partial_F \mathcal{E}(t, u) \Leftrightarrow \liminf_{w \rightarrow u} \frac{\mathcal{E}(t, w) - \mathcal{E}(t, u) \geq \langle \xi, w - u \rangle}{\|w - u\|_{\mathcal{B}}} \geq 0$$

- ▶  $\partial_F \mathcal{E}(t, u) \equiv \partial \mathcal{E}(t, u)$  if  $\mathcal{E}(t, \cdot)$  is convex
- ▶  $\partial_F \mathcal{E}(t, u)$  is **convex** for all  $u \in D(\mathcal{E}(t, \cdot))$
- ▶ The map  $u \mapsto \partial_F \mathcal{E}(t, u)$  is **not strongly-weakly closed** in the sense of graphs!!!

## Heuristics

### Step 2: estimates on the approximate solutions

- ▶ **Approximate solutions:** interpolants on  $(0, T)$  of the discrete solutions  $\{U_\tau^k\}_{k=1}^n$ :
  - ▶  $\{\bar{U}_\tau\}$  piecewise constant;
  - ▶  $\{\hat{U}_\tau\}$  piecewise linear.
- ▶ from the discrete Euler-Lagrange equation  $\rightsquigarrow$  **Approximate equation:**

$$\partial\Psi_\varepsilon\left(\hat{U}'_\tau(t)\right) + \partial_F\mathcal{E}(t, \bar{U}_\tau(t)) \ni 0 \quad t \in (0, T)$$

- ▶ **A priori estimates + compactness** (strong for  $\hat{U}_\tau$ , weak for  $\hat{U}'_\tau$ )
- ▶ convergence to some limit curve  $u \in W^{1,1}(0, T; \mathcal{B})$

**BUT:** you can't pass to the limit in

$$-\partial\Psi_\varepsilon\left(\hat{U}'_\tau(t)\right) \ni \partial_F\mathcal{E}(t, \bar{U}_\tau(t)) \ni 0 \quad t \in (0, T)$$

because  $u \rightharpoonup \partial_F\mathcal{E}(t, u)$  **in general** is **NOT strongly-weakly closed!!**

## The limiting subdifferential

### First idea:

Consider (a version of) the **strong-weak closure** of  $\partial_F \mathcal{E}(t, u)$ :

**the limiting subdifferential**  $\partial \mathcal{E}(t, \cdot)$  [Mordukhovich'84]

given  $u \in D(en(t, \cdot))$ ,

$$\xi \in \partial \mathcal{E}(t, u) \Leftrightarrow \exists \{u_n\}, \{\xi_n\} \subset \mathcal{B} : \begin{cases} \xi_n \in \partial_F \mathcal{E}(t, u_n) \quad \forall n \in \mathbb{N}, \\ u_n \rightarrow u, \\ \xi_n \rightarrow \xi, \\ \sup_n \mathcal{E}(t, u_n) < +\infty \end{cases}$$

The **limiting subdifferential** is our notion of subdifferential!

## Heuristics

### Second idea:

- Instead of passing to the limit in the **pointwise** equation

$$-\partial\Psi_\varepsilon\left(\widehat{U}'_\tau(t)\right) \ni \partial_F\mathcal{E}(t, \overline{U}_\tau(t)) \ni 0 \quad t \in (0, T)$$

pass to the limit in the **approximate energy inequality** (**technical point!**)

$$\begin{aligned} \int_0^t \Psi_\varepsilon\left(\widehat{U}'_\tau(s)\right) ds + \int_0^t \Psi_\varepsilon^*\left(-\partial_F\mathcal{E}(s, \overline{U}_\tau(s))\right) ds + \mathcal{E}(t, \overline{U}_\tau(t)) \\ \leq \mathcal{E}(0, u_0) + \int_0^t \partial_t\mathcal{E}(s, \overline{U}_\tau(s)) ds \end{aligned}$$

with  $\Psi_\varepsilon^*$  the Fenchel-Moreau conjugate of  $\Psi_\varepsilon$



## Heuristics

### Second idea:

- ▶ Instead of passing to the limit in the **pointwise** equation

$$-\partial\Psi_\varepsilon\left(\widehat{U}'_\tau(t)\right) \ni \partial_F\mathcal{E}(t, \overline{U}_\tau(t)) \ni 0 \quad t \in (0, T)$$

pass to the limit in the **approximate energy inequality** (**technical point!**)

$$\begin{aligned} \int_0^t \Psi_\varepsilon\left(\widehat{U}'_\tau(s)\right) ds + \int_0^t \Psi_\varepsilon^*\left(-\partial_F\mathcal{E}(s, \overline{U}_\tau(s))\right) ds + \mathcal{E}(t, \overline{U}_\tau(t)) \\ \leq \mathcal{E}(0, u_0) + \int_0^t \partial_t\mathcal{E}(s, \overline{U}_\tau(s)) ds \end{aligned}$$

with  $\Psi_\varepsilon^*$  the Fenchel-Moreau conjugate of  $\Psi_\varepsilon$

- ▶ By **LOWER-SEMICONITINUITY** obtain the **limit energy inequality**

$$\begin{aligned} \int_0^t \Psi_\varepsilon(u'(s)) ds + \int_0^t \Psi_\varepsilon^*(-\partial\mathcal{E}(s, u(s))) ds + \mathcal{E}(t, u(t)) \\ \leq \mathcal{E}(0, u_0) + \int_0^t \partial_t\mathcal{E}(s, u(s)) ds \end{aligned}$$

## Heuristics

### Third idea:

If  $\partial\mathcal{E}(t, \cdot)$  fulfils the **chain rule**

$$u \in W^{1,1}(0, T; \mathcal{B}) \Rightarrow \frac{d}{dt} \mathcal{E}(t, u(t)) = \langle \partial\mathcal{E}(t, u(t)), u'(t) \rangle + \partial_t \mathcal{E}(t, u(t)) \text{ a.e. in } (0, T)$$

## Heuristics

### Third idea:

If  $\partial\mathcal{E}(t, \cdot)$  fulfils the **chain rule**

$$u \in W^{1,1}(0, T; \mathcal{B}) \Rightarrow \frac{d}{dt} \mathcal{E}(t, u(t)) = \langle \partial\mathcal{E}(t, u(t)), u'(t) \rangle + \partial_t \mathcal{E}(t, u(t)) \text{ a.e. in } (0, T)$$

then

$$\int_0^t \Psi_\varepsilon(u'(s)) ds + \int_0^t \Psi_\varepsilon^*(-\partial\mathcal{E}(s, u(s))) ds + \mathcal{E}(t, u(t)) \leq \mathcal{E}(0, u_0) + \int_0^t \partial_t \mathcal{E}(s, u(s)) ds$$

## Heuristics

### Third idea:

If  $\partial\mathcal{E}(t, \cdot)$  fulfils the **chain rule**

$$u \in W^{1,1}(0, T; \mathcal{B}) \Rightarrow \frac{d}{dt} \mathcal{E}(t, u(t)) = \langle \partial\mathcal{E}(t, u(t)), u'(t) \rangle + \partial_t \mathcal{E}(t, u(t)) \text{ a.e. in } (0, T)$$

then

$$\begin{aligned} \int_0^t \Psi_\varepsilon(u'(s)) ds + \int_0^t \Psi_\varepsilon^*(-\partial\mathcal{E}(s, u(s))) ds + \mathcal{E}(t, u(t)) &\leq \mathcal{E}(0, u_0) + \int_0^t \partial_t \mathcal{E}(s, u(s)) ds \\ &= \mathcal{E}(t, u(t)) - \int_0^t \langle \partial\mathcal{E}(s, u(s)), u'(s) \rangle ds \end{aligned}$$

## Heuristics

### Third idea:

If  $\partial\mathcal{E}(t, \cdot)$  fulfils the **chain rule**

$$u \in W^{1,1}(0, T; \mathcal{B}) \Rightarrow \frac{d}{dt} \mathcal{E}(t, u(t)) = \langle \partial\mathcal{E}(t, u(t)), u'(t) \rangle + \partial_t \mathcal{E}(t, u(t)) \text{ a.e. in } (0, T)$$

then

$$\begin{aligned} \int_0^t \Psi_\varepsilon(u'(s)) ds + \int_0^t \Psi_\varepsilon^*(-\partial\mathcal{E}(s, u(s))) ds + \mathcal{E}(t, u(t)) &\leq \mathcal{E}(0, u_0) + \int_0^t \partial_t \mathcal{E}(s, u(s)) ds \\ &= \mathcal{E}(t, u(t)) - \int_0^t \langle \partial\mathcal{E}(s, u(s)), u'(s) \rangle ds \end{aligned}$$

$$\int_0^t \left( \Psi_\varepsilon(u'(s)) + \Psi_\varepsilon^*(-\partial\mathcal{E}(s, u(s))) + \langle \partial\mathcal{E}(s, u(s)), u'(s) \rangle \right) ds = 0$$

## Heuristics

### Third idea:

If  $\partial\mathcal{E}(t, \cdot)$  fulfils the **chain rule**

$$u \in W^{1,1}(0, T; \mathcal{B}) \Rightarrow \frac{d}{dt} \mathcal{E}(t, u(t)) = \langle \partial\mathcal{E}(t, u(t)), u'(t) \rangle + \partial_t \mathcal{E}(t, u(t)) \text{ a.e. in } (0, T)$$

then

$$\begin{aligned} \int_0^t \Psi_\varepsilon(u'(s)) ds + \int_0^t \Psi_\varepsilon^*(-\partial\mathcal{E}(s, u(s))) ds + \mathcal{E}(t, u(t)) &\leq \mathcal{E}(0, u_0) + \int_0^t \partial_t \mathcal{E}(s, u(s)) ds \\ &= \mathcal{E}(t, u(t)) - \int_0^t \langle \partial\mathcal{E}(s, u(s)), u'(s) \rangle ds \end{aligned}$$

$$\Psi_\varepsilon(u'(t)) + \Psi_\varepsilon^*(-\partial\mathcal{E}(t, u(t))) + \langle \partial\mathcal{E}(t, u(t)), u'(t) \rangle = 0 \text{ a.e. in } (0, T)$$

## Heuristics

### Third idea:

If  $\partial\mathcal{E}(t, \cdot)$  fulfils the **chain rule**

$$u \in W^{1,1}(0, T; \mathcal{B}) \Rightarrow \frac{d}{dt} \mathcal{E}(t, u(t)) = \langle \partial\mathcal{E}(t, u(t)), u'(t) \rangle + \partial_t \mathcal{E}(t, u(t)) \quad \text{a.e. in } (0, T)$$

then

$$\begin{aligned} \int_0^t \Psi_\varepsilon(u'(s)) ds + \int_0^t \Psi_\varepsilon^*(-\partial\mathcal{E}(s, u(s))) ds + \mathcal{E}(t, u(t)) &\leq \mathcal{E}(0, u_0) + \int_0^t \partial_t \mathcal{E}(s, u(s)) ds \\ &= \mathcal{E}(t, u(t)) - \int_0^t \langle \partial\mathcal{E}(s, u(s)), u'(s) \rangle ds \end{aligned}$$

whence

$$\partial\Psi_\varepsilon(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 \quad \text{for a.a. } t \in (0, T).$$

and the **energy identity**

$$\int_0^t \Psi_\varepsilon(u'(s)) ds + \int_0^t \Psi_\varepsilon^*(-\partial\mathcal{E}(s, u(s))) ds + \mathcal{E}(t, u(t)) = \mathcal{E}(0, u_0) + \int_0^t \partial_t \mathcal{E}(s, u(s)) ds$$

# An existence result

## To sum it up

- ▶ Fréchet subdifferential  $\partial_F \mathcal{E}(t, \cdot)$  naturally pops out from time-incremental minimization
- ▶ use with its strong-weak closure  $\partial \mathcal{E}(t, \cdot)$



## An existence result

### Existence theorem

Assume

- ▶  $u \mapsto \mathcal{E}(t, u)$  is **coercive**
- ▶  $t \mapsto \mathcal{E}(t, u)$  is **smooth** enough
- ▶  $\partial\mathcal{E}(t, \cdot)$  complies with the **chain rule**: for example if

$$\mathcal{E}(t, \cdot) = \mathcal{E}_{\text{convex}}(t, \cdot) + \mathcal{E}_{\text{concave}}(t, \cdot)$$

& the **convex part dominates concave part**

## An existence result

### Existence theorem

Assume

- ▶  $u \mapsto \mathcal{E}(t, u)$  is **coercive**
- ▶  $t \mapsto \mathcal{E}(t, u)$  is **smooth** enough
- ▶  $\partial\mathcal{E}(t, \cdot)$  complies with the **chain rule**: for example if

$$\mathcal{E}(t, \cdot) = \mathcal{E}_{\text{convex}}(t, \cdot) + \mathcal{E}_{\text{concave}}(t, \cdot)$$

& the **convex part dominates concave part**

Then, the approximate solutions converge to a curve  $u \in W^{1,1}(0, T; \mathcal{B})$  which **solves** the Cauchy problem for

$$\partial\Psi_\varepsilon(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 \quad \text{for a.a. } t \in (0, T).$$

## An existence result

### Existence theorem

Assume

- ▶  $u \mapsto \mathcal{E}(t, u)$  is **coercive**
- ▶  $t \mapsto \mathcal{E}(t, u)$  is **smooth** enough
- ▶  $\partial\mathcal{E}(t, \cdot)$  complies with the **chain rule**: for example if

$$\mathcal{E}(t, \cdot) = \mathcal{E}_{\text{convex}}(t, \cdot) + \mathcal{E}_{\text{concave}}(t, \cdot)$$

& the **convex part dominates concave part**

Then, the approximate solutions converge to a curve  $u \in W^{1,1}(0, T; \mathcal{B})$  which **solves** the Cauchy problem for

$$\partial\Psi_\varepsilon(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 \quad \text{for a.a. } t \in (0, T).$$

Moreover,  $u$  fulfils the **energy identity** (due to the chain rule) for all  $0 \leq s \leq t \leq T$

$$\int_s^t \Psi_\varepsilon(u'(r)) \, dr + \int_s^t \Psi_\varepsilon^*(-\partial\mathcal{E}(r, u(r))) \, dr + \mathcal{E}(t, u(t)) = \mathcal{E}(0, u_0) + \int_s^t \partial_t \mathcal{E}(r, u(r)) \, dr.$$

## Setting

Specialize to dissipation  $\Psi_\varepsilon$  on  $\mathcal{B}$

$$\Psi_\varepsilon(\cdot) = \overbrace{\Psi_1(\cdot)}^{\text{rate-independent}} + \varepsilon \overbrace{\Psi_2(\cdot)}^{\text{viscous}}$$

## Setting

Specialize to dissipation  $\Psi_\varepsilon$  on  $\mathcal{B}$

$$\Psi_\varepsilon(\cdot) = \overbrace{\Psi_1(\cdot)}^{\text{rate-independent}} + \varepsilon \overbrace{\Psi_2(\cdot)}^{\text{viscous}}$$

Further specialize to **dissipation given by norms!**

## Setting

Specialize to dissipation  $\Psi_\varepsilon$  on  $\mathcal{B}$

$$\Psi_\varepsilon(\cdot) = \overbrace{\Psi_1(\cdot)}^{\text{rate-independent}} + \varepsilon \overbrace{\Psi_2(\cdot)}^{\text{viscous}}$$

Further specialize to **dissipation given by norms!**

### Abstract rate-independent evolutions

- ▶ Dissipation:
  - ▶  $B_2$  reflexive
  - ▶  $B_2 \subset B_1 \equiv \mathcal{B}$
  - ▶ rate-independent dissipation:  $\Psi_1(\cdot) = |\cdot|_1$
- ▶ Energy:  $\mathcal{E} : (0, T) \times B_2 \rightarrow (-\infty, +\infty]$ ,  $\mathcal{E}(t, \cdot)$  with **compact sublevels** in  $B_2$ ,  $\mathcal{E}(t, \cdot)$  **nonconvex**, smooth with respect to  $t$

## Setting

Specialize to dissipation  $\Psi_\varepsilon$  on  $\mathcal{B}$

$$\Psi_\varepsilon(\cdot) = \overbrace{\Psi_1(\cdot)}^{\text{rate-independent}} + \varepsilon \overbrace{\Psi_2(\cdot)}^{\text{viscous}}$$

Further specialize to **dissipation given by norms!**

### Abstract rate-independent evolutions

- ▶ Dissipation:
  - ▶  $B_2$  reflexive
  - ▶  $B_2 \subset B_1 \equiv \mathcal{B}$
  - ▶ rate-independent dissipation:  $\Psi_1(\cdot) = |\cdot|_1$
- ▶ Energy:  $\mathcal{E} : (0, T) \times B_2 \rightarrow (-\infty, +\infty]$ ,  $\mathcal{E}(t, \cdot)$  with **compact sublevels** in  $B_2$ ,  $\mathcal{E}(t, \cdot)$  **nonconvex**, smooth with respect to  $t$

$$\partial\Psi_1(u'(t)) + \partial\mathcal{E}(t, u(t)) \ni 0 \quad \text{in } B_2', \quad t \in (0, T)$$

## Setting

Specialize to dissipation  $\Psi_\varepsilon$  on  $\mathcal{B}$

$$\Psi_\varepsilon(\cdot) = \overbrace{\Psi_1(\cdot)}^{\text{rate-independent}} + \varepsilon \overbrace{\Psi_2(\cdot)}^{\text{viscous}}$$

Further specialize to **dissipation given by norms!**

### Interaction of $L^1$ & $L^2$ norms

- ▶ Dissipation:
  - ▶  $B_2 \sim L^2(\Omega)$
  - ▶  $B_1 \sim L^1(\Omega)$
  - ▶ rate-independent dissipation:  $\Psi_1(\cdot) = |\cdot|_1$
- ▶ Energy:

$$\mathcal{E}(t, u) = \frac{1}{2} \int_{\Omega} |\nabla u|^2 + \overbrace{\int_{\Omega} \mathcal{W}(u)}^{\text{double-well potential}} - \langle \ell(t), u \rangle$$



## Setting

Specialize to dissipation  $\Psi_\varepsilon$  on  $\mathcal{B}$

$$\Psi_\varepsilon(\cdot) = \overbrace{\Psi_1(\cdot)}^{\text{rate-independent}} + \varepsilon \overbrace{\Psi_2(\cdot)}^{\text{viscous}}$$

Further specialize to **dissipation given by norms!**

### Interaction of $L^1$ & $L^2$ norms

► Dissipation:

- $B_2 \sim L^2(\Omega)$
- $B_1 \sim L^1(\Omega)$
- rate-independent dissipation:  $\Psi_1(\cdot) = |\cdot|_1$

► Energy:

$$\mathcal{E}(t, u) = \frac{1}{2} \int_{\Omega} |\nabla u|^2 + \overbrace{\int_{\Omega} \mathcal{W}(u)}^{\text{double-well potential}} - \langle \ell(t), u \rangle$$

$$\partial \Psi_1(u'(t)) + \partial \mathcal{E}(t, u(t)) \ni 0 \quad \text{in } B'_2, \quad t \in (0, T)$$

$$\text{Sign}(u_t) - \Delta u + \mathcal{W}'(u) = \ell(t) \quad \text{in } \Omega \times (0, T)$$

## Setting

Specialize to dissipation  $\Psi_\varepsilon$  on  $\mathcal{B}$

$$\Psi_\varepsilon(\cdot) = \underbrace{\Psi_1(\cdot)}_{\text{rate-independent}} + \varepsilon \underbrace{\Psi_2(\cdot)}_{\text{viscous}}$$

Further specialize to **dissipation given by norms!**

### Interaction of $L^1$ & $L^2$ norms

► Dissipation:

- $B_2 \sim L^2(\Omega)$
- $B_1 \sim L^1(\Omega)$
- rate-independent dissipation:  $\Psi_1(\cdot) = |\cdot|_1$
- **viscous approximation**  $\Psi_\varepsilon(\cdot) = |\cdot|_1 + \frac{\varepsilon}{2} \|\cdot\|_2^2$

► Energy:

$$\mathcal{E}(t, u) = \frac{1}{2} \int_{\Omega} |\nabla u|^2 + \int_{\Omega} \mathcal{W}(u) - \langle \ell(t), u \rangle$$

$$\varepsilon \partial \Psi_2(u'(t)) + \partial \Psi_1(u'(t)) + \partial \mathcal{E}(t, u(t)) \ni 0 \quad \text{in } B_2', \quad t \in (0, T)$$

$$\varepsilon u_t + \text{Sign}(u_t) - \Delta u + \mathcal{W}'(u) = \ell(t) \quad \text{in } \Omega \times (0, T)$$

## Step 1: energy identity

**Chain rule for  $\partial\mathcal{E}$  + convex analysis:** every solution of

$$\overbrace{\varepsilon \partial\Psi_2(u'(t)) + \partial\Psi_1(u'(t))}^{\partial\Psi_\varepsilon(u'(t))} + \partial\mathcal{E}(t, u(t)) \ni 0 \text{ in } B'_2, \quad t \in (0, T)$$

fulfils **energy identity**  $\forall 0 \leq t_1 \leq t_2 \leq T$

$$\begin{aligned} \int_{t_1}^{t_2} \Psi_\varepsilon(\dot{u}(r)) \, dr + \int_{t_1}^{t_2} \Psi_\varepsilon^*(-\partial\mathcal{E}(r, u(r))) \, dr + \mathcal{E}(t_2, u(t_2)) \\ = \mathcal{E}(t_1, u(t_1)) + \int_{t_1}^{t_2} \partial_t \mathcal{E}(t, u(r)) \, dr \end{aligned}$$

## Step 1: energy identity

**Chain rule for  $\partial\mathcal{E}$  + convex analysis:** every solution of

$$\overbrace{\varepsilon \partial\Psi_2(u'(t)) + \partial\Psi_1(u'(t))}^{\partial\Psi_\varepsilon(u'(t))} + \partial\mathcal{E}(t, u(t)) \ni 0 \text{ in } B'_1, \quad t \in (0, T)$$

fulfils **energy identity**  $\forall 0 \leq t_1 \leq t_2 \leq T$

$$\begin{aligned} \int_{t_1}^{t_2} \Psi_\varepsilon(\dot{u}(r)) \, dr + \int_{t_1}^{t_2} \Psi_\varepsilon^*(-\partial\mathcal{E}(r, u(r))) \, dr + \mathcal{E}(t_2, u(t_2)) \\ = \mathcal{E}(t_1, u(t_1)) + \int_{t_1}^{t_2} \partial_t \mathcal{E}(t, u(r)) \, dr \end{aligned}$$

where the conjugate  $\Psi_\varepsilon^*$  is:

$$\Psi_\varepsilon^*(-\partial\mathcal{E}) = \frac{1}{2\varepsilon} \min_{|z|_{1,*} \leq 1} \| -\partial\mathcal{E} - z \|_{2,*}^2 = \frac{1}{2\varepsilon} d_2(-\partial\mathcal{E}, K^*)^2$$

and  $K^*$  **unitary ball in  $B_1^*$** .

## Competition between viscous effects & rate-independent behaviour

$$\begin{aligned} & \int_{t_1}^{t_2} |u'(r)|_1 \, dr + \int_{t_1}^{t_2} \frac{\varepsilon}{2} \|u'(r)\|_2^2 + \frac{1}{2\varepsilon} d_2(-\partial\mathcal{E}(r, u(r)), K^*)^2 \, dr + \mathcal{E}(t_2, u(t_2)) \\ & = \mathcal{E}(t_1, u(t_1)) + \int_{t_1}^{t_2} \partial_t \mathcal{E}(t, u(r)) \, dr \end{aligned}$$

## Competition between viscous effects & rate-independent behaviour

$$\begin{aligned} & \int_{t_1}^{t_2} |u'(r)|_1 \, dr + \int_{t_1}^{t_2} \frac{\varepsilon}{2} \|u'(r)\|_2^2 + \frac{1}{2\varepsilon} d_2(-\partial\mathcal{E}(r, u(r)), K^*)^2 \, dr + \mathcal{E}(t_2, u(t_2)) \\ & = \mathcal{E}(t_1, u(t_1)) + \int_{t_1}^{t_2} \partial_t \mathcal{E}(t, u(r)) \, dr \end{aligned}$$

♣ Note that

$$d_2(-\partial\mathcal{E}(r, u(r)), K^*) = 0 \quad \Leftrightarrow \quad |-\partial\mathcal{E}(r, u(r))|_{1,*} \leq 1$$

**LOCAL version of the stability condition:**

$$|-\partial\mathcal{E}(r, u(r))|_{1,*} \leq 1 \quad \text{vs.} \quad \frac{\mathcal{E}(t, u(t)) - \mathcal{E}(t, z)}{\mathcal{D}(u(t), z)} \leq 1 \quad \forall z$$

## Competition between viscous effects & rate-independent behaviour

$$\begin{aligned} & \int_{t_1}^{t_2} |u'(r)|_1 \, dr + \int_{t_1}^{t_2} \frac{\varepsilon}{2} \|u'(r)\|_2^2 + \frac{1}{2\varepsilon} d_2(-\partial\mathcal{E}(r, u(r)), K^*)^2 \, dr + \mathcal{E}(t_2, u(t_2)) \\ & = \mathcal{E}(t_1, u(t_1)) + \int_{t_1}^{t_2} \partial_t \mathcal{E}(t, u(r)) \, dr \end{aligned}$$

♣ Note that

$$d_2(-\partial\mathcal{E}(r, u(r)), K^*) = 0 \quad \Leftrightarrow \quad |-\partial\mathcal{E}(r, u(r))|_{1,*} \leq 1$$

**LOCAL version of the stability condition:**

$$|-\partial\mathcal{E}(r, u(r))|_{1,*} \leq 1 \quad \text{vs.} \quad \frac{\mathcal{E}(t, u(t)) - \mathcal{E}(t, z)}{\mathcal{D}(u(t), z)} \leq 1 \quad \forall z$$

♣ Energy identity highlights the **competition between viscosity & rate-independence**

## Step 2: rescaling

### The technique by Efendiev-Mielke

- ▶ Jumps in the vanishing viscosity limit correspond to **viscous transitions** between stable states
- ▶ To capture the viscous transition path: **NOT SHRINK** jumps at a point, look at curves with their **arclength parametrization**
- ▶ Asymptotic analysis of (reparametrized) trajectories in an extended phase space



## Step 2: rescaling

### The technique by Efendiev-Mielke

- ▶ Jumps in the vanishing viscosity limit correspond to **viscous transitions** between stable states
- ▶ To capture the viscous transition path: **NOT SHRINK** jumps at a point, look at curves with their **arclength parametrization**
- ▶ Asymptotic analysis of (reparametrized) trajectories in an extended phase space

We reparametrize

$$\begin{aligned} \int_{t_1}^{t_2} |u'(r)|_1 dr + \int_{t_1}^{t_2} \frac{\varepsilon}{2} \|u'(r)\|_2^2 + \frac{1}{2\varepsilon} d_2(-\partial\mathcal{E}(r, u(r)), K^*)^2 dr + \mathcal{E}(t_2, u(t_2)) \\ = \mathcal{E}(t_1, u(t_1)) + \int_{t_1}^{t_2} \partial_t \mathcal{E}(t, u(r)) dr \end{aligned}$$

by the rescaling function:

$$s_\varepsilon(t) = t + \int_0^t (|u'(r)|_1 + \|u'(r)\|_2 \cdot d_2(-\partial\mathcal{E}(r, u(r)), K^*)) dr$$

**"energy arclength"**

## Step 2: rescaling

$$\begin{cases} s_\varepsilon(t) := t + \int_0^t (|u'(r)|_1 + \|u'(r)\|_2 \cdot d_2(-\partial\mathcal{E}(r, u(r)), K^*)) \, dr \\ \hat{t}_\varepsilon = s_\varepsilon^{-1}, \quad \hat{u}_\varepsilon = u \circ \hat{t}_\varepsilon \end{cases}$$

Hence we perform the asymptotic analysis for the **"extended" trajectory**  $\{(\hat{t}_\varepsilon, \hat{u}_\varepsilon)\}$

## Step 2: rescaling

$$\begin{cases} s_\varepsilon(t) := t + \int_0^t (|u'(r)|_1 + \|u'(r)\|_2 \cdot d_2(-\partial\mathcal{E}(r, u(r)), K^*)) \, dr \\ \hat{t}_\varepsilon = s_\varepsilon^{-1}, \quad \hat{u}_\varepsilon = u \circ \hat{t}_\varepsilon \end{cases}$$

Hence we perform the asymptotic analysis for the **"extended" trajectory**  $\{(\hat{t}_\varepsilon, \hat{u}_\varepsilon)\}$

### Rescaled energy identity

$$\begin{aligned} \int_{s_1}^{s_2} |\hat{u}'_\varepsilon|_1 + \int_{s_1}^{s_2} \frac{\varepsilon}{2\hat{t}'_\varepsilon} \|\hat{u}'_\varepsilon\|_2^2 + \frac{\hat{t}'_\varepsilon}{2\varepsilon} d_2(-\partial\mathcal{E}(\hat{t}_\varepsilon, \hat{u}_\varepsilon), K^*)^2 + \mathcal{E}(\hat{t}_\varepsilon(s_2), \hat{u}_\varepsilon(s_2)) \\ = \mathcal{E}(\hat{t}_\varepsilon(s_1), \hat{u}_\varepsilon(s_1)) + \int_{s_1}^{s_2} \partial_t \mathcal{E}(\hat{t}_\varepsilon, \hat{u}_\varepsilon) \hat{t}'_\varepsilon \end{aligned}$$

+ **"normalization" condition**

$$\hat{t}'_\varepsilon + |\hat{u}'_\varepsilon|_1 + \|\hat{u}'_\varepsilon\|_2 \cdot d_2(-\partial\mathcal{E}(\hat{t}_\varepsilon, \hat{u}_\varepsilon), K^*) \equiv 1$$

## Step 2: rescaling

$$\begin{cases} s_\varepsilon(t) := t + \int_0^t (|u'(r)|_1 + \|u'(r)\|_2 \cdot d_2(-\partial\mathcal{E}(r, u(r)), K^*)) dr \\ \hat{t}_\varepsilon = s_\varepsilon^{-1}, \quad \hat{u}_\varepsilon = u \circ \hat{t}_\varepsilon \end{cases}$$

Hence we perform the asymptotic analysis for the **"extended" trajectory**  $\{(\hat{t}_\varepsilon, \hat{u}_\varepsilon)\}$

### Rescaled energy identity

$$\begin{aligned} \int_{s_1}^{s_2} |\hat{u}'_\varepsilon|_1 + \int_{s_1}^{s_2} \frac{\varepsilon}{2\hat{t}'_\varepsilon} \|\hat{u}'_\varepsilon\|_2^2 + \frac{\hat{t}'_\varepsilon}{2\varepsilon} d_2(-\partial\mathcal{E}(\hat{t}_\varepsilon, \hat{u}_\varepsilon), K^*)^2 + \mathcal{E}(\hat{t}_\varepsilon(s_2), \hat{u}_\varepsilon(s_2)) \\ = \mathcal{E}(\hat{t}_\varepsilon(s_1), \hat{u}_\varepsilon(s_1)) + \int_{s_1}^{s_2} \partial_t \mathcal{E}(\hat{t}_\varepsilon, \hat{u}_\varepsilon) \hat{t}'_\varepsilon \end{aligned}$$

+ **"normalization" condition**

$$\hat{t}'_\varepsilon + |\hat{u}'_\varepsilon|_1 + \|\hat{u}'_\varepsilon\|_2 \cdot d_2(-\partial\mathcal{E}(\hat{t}_\varepsilon, \hat{u}_\varepsilon), K^*) \equiv 1$$

$\Rightarrow$  **A priori estimates, compactness** for  $\{(\hat{t}_\varepsilon, \hat{u}_\varepsilon)\}$

## The asymptotic analysis as $\varepsilon \downarrow 0$

### Theorem [Mielke, R., Savaré'09]

- ▶  $B_2$  reflexive,  $B_2 \subset B_1$
- ▶ Energy:  $\mathcal{E} : (0, T) \times B_2 \rightarrow (-\infty, +\infty]$ , with compact sublevels in  $B_2$ ,
- ▶  $\mathcal{E}$  complies with **chain rule**

## The asymptotic analysis as $\varepsilon \downarrow 0$

### Theorem [Mielke, R., Savaré'09]

Up to a subsequence, as  $\varepsilon \searrow 0$  the trajectory  $\{(\hat{t}_\varepsilon, \hat{u}_\varepsilon)\}$

$$\begin{aligned} \int_{s_1}^{s_2} |\hat{u}'_\varepsilon|_1 + \int_{s_1}^{s_2} \frac{\varepsilon}{2\hat{t}'_\varepsilon} \|\hat{u}'_\varepsilon\|_2^2 + \frac{\hat{t}'_\varepsilon}{2\varepsilon} d_2(-\partial\mathcal{E}(\hat{t}_\varepsilon, \hat{u}_\varepsilon), K^*)^2 + \mathcal{E}(\hat{t}_\varepsilon(s_2), \hat{u}_\varepsilon(s_2)) \\ = \mathcal{E}(\hat{t}_\varepsilon(s_1), \hat{u}_\varepsilon(s_1)) + \int_{s_1}^{s_2} \partial_t \mathcal{E}(\hat{t}_\varepsilon, \hat{u}_\varepsilon) \hat{t}'_\varepsilon \end{aligned}$$

## The asymptotic analysis as $\varepsilon \downarrow 0$

### Theorem [Mielke, R., Savaré'09]

Up to a subsequence, as  $\varepsilon \searrow 0$  the trajectory  $\{(\hat{t}_\varepsilon, \hat{u}_\varepsilon)\}$

$$\begin{aligned} \int_{s_1}^{s_2} |\hat{u}'_\varepsilon|_1 + \int_{s_1}^{s_2} \frac{\varepsilon}{2\hat{t}'_\varepsilon} \|\hat{u}'_\varepsilon\|_2^2 + \frac{\hat{t}'_\varepsilon}{2\varepsilon} d_2(-\partial\mathcal{E}(\hat{t}_\varepsilon, \hat{u}_\varepsilon), K^*)^2 + \mathcal{E}(\hat{t}_\varepsilon(s_2), \hat{u}_\varepsilon(s_2)) \\ = \mathcal{E}(\hat{t}_\varepsilon(s_1), \hat{u}_\varepsilon(s_1)) + \int_{s_1}^{s_2} \partial_t \mathcal{E}(\hat{t}_\varepsilon, \hat{u}_\varepsilon) \hat{t}'_\varepsilon \end{aligned}$$

converges to  $(\hat{t}, \hat{u}) \in C_{\text{Lip}}^0([0, \hat{T}]; [0, T] \times B_1)$

$$\begin{aligned} \int_{s_1}^{s_2} |\hat{u}'|_1 + \int_{s_1}^{s_2} \mathcal{M}_0(\hat{t}', \|\hat{u}'\|_2, -\partial\mathcal{E}(\hat{t}, \hat{u})) + \mathcal{E}(\hat{t}(s_2), \hat{u}(s_2)) \\ = \mathcal{E}(\hat{t}(s_1), \hat{u}(s_1)) + \int_{s_1}^{s_2} \partial_t \mathcal{E}(\hat{t}, \hat{u}) \hat{t}' \end{aligned}$$

( $|\hat{u}'|_1$  **metric derivative** of  $\hat{u}$ !), with

$$\mathcal{M}_0(\hat{t}', \|\hat{u}'\|_2, -\partial\mathcal{E}(\hat{t}, \hat{u})) = \begin{cases} \|\hat{u}'\|_2 \cdot d_2(-\partial\mathcal{E}(\hat{t}, \hat{u}), K^*) & \text{se } \hat{t}' = 0 \\ I_{\{0\}}(d_2(-\partial\mathcal{E}(\hat{t}, \hat{u}), K^*)) & \text{se } \hat{t}' > 0 \end{cases}$$

## The asymptotic analysis as $\varepsilon \downarrow 0$

### Theorem [Mielke, R., Savaré'09]

Up to a subsequence, as  $\varepsilon \searrow 0$  the trajectory  $\{(\hat{t}_\varepsilon, \hat{u}_\varepsilon)\}$

$$\begin{aligned} \int_{s_1}^{s_2} |\hat{u}'_\varepsilon|_1 + \int_{s_1}^{s_2} \frac{\varepsilon}{2\hat{t}'_\varepsilon} \|\hat{u}'_\varepsilon\|_2^2 + \frac{\hat{t}'_\varepsilon}{2\varepsilon} d_2(-\partial\mathcal{E}(\hat{t}_\varepsilon, \hat{u}_\varepsilon), K^*)^2 + \mathcal{E}(\hat{t}_\varepsilon(s_2), \hat{u}_\varepsilon(s_2)) \\ = \mathcal{E}(\hat{t}_\varepsilon(s_1), \hat{u}_\varepsilon(s_1)) + \int_{s_1}^{s_2} \partial_t \mathcal{E}(\hat{t}_\varepsilon, \hat{u}_\varepsilon) \hat{t}'_\varepsilon \end{aligned}$$

converges to  $(\hat{t}, \hat{u}) \in C_{\text{Lip}}^0([0, \hat{T}]; [0, T] \times B_1)$

$$\begin{aligned} \int_{s_1}^{s_2} |\hat{u}'|_1 + \int_{s_1}^{s_2} \mathcal{M}_0(\hat{t}', \|\hat{u}'\|_2, -\partial\mathcal{E}(\hat{t}, \hat{u})) + \mathcal{E}(\hat{t}(s_2), \hat{u}(s_2)) \\ = \mathcal{E}(\hat{t}(s_1), \hat{u}(s_1)) + \int_{s_1}^{s_2} \partial_t \mathcal{E}(\hat{t}, \hat{u}) \hat{t}' \end{aligned}$$

( $|\hat{u}'|_1$  **metric derivative** of  $\hat{u}$ !), with

$$\mathcal{M}_0(\hat{t}', \|\hat{u}'\|_2, -\partial\mathcal{E}(\hat{t}, \hat{u})) = \begin{cases} \|\hat{u}'\|_2 \cdot d_2(-\partial\mathcal{E}(\hat{t}, \hat{u}), K^*) & \text{se } \hat{t}' = 0 \\ I_{\{0\}}(d_2(-\partial\mathcal{E}(\hat{t}, \hat{u}), K^*)) & \text{se } \hat{t}' > 0 \end{cases}$$

+ **"normalization condition"**

$$\hat{t}' + |\hat{u}'|_1 + \|\hat{u}'\|_2 \cdot d_2(-\partial\mathcal{E}(\hat{t}, \hat{u}), K^*) \equiv 1$$



## Two key ideas

### Proof based on:

- ▶ reparametrization of trajectories
- ▶ energy identity: need for a notion of subdifferential for which the chain rule holds..

## Remarks on the limit problem

$$(\hat{t}, \hat{u}) \in C_{\text{Lip}}^0([0, \hat{T}]; [0, T] \times B_1), \quad \hat{t}(0) = 0, \quad \hat{t}(\hat{T}) = T$$

**Normalization:**  $\hat{t}' + |\hat{u}'|_1 + \|\hat{u}'\|_2 \cdot d_2(-\partial\mathcal{E}(\hat{t}, \hat{u}), K^*) \equiv 1$

**Energy id.:** 
$$\begin{aligned} \frac{d}{ds} \mathcal{E}(\hat{t}(s), \hat{u}(s)) - \partial_t \mathcal{E}(\hat{t}(s), \hat{u}(s)) \hat{t}'(s) \\ = -\langle -\partial\mathcal{E}(\hat{t}(s), \hat{u}(s)), \hat{u}'(s) \rangle \\ = -|\hat{u}'(s)|_1 - \|\hat{u}'(s)\|_2 \cdot d_2(-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s)), K^*) \quad \text{for a.a. } s \in (0, \hat{T}) \end{aligned}$$

**Constraint:**  $\mathcal{M}_0(\hat{t}', \|\hat{u}'\|_2, -\partial\mathcal{E}(\hat{t}, \hat{u})) \equiv \|\hat{u}'\|_2 \cdot d_2(-\partial\mathcal{E}(\hat{t}, \hat{u}), K^*)$

## Remarks on the limit problem

$$(\hat{t}, \hat{u}) \in C_{\text{Lip}}^0([0, \hat{T}]; [0, T] \times B_1), \quad \hat{t}(0) = 0, \quad \hat{t}(\hat{T}) = T$$

**Normalization:**  $\hat{t}' + |\hat{u}'|_1 + \|\hat{u}'\|_2 \cdot d_2(-\partial\mathcal{E}(\hat{t}, \hat{u}), K^*) \equiv 1$

**Energy id.:**  $\frac{d}{ds} \mathcal{E}(\hat{t}(s), \hat{u}(s)) - \partial_t \mathcal{E}(\hat{t}(s), \hat{u}(s)) \hat{t}'(s)$

$$= -\langle -\partial\mathcal{E}(\hat{t}(s), \hat{u}(s)), \hat{u}'(s) \rangle$$

$$= -|\hat{u}'(s)|_1 - \|\hat{u}'(s)\|_2 \cdot d_2(-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s)), K^*) \quad \text{for a.a. } s \in (0, \hat{T})$$

**Three regimes:**

$$\left\{ \begin{array}{l} \hat{t}'(s) = 1 \quad (\Leftrightarrow \hat{u}'(s) = 0) \\ \quad \Rightarrow \quad |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \leq 1 \\ \hat{t}'(s) \in (0, 1) \quad (\Leftrightarrow |\hat{u}'(s)|_1 \in (0, 1)) \\ \quad \Rightarrow \quad |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} = 1 \\ \hat{t}'(s) = 0 \quad (\Leftrightarrow |\hat{u}'(s)|_1 + \|\hat{u}'(s)\|_2 \cdot d_2(-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s)), K^*) = 1) \\ \quad \Rightarrow \quad |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \geq 1 \end{array} \right.$$

## Parametrized rate-independent evolutions

### Definition

We call **parametrized rate-independent evolution** a pair

$$(\hat{t}, \hat{u}) \in \text{AC}([0, \hat{T}]; [0, T] \times B_1), \quad \hat{t}(0) = 0, \quad \hat{t}(\hat{T}) = T,$$

satisfying

**Nondegeneracy:**  $\hat{t}'(s) + |\hat{u}'(s)|_1 > 0$  for a.a.  $s \in (0, \hat{T})$ ,

**Energy id.:** 
$$\begin{aligned} \frac{d}{ds} \mathcal{E}(\hat{t}(s), \hat{u}(s)) - \partial_t \mathcal{E}(\hat{t}(s), \hat{u}(s)) \hat{t}'(s) \\ = -\langle -\partial \mathcal{E}(\hat{t}(s), \hat{u}(s)), \hat{u}'(s) \rangle \\ = -|\hat{u}'(s)|_1 - \|\hat{u}'(s)\|_2 \cdot d_2(-\partial \mathcal{E}(\hat{t}(s), \hat{u}(s)), K^*) \quad \text{for a.a. } s \in (0, \hat{T}) \end{aligned}$$

**Three regimes:** 
$$\begin{cases} \hat{t}'(s) > 0 \Rightarrow |-\partial \mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \leq 1 \\ \hat{t}'(s) |\hat{u}'(s)|_1 > 0 \Rightarrow |-\partial \mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} = 1 \\ \hat{t}'(s) = 0 \Rightarrow |-\partial \mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \geq 1 \end{cases}$$

## Properties of parametrized rate-independent evolutions

**Nondegeneracy:**  $\hat{t}'(s) + |\hat{u}'(s)|_1 > 0$  for a.a.  $s \in (0, \hat{T})$ ,

**Energy id.:** 
$$\begin{aligned} \frac{d}{ds} \mathcal{E}(\hat{t}(s), \hat{u}(s)) - \partial_t \mathcal{E}(\hat{t}(s), \hat{u}(s)) \hat{t}'(s) \\ = -\langle -\partial \mathcal{E}(\hat{t}(s), \hat{u}(s)), \hat{u}'(s) \rangle \\ = -|\hat{u}'(s)|_1 - \|\hat{u}'(s)\|_2 \cdot d_2(-\partial \mathcal{E}(\hat{t}(s), \hat{u}(s)), K^*) \quad \text{for a.a. } s \in (0, \hat{T}) \end{aligned}$$

**Three regimes:** 
$$\begin{cases} \hat{t}'(s) > 0 \Rightarrow & |-\partial \mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \leq 1 \\ \hat{t}'(s) |\hat{u}'(s)|_1 > 0 \Rightarrow & |-\partial \mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} = 1 \\ \hat{t}'(s) = 0 \Rightarrow & |-\partial \mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \geq 1 \end{cases}$$

- ▶ **Intrinsically rate-independent** notion: invariant for time-rescalings
- ▶ Arises in the **vanishing viscosity limit**
- ▶ Under conditions on  $B_1$ , **notion equivalent to the doubly nonlinear PDE formulation**
- ▶ **Stability w.r.t. the problem data**

## Properties of parametrized rate-independent evolutions

**Nondegeneracy:**  $\hat{t}'(s) + |\hat{u}'(s)|_1 > 0$  for a.a.  $s \in (0, \hat{T})$ ,

**Energy id.:** 
$$\begin{aligned} \frac{d}{ds} \mathcal{E}(\hat{t}(s), \hat{u}(s)) - \partial_t \mathcal{E}(\hat{t}(s), \hat{u}(s)) \hat{t}'(s) \\ = -\langle -\partial \mathcal{E}(\hat{t}(s), \hat{u}(s)), \hat{u}'(s) \rangle \\ = -|\hat{u}'(s)|_1 - \|\hat{u}'(s)\|_2 \cdot d_2(-\partial \mathcal{E}(\hat{t}(s), \hat{u}(s)), K^*) \quad \text{for a.a. } s \in (0, \hat{T}) \end{aligned}$$

**Three regimes:** 
$$\begin{cases} \hat{t}'(s) > 0 \Rightarrow & |-\partial \mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \leq 1 \\ \hat{t}'(s) |\hat{u}'(s)|_1 > 0 \Rightarrow & |-\partial \mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} = 1 \\ \hat{t}'(s) = 0 \Rightarrow & |-\partial \mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \geq 1 \end{cases}$$

- ▶ **Intrinsically rate-independent** notion: invariant for time-rescalings
- ▶ Arises in the **vanishing viscosity limit**
- ▶ Under conditions on  $B_1$ , **notion equivalent to the doubly nonlinear PDE formulation**
- ▶ **Stability w.r.t. the problem data**
- ▶ **Three regimes reflect the different types of evolution of the system**

## The three regimes

$$\begin{cases} \hat{t}'(s) > 0, |\hat{u}'(s)|_1 = 0 \Rightarrow & |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \leq 1 & \text{(sticking)} \\ \hat{t}'(s) |\hat{u}'(s)|_1 > 0 \Rightarrow & |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} = 1 & \text{(sliding)} \\ \hat{t}'(s) = 0 \Rightarrow & |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \geq 1 & \text{(viscous slip)} \end{cases}$$

## The three regimes

$$\begin{cases} \hat{t}'(s) > 0, |\hat{u}'(s)|_1 = 0 \Rightarrow & |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \leq 1 & \text{(stationarity)} \\ \hat{t}'(s) |\hat{u}'(s)|_1 > 0 \Rightarrow & |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} = 1 & \text{(rate-independent)} \\ \hat{t}'(s) = 0 \Rightarrow & |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \geq 1 & \text{(jump)} \end{cases}$$

### Stationarity

- ▶ **local stability** condition

$$|-\partial\mathcal{E}(\hat{t}(s_0), \hat{u}(s_0))|_{1,*} \leq 1 \quad \text{vs.} \quad \frac{\mathcal{E}(s_0, u(s_0)) - \mathcal{E}(s_0, z)}{\mathcal{D}(u(s_0), z)} \leq 1 \quad \forall z$$

- ▶ in a neighbourhood  $I(s_0)$  there hold  $\hat{u}(s) \equiv \hat{u}(s_0)$  and the **energy identity**

$$\mathcal{E}(\hat{t}(s_2), \hat{u}(s_0)) - \mathcal{E}(\hat{t}(s_1), \hat{u}(s_0)) = \int_{s_1}^{s_2} \partial_t \mathcal{E}(\hat{t}(s), \hat{u}(s_0)) \hat{t}'(s) \, ds \quad \forall s_1 \leq s_2 \in I(s_0).$$



## The three regimes

$$\begin{cases} \hat{t}'(s) > 0, |\hat{u}'(s)|_1 = 0 \Rightarrow & |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \leq 1 & \text{(stationarity)} \\ \hat{t}'(s) |\hat{u}'(s)|_1 > 0 \Rightarrow & |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} = 1 & \text{(rate-independent)} \\ \hat{t}'(s) = 0 \Rightarrow & |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \geq 1 & \text{(jump)} \end{cases}$$

### Rate-independent evolution

- ▶ **local stability** condition

$$|-\partial\mathcal{E}(\hat{t}(s_0), \hat{u}(s_0))|_{1,*} = 1 \quad \sim \quad -\partial\mathcal{E}(\hat{t}(s_0), \hat{u}(s_0)) \in \overbrace{\partial\Psi_1(\hat{u}'(s_0))}^{\text{Sign}(\hat{u}'(s_0))}$$

- ▶ in a neighbourhood  $I(s_0)$  **energy identity**  $\forall s_1 \leq s_2 \in I(s_0)$

$$\mathcal{E}(\hat{t}(s_2), \hat{u}(s_2)) - \mathcal{E}(\hat{t}(s_1), \hat{u}(s_1)) = \int_{s_1}^{s_2} (\partial_t \mathcal{E}(\hat{t}(s), \hat{u}(s)) \hat{t}'(s) - |\hat{u}'(s)|_1) \, ds.$$

## The three regimes

$$\left\{ \begin{array}{l} \hat{t}'(s) > 0, |\hat{u}'(s)|_1 = 0 \Rightarrow \\ \hat{t}'(s) |\hat{u}'(s)|_1 > 0 \Rightarrow \\ \hat{t}'(s) = 0 \Rightarrow \end{array} \right. \begin{array}{l} |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \leq 1 \quad \text{(stationarity)} \\ |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} = 1 \quad \text{(rate-independent)} \\ |-\partial\mathcal{E}(\hat{t}(s), \hat{u}(s))|_{1,*} \geq 1 \quad \text{(jump)} \end{array}$$

### Jump regime

- ▶ in a neighbourhood  $I(s_0)$  there hold  $\hat{t}(s) \equiv \hat{t}(s_0)$  and **energy identity**  
 $\forall s_1 \leq s_2 \in I(s_0)$

$$\begin{aligned} & \mathcal{E}(\hat{t}(s_0), \hat{u}(s_2)) - \mathcal{E}(\hat{t}(s_0), \hat{u}(s_1)) \\ &= \int_{s_1}^{s_2} (-|\hat{u}'(s)|_1 - \|\hat{u}'(s)\|_2 \cdot d_2(-\partial\mathcal{E}(\hat{t}(s_0), \hat{u}(s)), K^*)) \, ds. \end{aligned}$$

- ▶ With a **rescaling** which depends on the

$$\text{viscous quantities} \quad \|\hat{u}'\|_2 \ \& \ d_2(-\partial\mathcal{E}(\hat{t}, \hat{u}), K^*)$$

we pass from  $\hat{u}(s)$  to  $\tilde{u}(\sigma)$  solution of the **viscous doubly nonlinear equation**

$$\partial\Psi_1(\tilde{u}'(\sigma)) + \partial\Psi_2(\tilde{u}'(\sigma)) + \partial\mathcal{E}(\hat{t}(s_0), u(\sigma)) \ni 0$$

## Differential characterization

In the case

$B_1$  has the Radon Nikodým property

(hence we have **pointwise derivatives!!** But  $L^1$  is forbidden) &

## Differential characterization

In the case

$B_1$  has the Radon Nikodým property

(hence we have **pointwise derivatives!!** But  $L^1$  is forbidden) &

$$\mathcal{E}(t, u) = E(u) - \langle \ell(t), u \rangle$$

## Differential characterization

In the case

$B_1$  has the Radon Nikodým property

(hence we have **pointwise derivatives!!** But  $L^1$  is forbidden) &

$$\mathcal{E}(t, u) = E(u) - \langle \ell(t), u \rangle$$

then  $(\hat{t}, \hat{u}) \in \text{AC}([0, \hat{T}]; [0, T] \times B_1)$  is a parametrized rate-independent evolution

**if and only if**

$$\begin{aligned} &\exists \lambda : (0, \hat{T}) \rightarrow \mathbb{R}^+ : \text{for a.a. } s \in (0, \hat{T}) \\ &\left\{ \begin{array}{l} \partial\Psi_1(\hat{u}'(s)) + \lambda(s)\partial\Psi_2(\hat{u}'(s)) + \partial E(\hat{u}(s)) \ni \ell(\hat{t}(s)) \\ \lambda(s)\hat{t}'(s) = 0 \end{array} \right. \end{aligned}$$

## Differential characterization

In the case

$B_1$  has the Radon Nikodým property

(hence we have **pointwise derivatives!!** But  $L^1$  is forbidden) &

$$\mathcal{E}(t, u) = E(u) - \langle \ell(t), u \rangle$$

then  $(\hat{t}, \hat{u}) \in \text{AC}([0, \hat{T}]; [0, T] \times B_1)$  is a parametrized rate-independent evolution

**if and only if**

$$\begin{cases} \exists \lambda : (0, \hat{T}) \rightarrow \mathbb{R}^+ : \text{for a.a. } s \in (0, \hat{T}) \\ \left\{ \begin{array}{l} \partial\Psi_1(\hat{u}'(s)) + \lambda(s)\partial\Psi_2(\hat{u}'(s)) + \partial E(\hat{u}(s)) \ni \ell(\hat{t}(s)) \\ \lambda(s)\hat{t}'(s) = 0 \end{array} \right. \end{cases}$$

The "Lagrange multiplier"  $\lambda$  (and thus viscous dissipation) is **activated by**  $\hat{t}'(s) = 0$ , i.e. at jumps

## BV rate-independent evolutions

### From virtual to real jumps

With a suitable rescaling, we pass

from  $(\hat{t}, \hat{u})$  parametrized rate-independent evolution

↓ ↑

to  $u$  **BV rate-independent evolution**

in jump points it follows the trajectory of a **viscous doubly nonlinear equation**

## BV rate-independent evolutions

### From virtual to real jumps

With a suitable rescaling, we pass

from  $(\hat{t}, \hat{u})$  parametrized rate-independent evolution

$\Downarrow \Uparrow$

to  $u$  **BV rate-independent evolution**

in jump points it follows the trajectory of a **viscous doubly nonlinear equation**

### Approximation by time discretization

Existence of BV rate-independent evolutions: passing to the limit in the **discretization scheme**:

$$\left\{ \begin{array}{l} U_\tau^n \in \operatorname{Argmin}_{U \in B_2} \left\{ |U - U_\tau^{n-1}|_1 + \frac{\varepsilon(\tau)}{\tau} \|U - U_\tau^{n-1}\|_2^2 + \mathcal{E}(t_n, U) \right\} . \\ \text{with } \varepsilon(\tau) \rightarrow 0 \text{ \& } \frac{\varepsilon(\tau)}{\tau} \uparrow \infty \text{ as } \tau \downarrow 0 \end{array} \right.$$



## Metric analysis popping out

$$\begin{aligned} \int_{s_1}^{s_2} |\hat{u}'|_1 + \int_{s_1}^{s_2} \mathcal{M}_0(\hat{t}', \|\hat{u}'\|_2, -\partial\mathcal{E}(\hat{t}, \hat{u})) + \mathcal{E}(\hat{t}(s_2), \hat{u}(s_2)) \\ = \mathcal{E}(\hat{t}(s_1), \hat{u}(s_1)) + \int_{s_1}^{s_2} \partial_t \mathcal{E}(\hat{t}, \hat{u}) \hat{t}' \end{aligned}$$

**BUT:**

$B_1 \sim L^1$  does **not** have the **Radon-Nikodým property**

## Metric analysis popping out

$$\begin{aligned} & \int_{s_1}^{s_2} |\hat{u}'|_1 + \int_{s_1}^{s_2} \mathcal{M}_0(\hat{t}', \|\hat{u}'\|_2, -\partial\mathcal{E}(\hat{t}, \hat{u})) + \mathcal{E}(\hat{t}(s_2), \hat{u}(s_2)) \\ &= \mathcal{E}(\hat{t}(s_1), \hat{u}(s_1)) + \int_{s_1}^{s_2} \partial_t \mathcal{E}(\hat{t}, \hat{u}) \hat{t}' \end{aligned}$$

**BUT:**

$B_1 \sim L^1$  does **not** have the **Radon-Nikodým property**

- ▶ Need to replace pointwise derivative  $|\hat{u}'|_1$  by **metric derivative**
- ▶ Metric analysis, theory of **gradient flows** in metric spaces:
  - ▶ **De Giorgi, Marino, Saccon, Tosques, Degiovanni, Ambrosio '80 ~ '90**  $\rightsquigarrow$  theory of **Curves of Maximal Slope** and **Minimizing Movements**

## Metric analysis popping out

$$\begin{aligned} & \int_{s_1}^{s_2} |\hat{u}'|_1 + \int_{s_1}^{s_2} \mathcal{M}_0(\hat{t}', \|\hat{u}'\|_2, -\partial\mathcal{E}(\hat{t}, \hat{u})) + \mathcal{E}(\hat{t}(s_2), \hat{u}(s_2)) \\ & = \mathcal{E}(\hat{t}(s_1), \hat{u}(s_1)) + \int_{s_1}^{s_2} \partial_t \mathcal{E}(\hat{t}, \hat{u}) \hat{t}' \end{aligned}$$

**BUT:**

$B_1 \sim L^1$  does **not** have the **Radon-Nikodým property**

- ▶ Need to replace pointwise derivative  $|\hat{u}'|_1$  by **metric derivative**
- ▶ Metric analysis, theory of **gradient flows** in metric spaces:
  - ▶ **De Giorgi, Marino, Saccon, Tosques, Degiovanni, Ambrosio '80 ~ '90**  $\rightsquigarrow$  theory of **Curves of Maximal Slope** and **Minimizing Movements**
  - ▶ [*Gradient flows in metric spaces*, **Ambrosio-Gigli-Savaré** 2005]  $\rightsquigarrow$  systematic theory of existence, approximation & uniqueness of solutions of metric gradient flows, with applications to gradient flows in Wasserstein spaces.

## Metric analysis popping out

$$\begin{aligned} \int_{s_1}^{s_2} |\hat{u}'|_1 + \int_{s_1}^{s_2} \mathcal{M}_0(\hat{t}', \|\hat{u}'\|_2, -\partial\mathcal{E}(\hat{t}, \hat{u})) + \mathcal{E}(\hat{t}(s_2), \hat{u}(s_2)) \\ = \mathcal{E}(\hat{t}(s_1), \hat{u}(s_1)) + \int_{s_1}^{s_2} \partial_t \mathcal{E}(\hat{t}, \hat{u}) \hat{t}' \end{aligned}$$

**BUT:**

$B_1 \sim L^1$  does **not** have the **Radon-Nikodým property**

- ▶ Need to replace pointwise derivative  $|\hat{u}'|_1$  by **metric derivative**
- ▶ Metric analysis, theory of **gradient flows** in metric spaces:
  - ▶ **De Giorgi, Marino, Saccon, Tosques, Degiovanni, Ambrosio '80 ~ '90**  $\rightsquigarrow$  theory of **Curves of Maximal Slope** and **Minimizing Movements**
  - ▶ [*Gradient flows in metric spaces*, **Ambrosio-Gigli-Savaré** 2005]  $\rightsquigarrow$  systematic theory of existence, approximation & uniqueness of solutions of metric gradient flows, with applications to gradient flows in Wasserstein spaces.
- ▶ subdifferentials replaced by **lopes**, doubly nonlinear equations formulated in a metric setting [**R., Mielke, Savaré'08**], [**Mielke, R., Savaré'08**]

## Future developments

- ▶ applications of this general method to **"concrete" problems in continuum mechanics..**
- ▶ **from two norms to two metrics..**